

Review responses

Review response I

Referee comment:

- 5 The only major suggestion would be to include more information and interpretations of the data regarding environmental and climate changes that the lake has experienced during deposition of the sedimentary record. In Lines 54-55 it is specifically stated that the main aim of the projects is a better understanding of Holocene climate in Central Asia, so this should also be included in this paper. It is the only main weakness of the manuscript and the paper would have much more value if more details on the climate changes in the region are included as well.
- 10 That is why a third focus should be included in line 57, which states something like “3) reconstruction of regional climate changes in Central Asia”. The discussion of climate changes could partly be included into chapter 5.1 or within the last chapter of the discussion, to extend what the authors describe very briefly. This could either be added to chapter 5.4 or as a new chapter 5.5.

Author’s response:

- 15 We appreciate the constructive and very helpful comments on of anonymous referee #1 and addressed almost all suggestions in our revision. However, we relinquish from extending the paper to a more climatic focus. It is the explicit aim of this study to develop a robust chronology for the Chatyr Kul sediment record based predominantly on varve counting. This is the first varve chronology for a Central Asian lake sediment record and detailed analyses of seasonal sedimentation processes were required to prove the existence of annual laminations. This chronology
- 20 further enabled us through comparison with radiocarbon dating on aquatic matter for the first time (i) to quantify reservoir ages in a Central Asian lake record and (ii) to demonstrate changes of reservoir ages throughout the Holocene. Our results are of general interest for lake sediment dating in this region in settings where only radiocarbon dating is possible. Hence, we consider our chronological data as valuable stand-alone results and, therefore, have chosen *Geochronology* as a target journal. We explicitly disclaim from discussing climate changes
- 25 because this would have been far beyond the scope of this paper and, in contrast, even would have blurred its focus. We have clarified the focus of this paper in the manuscript (lines 51-54 in the revised manuscript).

Specific, minor suggestions:

Referee comment:

30 Lines 14-17. These first two sentences of the abstract appear to be repetitive and could be combined in shortened form.

Author's response:

corrected

Author's change in the manuscript:

35 Microfacies analysis of a sediment record from Lake Chatyr Kol (Kyrgyz Republic) reveals the presence of seasonal laminae (varves) from the sediment basis dated at $11,619 \pm 603$ years BP up to $\sim 360 \pm 40$ years BP. The Chatvd19 floating varve chronology relies on replicate varve counts on overlapping petrographic thin sections with an uncertainty of ± 5 %.

40 Referee comment:

Line 44: Typo – “n” to be removed

Author's response:

corrected

Author's change in the manuscript:

45 ...2) human influence $\#$ (Boomer et al., 2000; Mathis et al., 2014; Schröter et al., 2019 in review),...

Referee comment:

Lines 53-54: This information should be included into supplements. The aim of these projects is obviously the reconstruction of Holocene climate and so more information on this should be provided in the paper.

50 Author's response:

This was indeed misleading since we did not clearly distinguish between the overall project goals on climate reconstruction (which involves also other research groups) and the scope of this study, i.e. the construction of a robust age model for the entire project team. We have clarified the goal of this study in the revised version (lines 51-54)

55 Author's change in the manuscript:

The sediment record from Lake Chatyr Kol is the first varved record from CA covering most of the Holocene and the main goal of this study is to establish a robust age model through an integrated dating approach primarily based on varve counting. Varve counting requires an in-depth understanding of seasonal deposition of all varve types occurring in the sediment record.

60

Referee comment:

Line 62: remove dash

Author's response:

65 corrected

Author's change in the manuscript:

Geologically, the surrounding mountain ranges belong to Silurian –to Carboniferous sedimentary-volcanogenic complexes of marine-continental collision zones, consisting of limestones and dolomites, that crop out directly along the northern lake shore, as well as siliceous rocks, shales and scattered Permian granites that crop out in the south and north-east (Academy of Science of the Kyrgyz SSR, 1987).

70

Referee comment:

Line 70: delete amounts

Author's response:

75 corrected

Author's change in the manuscript:

Mean annual precipitation is ~275 mm/a as indicated by Aizen's (2001) evaluation and spatial averaging of annual precipitation ~~amounts~~ of historical records published by Hydrometeo (Reference Book of Climate USSR, Kyrgyz SSR, 1988).

80

Referee comment:

Line 133: How long was the in-growth time (check also spelling to change to “in-growth-time”. The term photo peak activity sounds incorrect to me and should be replaced with a more appropriate term. It is the gamma energy that is recorded in the gamma spectrometry.

85 Author’s response:

We changed “photo peak activities” to gamma energies and changed ingrowth-time to in-growth-time.

Author’s change in the manuscript:

After sufficient in-growth-time, the gamma energies of ^{210}Pb ($T_{1/2}= 22$ a) and ^{214}Pb ($T_{1/2}= 26.8$ min), which is a daughter nuclide of ^{222}Rn ($T_{1/2}= 3.8$ d), were measured at 46.54, 295.24 and 351.93 keV.

90 Hardware control, data storage, and spectrum analysis were realized with the software Genie 2000 (Canberra Industries). Measurements were taken out for 1.5 to 7 days (Suppl. Tab. 1).

Referee comment:

Line 139: lab-internal – please note which lab, and where these samples were analyzed.

95 Author’s response:

Information has been added.

Author’s change in the manuscript:

For this purpose, the Kryal[®] tubes were placed into two well-type germanium detectors G1 and G2 (Canberra Industries) located in a basement lab of a concrete building at GFZ Potsdam which is actively ventilated (Schettler et al., 2006).

100

Referee comment:

Line 167: use a, b, c to refer to each group of laminae more easily and use this instead of “LZ + number” in the references to Fig 4 throughout the text

Author’s response:

105 corrected

Author’s change in the manuscript:

Figure captions and figure 4 have been changed accordingly, also in the text.

Referee comment:

110 Line 204: Add current institute/university of Ms Schwarz within brackets as well

Author's response:

Information has been added.

Author's change in the manuscript:

115 The third sublayer is formed by diatom blooms exclusively consisting of the planktic diatom species Cyclotella choctawhatcheeana (pers. comm. Anja Schwarz, [TU Braunschweig](#)) (Fig. 4.1 b upper part).

Referee comment:

Line 231: Add picture of homogeneous sediment to Fig 4 as well to see how it compares to the varved intervals. In particular, this is useful to show the faint, discontinuous laminae in the uppermost cm

120 Author's response:

agreed.

Author's change in the manuscript:

Pictures of homogenous sediments with faint laminae have been added to figure 4.

125 Referee comment:

Lines 269-270: Is this assumption justified? +/- 40 years BP uncertainty could be higher or lower? Why is it not possible to more precise that this? On the basis of the data presented, I would be surprised if the error is as high as 40 year

Author's response:

130 We consider an uncertainty of +/- 40 years at the anchor point as justified for the non-varved interval given the counting uncertainty of ca 5% in laminated sections. We agree to the reviewer that this error might be over-

estimated but we prefer to provide a conservative estimate and clarified in the revised text that this uncertainty is considered a maximum range.

Author's change in the manuscript:

135 We assume a conservative uncertainty of ca. 10% as a maximum error for our interpolation.

Referee comment:

Line 386: change to effect

Author's response:

140 done.

Author's change in the manuscript:

The abrupt decrease of the reservoir [effect](#) after ~AD 1150, despite an increase in detrital carbonate supply (Sect. 5.3.5, Fig. 9) might be related to the silting up of the basin leading to a shallower water depth, which is more susceptible to water circulation and an enhanced atmospheric CO₂ exchange (c.f. Geyh et al., 1997).

145

Referee comment:

150 Line 491 (data availability statement): Please add the data into this database during the review process, so you can include the doi of the dataset in this statement. I think it is very important to add the doi to the final paper, so the future reader can access the datasets easily

Author's response:

done. We further added a link to the newly established VARDA database.

155 Author's change in the manuscript:

<https://doi.pangaea.de/10.1594/PANGAEA.909981>

<https://varve.gfz-potsdam.de>

160

165

170

175

Review response II

Referee comment:

180 Although I have no major comment on the central topic of this paper (that is suitable for the journal), i.e. the chronology, I am puzzled why there is no μ -XRF data (e.g. Itrax) shown in your study. For example, the authors describe periods of prevailing anoxic bottom water conditions, calcitic materials/diatoms, coarse vs finer sediments, etc. In my opinion, it would be very helpful to show μ -XRF elements (and elemental ratios) to support your visual microscopic analysis. Have you made such analysis (XRF)? If you are to interpret the paleoenvironments from this site in the paper, I think that would be very valuable.

185

Author's response:

We appreciate the constructive comments and suggestions of referee #2 and we now included XRF mapping to strengthen detailed microfacies analysis presented in our MS. The main focus of the MS is a detailed microfacies analysis for 1) the development of process-based deposition models and 2) the establishment of detailed varve-
190 based chronology. For these purposes, detailed microfacies analyses is crucial and allows to distinguish, for example, detrital from endogenic calcite and detrital quartz from diatom SiO_2 . Such differentiations are not possible using μ -XRF scanning as element abundances, as these sediment fractions are geochemically identical and it is not apparent from, for example, relative variations of calcium and silicon respectively. However, we do agree that XRF element scanning is a powerful way to complement and support visual microfacies observations. Therefore, XRF
195 scanning maps of sediment blocks are used, which are the equivalent of the thin sections used for the microfacies analyses. These new mapping results for selected intervals with characteristic varve types confirm the occurrence of both detrital and endogenic calcite. These data are presented in a new Figure 4.2 and in the main text (new chapters 3.3 and 4.4 μ XRF element mapping, discussion within chapters 5.4.2 and 5.4.3) of the manuscript. To our
200 opinion, XRF mapping results are most suitable for a precise linking of sediment compositions and microscopic observations.

We add Rik Tjallingii as a co-author because he conducted the μ XRF element mapping and helped with revising the manuscript.

Moderate comments:

205 Referee comment:

1. There is an excellent matching between the varve counts with the 2 dated wood samples. However, there is almost 6000 years (first~360 cm) without chronological constraint. Given that many varves are qualified as 'unclear' from 130 cm to~270 cm of the composite depth, perhaps some other dating techniques could be added such as paleomag, OSL, 14C, etc. I would encourage the authors to at least comment on this.

210 Author's response:

The reviewer is right that the age uncertainties are higher in this interval of less well preserved varves which we have addressed by allocating higher uncertainty ranges. Nevertheless, varves in this interval can still be counted and represent the only applicable dating method. We have sieved the entire interval in order to find terrestrial plant remains for radiocarbon dating but, unfortunately, without any success. All five ¹⁴C dates obtained from this interval are from aquatic material and thus revealed too old ages due to reservoir effects.

215

Independently paleosecular variation (PSV) records in this region are only available from Lake Issyk Kol, Lake Baikal and Lake Aslikul. However, these records suffer from dating problems and show significant temporal offsets before 500 AD between the records and to global geomagnetic field models (Gómez-Paccard, 2012) so that they cannot be used for the Chatyr Kol chronology.

220 OSL dating is also not applicable because the Chatyr Kol sediments are mainly composed of materials not suitable for reliable luminescence dating including carbonates, organics and non-aeolian siliciclastic.

Author's change in manuscript:

We have changed 'unclear' to 'less well preserved' varves (chapter 3.2).

225 Referee comment:

2. Have you used any particular software to count the varves, please provide what you used.

Author's response:

No software was used for varve counting. Counting was exclusively performed on the Axioplan microscope using different magnifications and based on expert knowledge.

230 Author's change in manuscript: none

Referee comment:

- 235 3. The names of the cores and their depth are indicated in Fig.3. However, it is unclear in my opinion which cores were used for the composite. I assume A1o, and some part of the A3o, A3u...In brief how much sediment was used from each core sections?

Author's response:

The core sections used for the composite profile are colored in grey, as indicated by the legend.

Author's change in manuscript:

240 We add the depth sections of each core used for the composite profile in fig. 3.

Referee comment:

4. Fig. 1: Have you obtained several (7) gravity cores that are not in the same location of the composite core?

Author's response:

245 As indicated in Figure 1, the gravity cores 3, 5, 6 and 7 were obtained close to the composite core location while gravity cores 1 and 2 were recovered about 1-1.5 km further north-east and number 4 was recovered ~ 10 km further east in the shallow eastern lake basin.

Author's change in manuscript: none

Referee comment:

- 250 5. Solar activity: Lines 414-416: Raspopov et al., (2008) use a 100-300 year band-pass filter and find 'great correlation' with solar activity (inferred from 14C) from three locations or so, and with lags (as high as 150 years). One can do the same analysis with white noise and find similar correlation (for example see Turner et al. 2016: Solar cycles or random processes?). But more importantly, they filter out (bandpass) the data which make any high correlation not surprising at all. The comparison of the tree-rings and 14C prod rate (Fig. 1; Raspopov et al., 2008) without filtering is not very convincing either. Finally, they don't
- 255

use the actual instrumental sunspots data spanning the past~300 years to compare with their tree-ring records, which is a little bit curious. To be honest, I don't reject the influence of solar forcing on regional climate, but based on this paper, it does not help your interpretation of the connection between solar forcing and your site.

260 5b: Lines 414: "which show decadal-to centennial periodicities". The authors refer to Fig. 4 LZ II. This is an image; hard to see any decadal-to centennial periodicities. Can you make spectral analysis of these layers characterizing lithozone II to prove these periodicities? It could be challenging without i.e. μ -XRF data.

265 Author's response:

We agree to the reviewer and delete the discussion on solar cycles. We restrain to a pure description of the observed intercalations and point out that it remains unclear if they are related to external triggers or random processes citing the reference suggested by the reviewer (Turner et al., 2016). We also change the term "periodicities" to intercalating /recurring patterns.

270 5b. As we agree to the reviewer concerning the occurrence of petrographic periodicities and avoid this term in the revised manuscript, there is no further need for spectral analyses. The deposition of differing varve types showing the decadal-centennial intercalating pattern is purely based on varve counting. To better visualize the intercalations we add a figure linking sediment compositions and microfacies observations using XRF mapping of the varve type distribution for selected intervals of LZ II and LZ III (new Fig. 4.2 and 4.3).

275

Author's change in manuscript:

The causes for these clear intercalations remain speculative and include either external (climatic) triggers or unknown lake-internal or sedimentation variability (Turner et al., 2016 and reference herein). Fig. 4.2 (μ XRF mapping) has been added.

280

Referee comment:

6. In the text the authors use AD, please add AD/BC in your plots.

Author's response:

agreed.

285 Author's change in manuscript: An axis of AD/BC ages is added to the plots.

Minor comments:

Referee comment:

Lines 37-38: Why Lake Telmen is varved~1940-2013? Human influence (N & P) in the watershed? If so, this is not the case for your site?

290 Author's response:

This was misunderstood by the reviewer because the varve record from 1940 – 2013 is from Lake Sary Chelek in Kyrgyzstan and not from Lake Telmen. From Lake Telmen discontinuous varved intervals are reported for the time period from 4,390 cal years BP on (Peck, 2002). We clarify the sentences about other regional varve records in the text.

295 Author's change in manuscript:

In Kyrgyzstan, varves have so far been only reported from Lake Sary Chelek (Kyrgyzstan) for the short time interval from ~1940's to 2013 (Lauterbach et al., 2019). Other varved records in the larger region are Lake Telmen in northern Mongolia which exhibits discontinuously varved intervals from approximately 4,390 cal years BP (Peck, 2002) and Lake Sugan in north western China covering the last ~2,670 years BP (Zhou et al., 2007).

300

Referee comment:

Figures 5 and 6: add error bars on CRS/CIC model

Author's response:

305 done.

Author's change in manuscript:

Error bars added in Figures 5 & 6.

Referee comment:

310 Lines : 164-233-763 : change centimetre to centimeter

Author's response:

agreed.

Author's change in manuscript:

Line 164: The uppermost centimeter is enriched in calcite and exhibits greyish faint laminations.

315 Line 233: Faint and discontinuous calcite laminae occur in the uppermost centimeter (Fig. 4.1 f).

Line: 763: (Fig.6) Core pictures of the upper part of the composite profile CHAT12 (right) and the gravity core SC17_7 (left) illustrate the facies change to calcite-enriched sediments in the uppermost centimeter.

Referee comment:

320 Figure 1: should add labelling to isobaths.

Author's response:

agreed.

Author's change in manuscript:

Isobaths are labelled in figure 1.

325

Referee comment:

Line 301: laminar denudation: please describe this.

Author's response:

330 By "laminar denudation" we mean the superficial catchment runoff probably associated with an activation of widely dispersed smaller tributaries during precipitation events. We will clarify this in the text.

Author's change in manuscript:

Runoff with suspended sediment load is then likely directed through the Kegagyr River in the east but may also be the result of surface runoff through the activation of several and widely distributed smaller tributaries in the catchment.

335

Referee comment:

Line 461: Why such an increase of precipitation at AD 1150? MCA? However, it seems to last until recent, so occurring in the LIA as well. A change in boundary conditions in the watershed? High-resolution grain-size analysis could shed some light about this.

340

Author's response:

It is correct that the onset of additional detrital sub-layers during summer started at AD 1150 but it lasted not until recent but until ca. AD 1730, about the time when varve preservation became poor and finally ceased. We do not know the reason for this increase in summer runoff events and can only state that there is no coincidence with climatic periods reported from other records (MCA, LIA). We did not carry out grain size analysis because our continuous microfacies analyses does not reveal any significant shift in grain size. Therefore, changes in boundary conditions in the catchment appear unlikely. We have revised and clarified the text accordingly.

345

Author's change in the manuscript:

Chapter 5.4.5 revised

Clastic-organic varves constitute 59 % of the observed varves in LZ V, clastic-calcitic varves 26 % and organic-clastic varves 15 %, the latter ceasing at 110.5 cm (AD 1260 ± 50). Varve microfacies changes abruptly at 130 cm depth or AD 1150 from the dominance of organic-clastic varves to dominating clastic-organic and clastic-calcitic varves. Within 5 years, varve thickness drastically increase from Ø 0.43 mm in LZ IV to Ø 1.52 mm in LZ V due to thicker summer sublayers. Thicker summer sublayers result from both thicker mixed sublayers rich in algae remains (*Botryococcus*, chrysophytes, diatoms) and additional late summer detrital sublayers (Fig. 4.1.e, Suppl. Fig. 2f). The increase in summer layer thickness, therefore, suggest both, higher lacustrine productivity and an increase in summer runoff. However, the reasons for these changes remain elusive and a relation to known climatic

355

360 periods like the Medieval Climate Anomaly and the Little Ice Age is not found. One might speculate that the
frequent occurrence of late summer runoff layers either reflects convective rainfall events due to recycling of local
moisture sources (Aizen et al., 2001), or changing atmospheric circulation regimes. Changes in boundary conditions
in the catchment of the lake are unlikely since microfacies analyses does not show pronounced changes in grain
size distribution of the detrital material. Human impact cannot fully be excluded but low indices of human and
livestock fecal biomarkers (Schroeter et al., 2020) are an argument against major human impact. The presence of
lake deposits at the northern and southern shores ca 1.5 - 1 m above present day lake level dated at AD 1420 ± 204
365 cal years BP, AD 1044 ± 160 and AD 858± 166 (Shnitnikov, 1978) suggests that increased summer runoff might
have resulted in a more positive water budget and lake level rise.

References

370 Gómez-Paccard, M., Larrasoaña, J. C., Giralt, S., & Roberts, A. P. (2012). First paleomagnetic results of mid-to
late Holocene sediments from Lake Issyk-Kul (Kyrgyzstan): Implications for paleosecular variation in central Asia.
Geochemistry, Geophysics, Geosystems, 13(3).

Schroeter, N., Lauterbach, S., Stebich, M., Kalanke, J., Mingram, J., Yildiz, C., .Shouten, S. & Gleixner, G. (2020).
Biomolecular evidence of early human occupation of a high-altitude site in Western Central Asia during the
375 Holocene. *Frontiers in Earth Science*, 8, 20. <https://doi.org/10.3389/feart.2020.00020>

380

385

Review response III:

We thank reviewer #3 for his appreciation of our work and his valuable suggestion that we include in our revision.

Referee comment:

- 390 1. There are several points in Section 5 (eg. Line 298; Line 401; Line 435) that refer to the role of glacial meltwater in the supply of detrital clastic material into the basin and their presence in the varve microfacies. Is it possible to include the area that is/was glaciated in Figure 1? There is no mention of this in Section 2 (Study Site) other than meltwater run-off and it is not clear if this is from a glaciated catchment. A little more detail on this would be helpful to the reader. Also permafrost thaw is considered a possible contributor to older carbon in the lake water to explain the reservoir but this is not described within the site context.
- 395 Could this also be included in the site context?

Author's response:

The area displayed in figure 1 does not exhibit recent glaciers, but several glaciers exist further north-east on the central At Bashy range (Narama et al., 2007) at a level above ~4000 m a.s.l. and some of them drain into the Chatyr Kol via the Kegagyr River. Recent glaciers are also located on the western Torugat range (not shown in Fig.1) but 400 these glaciers do not drain into Chatyr Kol. We clarify glacier information in the chapter "study site". We further extend the formulation 'glacier and snow meltwater' because runoff includes also seasonal snowmelt. Finally, a photo of field observations displaying thermokarst/permafrost thawing structures observed at the Maloye Lake in < 2 km distance from Chatyr Kol is added.

Author change in manuscript:

405 Line 65-68:

The modern lake, which has a maximum length of 23 km, a width of 10 km and a maximum depth of 20 m in its western-central part, is endorheic and separated from the neighboring Arpa river basin in the north-west by a moraine (Shnitnikov, 1978). *The moraine originated from glacial advances of unknown age from the western Torugat range. Present day glaciers on the Torugat and At Bashy range exist above ~4000 m a.s.l. but only some 410 of the At Bashy glaciers drain into Chatyr Kol via the Kegagyr River.* The lake is moreover fed by convective rainfall events in summer (Aizen et al., 2001). *A shallow watershed hinders outflow to the east.*

Line 83-84

The permafrost level is located at a depth of 2.5-3 m in the littoral coast zones and the lake is covered by ice from October to April (Shnitnikov, 1978). Modern permafrost thawing results in instable shores visible at the Maloye lake located < 2 km to the South of Chatyr Kol(Fig. 1 Photo) and the development of small ponds on the shallow south-western shore of this lake and lake Chatyr Kol during the summer season.

Referee comment:

2. Microfacies section – the introduction to section 4.2 might be considered contradictory in that Line 166 states ‘consists of mainly clastic lamination’ but Figure 4 has clastic material in all of the microfacies. Later in the paragraph, it is stated that ‘sub-types were named according to the order of their dominant contents’, which has two occasions where either organic or calcitic laminations dominate the microfacies making the earlier statement invalid. If the first sentence said ‘Clastic material is present in all of the macroscopically visible laminations below 63.0 cm depth, and intercalates with calcitic, aragonitic and organic sublayers that build-up cyclic successions. And then the final three sentences can remain and it is a truer reflection of the microfacies. Also it could be useful to state how the subtypes are named according to the their dominant contents (I assume that the dominant component comes first?).

Author’s response:

The referee is right that our formulation was not sufficiently clear and we will change the introduction to chapter 4.2 accordingly. We only want to point out, that (i) all six varve types include a clastic sublayer and that (ii) the differentiation of varve types relies on the composition of the alternating sublayers and the dominance of sublayers within a varve cycle. The latter criterium is used for varve type names. For example, the clastic-organic varve type is characterized by the dominance of the clastic sublayer, while in the organic-clastic varve type the organic sublayer prevails. The varve types names are not related to the order of sublayer succession within the varves. In addition to the revision of the introduction of chapter 4.2, we re-name the subchapters from ‘Clastic-organic laminae’ to Clastic-organic type’ etc. to clarify that we are not presenting individual sublayers but varve types.

Author change in manuscript:

Line 166:

440 Microscopic sediment analysis revealed, that clastic sublayers are present throughout the finely laminated
sediments below 63.0 cm depth (Fig. 4). These clastic sublayers are variably intercalated with calcitic, aragonitic
and organic sublayers and thus form different types of cyclic successions. In total, we classified six different types
of sublayer successions as described below. The name for these types reflects the dominant sublayer for each of
the six types. For example, the 'clastic-organic type' is characterized by the dominance of clastic sublayers, while
in the organic-clastic type organic sublayers dominate. The names are not related to the order of sublayer succession
445 within each type.

Referee comment:

3. Figure 4 is good at showing the broader differences in the microfacies in each of the LZ's. However, the
detail in the schematic (varve depositional model) is difficult to evaluate within the images from the thin
450 sections at their current magnification. Could a higher magnification image that reflected more closely the
elements shown in the schematic also be included? Also a key for the symbols in the schematic is necessary,
and I note that there is no obvious winter layer detected in the clastic-diatom and clastic organic/clastic
aragonitic microfacies. Related to this, in the text is 'section 4.2.1 Clastic-organic laminae' the lower
schematic or the upper schematic? What is the difference between these two? It appears to be the aragonite
455 and this is what is identified in the text, but the clastic-organic coming first in the Figure confuses this
distinction. It would also be helpful that the order that is in the text was followed by the order in the figure
to remove this confusion.

Author's response:

We include additional microscopic images at higher magnification in the supplement. (Suppl. Fig. 2) In addition,
460 we include μ XRF element mapping of selected thin sections in order to better visualize the varve facies (Fig. 10).

Keys/Legends will be added for the used symbols in figure 4.

Winter layers in clastic-organic/clastic-aragonitic varve deposition model are added in fig. 4.

The reviewer is also right that the relation of schematics in figure 4 to section 4.2.1 is not clear. Therefore, we
modified fig. 4 accordingly.

465 Author change in manuscript:

We will change the order of the sub chapters in 4.2 according to figure 4 and revise the varve schemata label in figure 4.

Referee comment:

- 470 4. Section 5.4 provides a nice explanation of the broad environmental changes that lead to variations in the microfacies through the sequence. A criticism is that it is difficult to evaluate the thickness data of the different microfacies against the text, which starts by describing the frequency of the different microfacies in each of the Lithozones. A suggestion that could help the reader and also highlight the differences in microfacies that are observed between the LZ's would be to include on Figure 9 some percentage bar charts that collate the relative proportion of the different microfacies in each LZ. Such that for LZ I with clas-org 475 57%, clas-calc 29% and clas-arag 14%, clas-dia 0%,org-clas 0% calc-clas 0%. Then using the same order for the microfacies there could be a bar chart for LZ II, LZ III etc and then if aligned vertically the reader could draw a direct comparison between LZ's seeing the changes through the sequence. This could be a column on the right hand side of the current Figure. It would also be useful to arrange the thickness graphs 480 for each of the microfacies in the same order as their description in Figure 4 and in the text of Section 4.

Author's response:

The referee is right and changes in figure 9 have been made.

Author change in manuscript:

485 Instead of using bar plots, pie charts are displayed for each lithozone in figure 9. The thickness graphs in figure 9 are ordered according to the order in the text in chapter 4.2.

Technical corrections:

Referee comment:

490 Throughout the manuscript superscript is used inconsistently when it should be used e.g. for ^{14}C . Lead-210 is used interchangeably with ^{210}Pb , and cm-1 should be cm⁻¹. Spaces should be included between ages and the \pm symbol.

Author's response:

agreed. The text will be changed accordingly.

Referee comment:

495 Line 39 – states ‘.....which cover approximately 7,100 cal years BP....’ , is that the duration of the record or the base of the sequence is dated using varves to 7,100 cal years BP. Is this also the case for Lake Sugan and is this also in cal yrs BP?

Author’s response:

The lake sediment record of Lake Telmen extends back to 7,100 cal years BP according to AMS radiocarbon dating
500 of bulk sediment and pollen extracts. Varves in the Telmen record were only found in sediments younger than 4,390 cal yr BP (AMS) and have not been counted because varve preservation is reported as discontinuous. Instead, Peck et al. (2012) extrapolated average couplet thickness for an indirect varve age estimate.

For Lake Sugan, the authors counted laminae couplets on digital images of split core surfaces and compared these with ²¹⁰Pb dating (CRS). The ages were not provided as cal yrs BP.

505 Author change in manuscript:

In [Kyrgystan](#), varves have been only reported from Lake Sary Chelek [for the short time interval](#) from ~1940’s to 2013 (Lauterbach et al., 2019). [Other varved records in the wider region](#) are Lake Telmen in northern Mongolia which [exhibits discontinuous varved intervals during the last ca. 4,390 cal years BP](#) (Peck, 2002) and Lake Sugan in north western China covering the last ~2,670 years BP (Zhou et al., 2007).

510

Referee comment:

Line 44 – remove ‘n’

Author’s response:

Will be removed.

515

Referee comment:

Line 92 – where they are archived in a cold store at 4°C

Author's response: agreed

Author change in manuscript:

520 All cores were opened, split and photographed at GFZ Potsdam, where they are [archived in a cold store at 4°C](#).

Referee comment:

Line 97 – remove ‘continuously’ and put ‘Continuous’ at the start of the sentence.

Author's response: agreed

525 Author change in manuscript:

Continuous 10-cm-long sediment slabs with an overlap of 2 cm were taken from the whole composite profile to prepare large-scale petrographic thin sections.

Referee comment:

530 Line 138- should a value for keV be included after 5.9%?

Author's response: The keV for ^{210}Pb and ^{137}Cs were mentioned already earlier in the text in line 134-135.

Author change in manuscript: none

Referee comment:

535 Line 175 – is there an image that illustrates how it is possible to distinguish between detrital and endogenic calcite?

Author's response: agreed

Author change in manuscript:

An image (Fig. 4.2) (microscopic pictures) illustrating the differences is added to the manuscript.

540 Referee comment:

Line 301- I was unclear on ‘laminar denudation’ is that erosion of the lamination?

Author's response:

545 By “laminar denudation” we mean the superficial runoff associated with catchment runoff through the activation of widely dispersed smaller tributaries. We changed the wording from “laminar denudation” to “surface runoff” and clarify the meaning.

Author change in manuscript:

Runoff with suspended sediment load is then likely directed through the Kegagy River in the east but may also be the result of surface runoff through the activation of several widely distributed smaller tributaries in the catchment.

550 Referee comment:

Line 311 – replace ‘overserved’ with ‘observed’?

Author’s response: agreed.

Author change in manuscript:

555 Aragonite precipitates were only observed in the intervals between 600.0-605.0 and 609.0-616.0 cm composite depth.

Referee comment:

560 Line 343-344 – I was not clear on the meaning of ‘for each individual thin section comprising 324 and 13 years varve. My assumption is that this is the range, or maximum and minimum, in total number of varves observed on a single 10 cm thin section. However, I may have misread this.

Author’s response: This is correct, this refers to maximum and minimum number of varves within individual thin sections.

Author change in manuscript:

565 For the floating varve chronology we therefore compare the results for each individual thin section comprising between a maximum of 324 (506.8-497.6 cm) and a minimum of 13 (varves) (65.4-63.0 cm) (Fig. 5a, Fig. 8).

Referee comment:

Line 474 – unclear on the meaning of ‘robust fundament’. Do you mean ‘This robust chronology is fundamental for further detailed palaeoenvironmental.....’?

570 Author’s response: Yes.

Author change in manuscript:

This **robust** chronology **forms the base** for further detailed palaeoenvironmental and palaeoclimatic reconstructions.

Referee comment:

575 Line 479 – I assume that the increased windiness enable increased mixing of the lake waters and CO₂ exchange with the atmosphere. Perhaps be explicit here.

Author's response: agreed.

Author change in manuscript:

580 Lower reservoir ages of ~1000 years and less in the late Holocene might be related to enhanced atmospheric CO₂ exchange when the lake was shallower due to silting-up of the lake basin and/or increased windiness **inducing increased water column mixing favoring CO₂ exchange with the atmosphere.**

Referee comment:

Line 480 replace 'which allowed developing' with....'that allowed the development of seasonal deposition models

585 Author's response: agreed.

Author change in manuscript:

The construction of the varve-based chronology was only possible through detailed micro-facies analyses of the entire sediment sequence in overlapping thin sections **that allowed the development of** seasonal deposition models for all observed types of fine laminations.

590

References

Narama, C., Kääh, A., Duishonakunov, M., & Abdrakhmatov, K. (2010). Spatial variability of recent glacier area changes in the Tien Shan Mountains, Central Asia, using Corona (~ 1970), Landsat (~ 2000), and ALOS (~ 2007) satellite data. *Global and Planetary Change*, 71(1-2), 42-54.

595

List of relevant changes in the manuscript

- 600 1. Text changes have been made according to the suggestions of the three reviews in (deleted words etc. are not mentioned), Lines of the revised manuscript are displayed:
- Lines 14-17: text changes according to review I
 - Line 36-40: text changes according to review I, II and III
 - Line 49-54: text changes according to review I and III
 - 605 • Line 63-68: text changes according to review III
 - Line 84-86: text changes according to review III
 - Line 94: text changes according to review III
 - Line 99: text changes according to review III
 - Line 108: text changes according to review II
 - 610 • Line 146,148, 150: text changes according to review I
 - Line 176, 248, 885: text changes according to review II
 - Line 178-183: text changes according to review III
 - Line 224: text changes according to review I
 - Line 281: text changes according to review III (technical correction)
 - 615 • Line 296: text changes according to review I
 - Line 329-330: text changes according to review II and III
 - Line 380-381: text changes according to review III
 - Line 423: text changes according to review I
 - Line 450-451: text changes according to review II
 - 620 • Line 500-509: text changes according to review II
 - Line 523-525: text changes according to review III
 - Line 527: text changes according to review III
 - Line 542-543: text changes according to review I (inclusion of data availability doi)

 - 625 • Line 124-131, 247, 249-261, 284, 287, 312-321, 336, 338-352, 355, 443-444, 451-453, 466-473, 535-538, 549-550, 552-560: minor text changes after own revision and implementation of XRF mapping results
- 630 2. Changes in the order of chapters (due to the inclusion of additional chapters following the suggestions of review II and also following review III)
- Line 3: Rik Tjallingii was added to the list of Co-Authors due to the inclusion of XRF element maps
 - new chapters: 3.3 & 4.3 XRF element mapping changes order of subchapters within chapter 3 and 4
 - changed order of subchapters of chapter 4.2 according to the appearance of microscopic pictures in figure 4 (review III)

635

3. Changes of Figures:

- Figure 1: Isobath were labelled (review II) and one photo showing permafrost thawing structures was included (review III)
- Figure 3: Onset and basis depths of used composite sections are displayed (review II)
- 640 • Figure 4.1 Keys/Legend has been added, microscopic pictures of homogenous sediments have been added (review III)
- New figure 4.2 and 4.3 displaying XRF element mapping results (review II)
- Figure 5 AD/BC axis has been added, Errors have been added to CIC and CRS models (review II)
- Figure 6 Errors have been added to CIC and CRS models (review II)
- 645 • Figure 7 AD/BC axis has been added
- Figure 9 Varve types have been colored and their order of appearance follows figure 4.1 and the order of subchapters 4, pie charts displaying varve type percentages per lithozone have been added (review III)

650 4. Added references:

- Line: 709-710: Shapley, M. D., Ito, E., & Donovan, J. J.: Authigenic calcium carbonate flux in groundwater-controlled lakes: implications for lacustrine paleoclimate records. *Geochimica et Cosmochimica Acta*, 69(10), 2517-2533, <https://doi.org/10.1016/j.gca.2004.12.001>, 2005
- 655 • Turner, T. E., Swindles, G. T., Charman, D. J., Langdon, P. G., Morris, P. J., Booth, R. K., Parry, L. E., and Nichols, J. E.: Solar cycles or random processes? Evaluating solar variability in Holocene climate records, *Sci Rep*, 6, 23961, <https://doi.org/10.1038/srep23961>, 2016. (review II)

5. Changes of supplement:

- New supplementary figure 2 showing more details of individual varve structures (review III)
- 660 • Minor changes of Suppl. Table. 2 : inclusion of age errors, correction of depth 6.25 (SC17_7)
- Minor changes of Suppl. Table 3: following changes of Suppl. Table 2

665

Revised manuscript (revisions in blue):

Seasonal deposition processes and chronology of a varved Holocene lake sediment record from Lake Chatyr Kol (Kyrgyz Republic)

Julia Kalanke¹, Jens Mingram¹, Stefan Lauterbach², Ryskul Usubaliev³, Rik Tjallingii¹, Achim Brauer^{1,4}

¹GFZ German Research Centre for Geosciences, Section 'Climate Dynamics and Landscape Evolution', Potsdam, Germany

5 ²University of Kiel, Leibniz Laboratory for Radiometric Dating and Stable Isotope Research, Kiel, Germany

³Central Asian Institute for Applied Geoscience, Bishkek, Kyrgyzstan

⁴University of Potsdam, Institute of Geoscience, Potsdam, Germany

Correspondence to: Julia Kalanke (juliak@gfz-potsdam.de)
10 Telegrafenberg C322
14473 Potsdam/Germany
+49 331 288-1379

Abstract.

15 Microfacies analysis of a sediment record from Lake Chatyr Kol (Kyrgyz Republic) reveals the presence of seasonal laminae (varves) from the sediment basis dated at $11,619 \pm 603$ years BP up to $\sim 360 \pm 40$ years BP. The Chatyrd19 floating varve chronology relies on replicate varve counts on overlapping petrographic thin sections with an uncertainty of $\pm 5\%$.

The uppermost non-varved interval was chronologically constrained by ^{210}Pb and ^{137}Cs γ -spectrometry and interpolation based on varve thickness measurements of adjacent varved intervals with an assumed maximum uncertainty of 10%. Six varve types were distinguished, are described in detail and show a changing predominance of clastic-organic, clastic-calcitic or -aragonitic, 20 calcitic-clastic, organic-clastic and clastic-diatom varves throughout the Holocene. Variations in varve thickness and the number and composition of seasonal sublayers are attributed to 1) changes in the amount of summer or winter/spring precipitation affecting local runoff and erosion and/or to 2) evaporative conditions during summer. Radiocarbon dating of bulk organic matter, daphnia remains, aquatic plant remains and *Ruppia maritima* seeds reveal reservoir ages with a clear decreasing trend up core from $\sim 6,150$ years in the early Holocene, to $\sim 3,000$ years in the mid-Holocene, to $\sim 1,000$ years and less in the 25 late Holocene and modern times. In contrast, two radiocarbon dates from terrestrial plant remains are in good agreement with the varve-based chronology.

1 Introduction

30 The interplay of the large atmospheric circulation systems in Central Asia (CA), including the Siberian High, the Westerlies and the Indian Monsoon and their influences on regional climate is still not fully understood. This is partly due to the large contrasts of landscapes (high mountains, deep basins, large water bodies and deserts), the low spatial and temporal coverage of high-resolution paleo-climate archives and the partly problematic dating of these archives in this area. Information about Holocene climate variability in CA derive from several types of archives, including tree rings (Esper et al., 2003), speleothems

(Fohlmeister et al., 2017; Wolff et al., 2017), ice cores (Aizen, 2004), aeolian deposits (Huayu et al., 2010) and lakes (Heinecke et al., 2017; Lauterbach et al., 2014; Mathis et al., 2014; Rasmussen et al., 2000; Ricketts et al., 2001; Schwarz et al., 2017). However, none of these lake records has reported annually laminated sediments. In Kyrgyzstan, varves have been only reported from Lake Sary Chelek for the short time interval from ~1940's to 2013 (Lauterbach et al., 2019). Other varved records in the wider region are from Lake Telmen in northern Mongolia which includes discontinuous varved intervals during the last ca. 4,390 cal years BP (Peck, 2002) and from Lake Sуган in north western China covering the last ~2,670 years BP (Zhou et al., 2007). Deciphering Holocene climate changes based on limnic records in CA is challenging due to the influences of several factors: 1) chronological uncertainties caused by the scarcity of datable terrestrial plant material at high altitudes and often large ¹⁴C reservoir effects of aquatic organic material (Hou et al., 2012; Lockot et al., 2016; Mischke et al., 2013), 2) human influence (Boomer et al., 2000; Mathis et al., 2014), possibly overprinting the natural climate signals in the archives and 3) variations in the dominance of the mid-latitude Westerlies, the Siberian High and the Asian Monsoon system leading to different spatial and temporal climate effects over CA (Chen et al., 2008; Herzsuh, 2006; Mischke et al., 2017; Schroeter et al., 2020). All these factors can hamper data comparison and may lead to different paleo-environmental interpretations (Chen et al., 2008; Hou et al., 2012; Mischke et al., 2017). The investigation of varved lake sediments offers the unique opportunity for independent dating through varve counting. In addition, the description of varve micro-facies has the potential to provide detailed insights into environmental and climate variations at a seasonal scale. The sediment record from Lake Chatyr Kol is the first varved record from CA covering most of the Holocene and the main goal of this study is to establish a robust age model through an integrated dating approach primarily based on varve counting. Varve counting requires an in-depth understanding of seasonal deposition of all varve types occurring in the sediment record. Therefore, we apply continuous microfacies analysis for the entire sediment profile to describe the Holocene evolution of varve formation in detail and discuss fundamental deposition processes.

55

2. Study site

Lake Chatyr Kol (40°36' N, 75°14' E) (Fig.1) is located at ~3,530 m above sea level (a.s.l.) in the intramontane Aksai Basin (De Grave et al., 2011; Koppes et al., 2008) in the southern Kyrgyz Republic. In the north, the basin is restricted by the At Bashy Range and in the south by the Torugat Range resulting in a catchment area of about 1,084 km². Geologically, the surrounding mountain ranges belong to Silurian to Carboniferous sedimentary-volcanogenic complexes of marine-continental collision zones, consisting of limestones and dolomites, that crop out directly along the northern lake shore, as well as siliceous rocks, shales and scattered Permian granites that crop out in the south and north-east (Academy of Science of the Kyrgyz SSR, 1987). The modern lake, which has a maximum length of 23 km, a width of 10 km and a maximum depth of 20 m in its western-central part, is endorheic and separated from the neighbouring Arpa river basin in the north-west by a moraine (Shnitnikov, 1978). The moraine originates from glacial advances of unknown age from the western Torugat mountain range.

65

Present day glaciers exist above ~4000 m a.s.l. on the Torugat and At Bashy mountain ranges but only some of the At Bashy glaciers drain into Chatyr Kol via the Kegagyr River. The lake further receives convective rainfall in summer (Aizen et al., 2001). A shallow dam at ~ 3,550 m a.s.l. hinders outflow to the east. Modern climate conditions are generally dry and mainly controlled by the Westerlies and the Siberian Anticyclone Circulation (Aizen et al., 2001; Koppes et al., 2008). Mean annual precipitation is ~275 mm/a as indicated by Aizen's (2001) evaluation and spatial averaging of annual precipitation of historical records published by Hydrometeo (Reference Book of Climate USSR, Kyrgyz SSR, 1988). This is comparable to long-term instrumental data from nearby (about 50 km away) weather stations at comparable altitudes, where annual precipitation means are 237 mm/a (station "Chatirkul", 75°8'E, 40°6'N, 3,540 m a.s.l, AD 1961–1990) and 294 mm/a (station "TienShan", 78°2'E, 41°9'N, 3,614 m a.s.l., AD 1930–2000) (Williams and Konovalov, 2008). Monthly mean temperatures range from -26.0 to 8.0 °C (Koppes et al., 2008; Academy of Science of the Kyrgyz SSR, 1987) with means of -5.4 °C (station "Chatirkul") and -7.6 °C (station "TienShan") (Williams and Konovalov, 2008). The salinity of the lake water ranges from 1.06–1.15 g/l in the deeper western part of the lake to 0.24 g/l in the shallower eastern part near the inflow (Romanovsky, 2007). Measurements of oxygen concentrations (YSI Pro 6600 V2) during a field trip in July 2012 ranged from ~6 mg/l at the water surface to ~1 mg/l in 19 m depth with a clear oxygen minimum zone below ~11 m depth (Fig. 2). Water surface and bottom water temperatures (YSI Castaway CTD) at 19 m depth reached 13.2 °C and 9.4 °C. Specific conductivity (CTD) ranged from 1,902 $\mu\text{S}/\text{cm}^{-1}$ at 19 m to 1,825 $\mu\text{S}/\text{cm}^{-1}$, pH was ~9 (YSI Pro 6600 V2). Secchi depth was about 4 m. Nowadays, the lake is largely occupied by the amphipod *Gammarus alius* sp. nov. (Sidorov, 2012) and no fish live in the lake (Shnitnikov, 1978). The permafrost level is located at a depth of 2.5-3 m in the littoral coast zones and the lake is covered by ice from October to April (Shnitnikov, 1978). Modern permafrost thawing results in instable shores visible at the Maloye lake located < 2 km to the South of Chatyr Kol (Fig. 1 Photo) and the development of small ponds on the shallow south-western shore of this lake and lake Chatyr Kol during summer. Several terraces in the north, south and east of the lake result from Pleistocene-Holocene lake level fluctuations (Romanovsky and Shatravain, 2007; Shnitnikov, 1978). Vegetation around the lake is generally poor and represented by high alpine meadows (Shnitnikov, 1978; Taft et al., 2011).

90 3. Methods

3.1 Coring and Composite profile

Five parallel cores each of 3 to 6 m length have been retrieved in 2012 from the deepest part of the lake (40°36.37' N, 75°14.02' E) by using an UWITEC piston corer (Fig. 1, Tab. 1). All cores were opened, split and photographed at GFZ Potsdam, where they are archived in a cold store at 4°C. A continuous composite profile of 623.5 cm length (CHAT12) was established by correlating the individual, overlapping cores via macroscopically visible marker layers (Fig. 3). Furthermore, seven parallel gravity cores (SC17_1-7) have been retrieved with a UWITEC gravity corer in 2017 (Fig.1, Tab.1) to recover the undisturbed sediment-water interface, from which the best preserved parallel core SC17_7 was used for gamma spectrometric analysis.

3.2 Sediment microfacies analysis and varve counting

100 **Continuous** 10-cm-long sediment slabs with an overlap of 2 cm were taken from the whole composite profile to prepare large-scale petrographic thin sections. Thin section preparation followed the method described by Brauer and Casanova (2001) and included freeze-drying and vacuum impregnation of the sediment slabs with Araldite epoxy resin. Microfacies analysis, including a semi-quantitative evaluation of planktic and periphytic diatoms, aquatic plant remains (e.g. *Potamogeton* sp., *Ruppia maritima*), ostracods, daphnia, characeae and chrysophytes, was carried out on a Zeiss Axioplan microscope using different magnifications (25–400 x) and included measurement of varve thicknesses, microfacies/varve type characterization, 105 the definition of varve boundaries and the development of process-related deposition models. A varve quality index (VQI) ranging from 0-5 was given for each varve, comparable to the method from (Żarczyński et al., 2018) and references therein.

- VQI 0 = no varves or strongly disturbed varved sequences, no reliable counting (interpolation)
- VQI 1 = very low varve preservation, horizontally discontinuous varve and **less well-preserved** sublayer boundaries, difficult counting
- 110 • VQI 2 = low varve preservation, occasional horizontally discontinuous varve and sublayer boundaries, reliable counting
- VQI 3= medium varve preservation, horizontally continuous varve and sublayer boundaries, only small disturbances, reliable counting
- VQI 4= high varve preservation, clearly distinguishable varve and sublayer boundaries, reliable counting
- 115 • VQI 5= highest varve preservation, clearly distinguishable varve and sublayer boundaries, no disturbances, reliable counting

Varve counting was performed to establish a floating varve chronology. Non-varved intervals (VQI=0) between varved sediment sections were therefore interpolated by using the mean of sedimentation rates derived from about 20 varves above and below the non-varved part. Varves were counted twice by the same author. Counting uncertainty estimates were first 120 assessed by the percentage deviation of the second to the first count within one thin section. The mean of these deviations was used as an overall counting uncertainty estimate and assigned to the entire varved record. The uncertainty estimates were thus also assigned to interpolated sequences.

3.3 XRF element mapping

125 **X-ray Fluorescence (XRF) element mapping** was performed on two selected Araldite impregnated sediment blocks (ca. 2 x 10 cm), which were prepared for thin sections used for the microfacies analyses. XRF element mapping of these two sediment blocks allows linking micro-facies analyses of typical varve types directly with geochemical sediment compositions. Element mapping was performed at 50 µm resolution and covering most of the surface of the sediment block (15 x 100 mm) using a Bruker M4 Tornado at GFZ Potsdam. This scanner is equipped with a Rh X-ray source operated at 50 kV and 600 mA in 130 combination with poly-capillary X-ray optics that irradiate a spot of 20 µm for 50 ms. After measuring and an initial spectrum deconvolution, normalized element intensities are used to visualize relative element abundances as 2D maps.

135 3.4 Radiometric dating

3.4.1 Radiocarbon dating

In total, 36 accelerator mass spectrometry (AMS) ^{14}C measurements were carried out at the Poznań Radiocarbon Laboratory in Poland. Samples for ^{14}C measurements comprised two pieces of wood, bulk TOC samples, aquatic plant macro remains, daphnia remains and *Ruppia maritima* seeds (Tab.2). Additional samples of recent living daphnia and aquatic plants have been
140 collected to assess the modern ^{14}C reservoir effect. The resulting conventional ^{14}C ages were calibrated using OxCal 4.3 (Ramsey, 2009) with the IntCal13 calibration curve (Reimer et al., 2013).

3.4.2 Gamma spectrometry dating

Gamma spectrometry measurements were performed on 0.5-cm-thick sediment slices that were continuously sampled from the upper 15.0 cm of gravity core SC17_7 (Suppl. Table. 1). The samples were freeze-dried and sieved through a 200 μm mesh
145 for homogenization and removal of larger plant particles. Individual sample-aliquots were filled into gas-tight sealable low-activity Kryal[®] tubes at identical fill heights and accurately weighted. After sufficient in-growth-time, the gamma energies of ^{210}Pb ($T_{1/2}= 22$ a) and ^{214}Pb ($T_{1/2}= 26.8$ min), which is a daughter nuclide of ^{222}Rn ($T_{1/2}= 3.8$ d), were measured at 46.54, 295.24 and 351.93 keV. In addition, the gamma energies of ^{137}Cs ($T_{1/2}= 30.1$ a) were measured at 661.66 keV. For this purpose, the Kryal[®] tubes were placed into shielded measurement chambers equipped with two well-type germanium detectors G1 and
150 G2 (Canberra Industries) for ~1.5 to 7 days at GFZ Potsdam (Suppl. Tab. 1) (Schettler et al., 2006). Hardware control, data storage, and spectrum analysis were realized with the software Genie 2000 (Canberra Industries). The average counting uncertainty for ^{210}Pb was 5.9 %, for ^{214}Pb 7.7 % (295 keV) and 3.7 % (351 keV) and for ^{137}Cs 5.2 %. Efficiency calibrations were carried out for ^{210}Pb , ^{214}Pb and ^{137}Cs with the same analytical setup using a lab-internal standard and the “Loess Nussloch” standard (Potts et al., 2003). Blank activities for ^{137}Cs were negligible while average ^{210}Pb blank activities of 10 mBq/g for
155 detector G2 and ^{214}Pb blank activities of 9 mBq/g for the detectors G1 and G2 were considered. The activity measurements of ^{214}Pb were used to quantify the proportion of supported ^{210}Pb ($^{210}\text{Pb}_{\text{supp}}$) produced by the decay of ^{226}Ra in the sediment. The activity of unsupported ^{210}Pb ($^{210}\text{Pb}_{\text{unsupp}}$) in the sediment, which originates from the decay of ^{222}Rn in the atmosphere and associated aeolian deposition, is quantified by the difference between measured $^{210}\text{Pb}_{\text{total}}$ and $^{210}\text{Pb}_{\text{supp}}$. We selected sections that showed linear correlations in the semi-logarithmic plot of $^{210}\text{Pb}_{\text{unsupp}}$ versus depth to infer average sedimentation rates
160 using the constant initial concentration (CIC) model (c.f. Appleby, 2002) (Suppl. Tab. 2). Intercalated sediment sections showed nearly uncorrelated $\ln(^{210}\text{Pb}_{\text{unsupp}})$ vs. depth relationships at 10.25–9.25, 6.25–4.25 and 2.25–1.75 cm depth (Suppl. Fig.2). Therefore, the initial $^{210}\text{Pb}_{\text{unsupp}}$ activities of samples that bridged these sections were used alternatively to determine time intervals between these samples to infer a chronology. To assess possible changes of the sedimentation regime we

165 additionally calculated sedimentation rates of each 0.5-cm-thick sediment slice using the CRS model (constant initial $^{210}\text{Pb}_{\text{unSUPP}}$ supply) (c.f. Appleby, 2002; Appleby and Oldfield, 1978) (Suppl. Tab.3).

4. Results

4.1 Lithology

The composite profile can be subdivided into six lithological units (Fig. 3). Lithozone (LZ) I from 623.5 to 566.0 cm depth consists of greyish-brownish clastic-calcareous sediments. It shows mm-scale laminations of fine sandy and silty to clayey layers. LZ II (566.0–480.0 cm) exhibits intercalations between horizons of very fine, mm-scale laminated brownish-reddish organic-rich sediments and sections with greyish calcareous sediments. Brownish-reddish intercalating horizons of mm-scale laminated organic and calcareous sediments characterize LZ III from 480.0 to 273.0 cm depth. LZ IV (273.0–130.0 cm) is characterized by brownish-reddish mm-scale laminated organic-rich sediments with intercalated horizons rich in aquatic plant remains, which occur at 232.0–223.0 cm, 185.0–180.0 cm and 164.0–130.0 cm depth. LZ V (130.0–41.0 cm depth) starts with a 16-cm-thick interval of dark grey mm-scale laminated calcareous sediments, followed by brown mm- to cm-scale laminated sediments until 41.0 cm depth. Laminations are only poorly preserved between 63.0 and 41.0 cm depth. The uppermost sediments of LZ VI (41.0–0.0 cm depth) consist of homogenous, brownish-greyish calcareous sediments, which are rich in aquatic plant remains. The uppermost centimeter is enriched in calcite and exhibits greyish faint laminations.

4.2 Sediment microfacies analysis

180 Microscopic sediment analysis revealed, that clastic sublayers are present throughout the finely laminated sediments below 63.0 cm depth (Fig. 4.1). These clastic sublayers are variably intercalated with calcitic, aragonitic and organic sublayers and thus form different types of cyclic successions. In total, we classified six different types of sublayer successions as described below. The name for these types reflects the dominant sublayer for each of the six types. For example, the ‘clastic-organic type’ is characterized by the dominance of clastic sublayers, while in the organic-clastic type organic sublayers dominate. The names are not related to the order of sublayer succession within each type. Changing dominances of different sublayer successions reflect the lithozones.

4.2.1 Clastic-aragonitic type

190 Clastic-aragonitic laminae are rare (2.7 %), mainly occur in LZ I and particularly at 600.0-605.0, and 609.0-616.0 cm composite depth. This subtype is composed of three sublayers and the mean thickness is 0.59 mm (Fig. 4.1a, Suppl. Fig. 2a). These laminae exhibit the general pattern of clastic-organic laminae in LZ I, with a coarse-grained and thick basal detrital sublayer, but the overlying mixed (detrital calcite, mica, fsp, qtz and medium amounts of endogenic calcite) fine-grained sublayer additionally contains idiomorphic aragonite needles that are not found in clastic-organic varves. The sublayer succession ends with an amorphous organic matter sublayer.

4.2.2 Calcitic-clastic type

195 The deposition of calcitic-clastic laminae (6 %) with a dominating endogenic calcite sublayer is restricted to LZ II. This subtype is composed of three sublayers and the mean thickness is 0.41 mm, with a maximum of 2.0 mm. Calcitic-clastic laminae (Fig. 4.1b lower part, Suppl. Fig. 2b) are usually characterized by a basal detrital sublayer which, however, is not developed in all calcitic-clastic laminae. The overlying sublayer generally exhibits low species abundancies of diatom frustules, chrysophyte cysts, aquatic plant remains, daphnia, ostracods and characeae but massive and fine-grained endogenic calcite, which is not
200 the case in the clastic-calcitic laminae subtype. Endogenic calcite formed in the water column (Fig. 4.2a) is recognized by its well-developed idiomorphic rhombohedral shapes. Scattered detrital grains occasionally occur within the endogenic calcite matrix. One depositional cycle ends with an amorphous organic matter sublayer.

4.2.3 Clastic-diatom type

Clastic-diatom laminae (20 %) occur in LZ II, III and IV. This subtype is composed of three sublayers and the mean thickness
205 vary between 0.28 mm (LZ II), 0.34 mm (LZ III) and 0.35 mm (LZ IV). The depositional cycle starts with a basal detrital sublayer, which is overlain by a finer-grained mixed sublayer (detrital calcite, mica, fsp, qtz) occasionally containing chrysophytes and different diatom taxa. The third sublayer is formed by diatom blooms exclusively consisting of the planktic diatom species *Cyclotella choctawhatcheeana* (pers. comm. Anja Schwarz, TU Braunschweig) (Fig. 4.1b upper part, Suppl. Fig. 2c).

210 4.2.4 Clastic-calcitic type

The second most common laminae subtype (23.5 %) are clastic-calcitic laminae (Fig. 4.1c lower part, Suppl. Fig. 2d), which are most abundant in LZ I, II, III and V. This subtype is composed of four to five sublayers and the mean total thickness varies between 0.95 mm (LZ I), 0.35 mm (LZ II), 0.72 mm (LZ III), 1.56 mm (LZ IV) and the maximum value of 5.0 mm in LZ V. Clastic-calcitic laminae exhibit a basal detrital sublayer with a sharp lower boundary, which is followed by a bloom layer of
215 chrysophytes and/or diatoms, occurring sporadically after and/or within the detrital sublayer. The third, overlying mixed sublayer contains medium amounts of endogenic as well as fine-grained detrital calcite (Fig. 4.3a), as well as mica, fsp and qtz grains but low amounts of diatom frustules and chrysophyte cysts. One depositional cycle typically ends with an amorphous organic matter sublayer. In LZ V, these clastic-calcitic laminae occasionally contain a very fine-grained, light greyish, micritic sublayer before the cycle ends with the amorphous organic sublayer.

220 4.2.5 Organic-clastic type

Horizons of organic-clastic laminae (5.2 %) with dominating organic sublayers are mainly present within LZ IV and V (Fig. 4.1d, Suppl. Fig. 2e) particular at 261.0-252.0, 176.0-173.0, 150.0-126.0 and 122.0-110.0 cm depth. This subtype is composed of three sublayers and the mean thickness is 0.49 mm (LZ IV) and 1.64 mm (LZ V) with a maximum of 9 mm in LZ V.

Organic-clastic laminae exhibit an often horizontally discontinuous basal detrital sublayer (lens-shaped) in LZ IV, which is
225 overlain by a mixed sublayer that contains detrital calcite, mica, fsp and qtz grains and many aquatic plant remains and
periphytic diatoms (*Achnanthes brevipes*, pers. comm. Anja Schwarz, TU Braunschweig), whose colony chains are often
preserved. One deposition cyclic ends with a yellowish amorphous organic matter layer.

4.2.6 Clastic-organic type

Clastic-organic laminae are present in all lithozones and most abundant in the record (42.5 % of all observed and measured
230 laminae). This microfacies type is composed of four sublayers of which the most prominent is a basal clastic-detrital sublayer
with a sharp lower boundary. The basal detrital sublayer contains mainly detrital calcite, which is distinguished from endogenic
calcite by microscopic analyses. Detrital calcite is characterized by irregularly shaped grains and generally larger grain sizes of
average of 0.6 mm and up to 1.82 mm in LZ I. Detrital layers further contain siliciclastic minerals as mica, quartz (qtz) and
feldspars (fsp). This basal layer is often, but not regularly overlain by chrysophyte and/or diatom blooms and a third, mixed
235 sublayer containing mainly fine-grained detrital calcite, mica, fsp and qtz with low amounts of endogenic calcite and varying
amounts of diatom frustules, chrysophytes, characeae, ostracods and daphnia. The deposition cycle ends with a yellowish layer
of amorphous organic material.

The mean thickness of clastic-organic laminae differs between the lithozones. In LZ I, the mean thickness is 0.59 mm with a
maximum thickness of 3.1 mm. In LZ I, the basal detrital sublayer is thick and coarse-grained, rich in pyrite, and contains
240 mainly silt-to fine sand-sized grains and occasionally sand-sized qtz, calcite and fsp grains, whereas diatoms and chrysophytes
are rare. In LZ II and III, clastic-organic laminae are less thick with a mean thickness of 0.27 mm and 0.48 mm respectively.
In these lithozones, the basal sublayer contains no sand-sized particles. In LZ IV, mean varve thickness is 0.43 mm and the
basal sublayer is often lens-shaped and horizontally discontinuous. In LZ V between 130.0 and 63.0 cm depth thickest clastic-
organic laminae occur with a mean thickness of 1.5 mm and a maximum thickness of up to 7.0 mm (Fig. 4.1e, Suppl. Fig. 2f).
245 These clastic-organic laminae often include an additional detrital sublayer intercalated in the finer grained mixed sublayer.

4.2.7 Homogenous sediments

The uppermost 41.0 cm of the sediment record consist of homogenous sediments, containing a fine-grained mix of
autochthonous and allochthonous calcite, mica, qtz and fsp. The sediments are generally rich in organic remains, such as
aquatic plant remains, chrysophytes, diatoms and chlorophytes (*Botryococcus*). Faint and discontinuous calcite laminae occur
250 in the uppermost centimeter (Fig. 4.1f).

4.3 XRF element mapping

The two selected impregnated sediment blocks from 507 to 497.5 cm (XRF-Map 1 Fig. 4.2) and from 346.5 to 338.5 cm depth
(XRF-Map 2 Fig. 4.3) contain calcitic-clastic, clastic-diatom, clastic-calcitic and clastic-organic microfacies types. These
sediments are dominated by alternating calcitic and siliciclastic sediments represented by the elements Ca, Sr, Mg and Si, Al

255 respectively (Fig. 4.2 and 4.3). Color variations of the element maps show that the calcitic and siliciclastic sediments are clearly separated in sample the XRF-Map 1 (Fig. 4.2) but slightly more mixed in the XRF-Map 2 (Fig. 4.3). In both XRF-Maps 1 and 2, the carbonate sublayers are enriched in Sr (Fig. 4.2 and Fig. 4.3), whereas in the XRF-Map 1 the additional enrichment of Mg (Fig. 4.2) indicates the presence of Sr- and Mg-rich carbonates. Microfacies analysis show that Mg- and Sr- rich calcite sublayers in XRF-Map 1 are predominantly of endogenic origin (Fig. 4.2a), whereas the Sr- rich calcite layers in XRF-Map 2 contain mixed endogenic and resuspended calcites (Fig. 4.3a). Moreover, detrital carbonates occur predominantly as individual grains in the siliciclastic sediments and sublayers and are shown by Ca and Mg in the XRF element maps (Fig. 4.2 b and Fig. 4.3 b). Siliciclastic muds of clastic-organic and clastic-diatom varves are represented by the co-occurrence of Al and Si in the XRF element maps, whereas Al is absent in diatomaceous sublayers (Fig. 4.2 and 4.3).

4.4 Chronology

265 4.4.1 Floating varve chronology (623.5 - 63.0 cm)

A floating varve chronology labelled as Chatvd19 (Fig. 5b) was established for the composite profile below 63.0 cm depth and comprises a total of 11,259 counted and interpolated varves. Based on the interpretation of laminations as varves, 9,026 of the total 11,259 varves were counted which is equal to 80.2 %. The first varve count reveals 9,026 varves and is the base for the floating varve chronology. Although the total varve number of 8,955 obtained by the second count is very similar to the first count, larger deviations between the two varve counts in individual sediment sections occur throughout the sediment record due to varying stages of varve preservation as expressed in the VQI (Fig. 5a). Largest deviations occur in LZ I (603.0-595.0 cm) with 23.7 %, in LZ II (490.0-484.0 cm) with ~13 %, in LZ III (413.0-405.0 cm) with ~16 %, in LZ IV (141.0-134.0 cm depth) with 19.5 % and in LZ V (65.0-63.0 cm depth) with 7.7 %. Lowest deviations (<1 %) were obtained in LZ II at 539.0-530.0 cm and 498.0-490.0 cm, in LZ III at 451.0-445.0 cm, 421.0-413.0 cm, 375.0-369.0 cm, 299.0-290.0 cm and 282.0-275.0 cm, in LZ IV at 197.0-190.0 cm and in LZ V at 125.0-119.0 cm, 116.0-113.0 cm and 73.0-65.0 cm depth. Interpolated sequences are unevenly distributed within the record and are mainly present within LZ IV. The longest interpolated sequences occur in LZ IV with ~5 cm from 198.0-193.0 cm depth and in LZ III with almost 7 cm between 444.0-437.0 cm depth. A VQI (Fig.5a) of 1 is represented by 5.6 % of the total varves, VQI 2 by 8.4 %, VQI 3 by 26.4 %, VQI 4 by 18.3 % and VQI 5 by 21.4 %. The calculated mean deviation between the two varve counts of ~5 % (Fig. 5a) is used as a conservative uncertainty for the floating varve chronology to consider high uncertainties in individual sediment sections in a more realistic way, despite the similar total number of varves counted. The floating varve chronology has basal age of 11619 ± 603 years BP.

4.4.2 Chronology of the non-varved uppermost sediments

The uppermost 63.0 cm of the sediment profile are not varved and thus require alternative dating approaches including ^{210}Pb dating, activity profiles of ^{137}Cs and sedimentation-rate based interpolation. First, we measured ^{210}Pb activity concentrations of the uppermost 15 cm of short core SC17_7 and applied the CIC and CRS models (Fig. 5c & 6, Suppl. Tab. 3). SC17_7 is

correlated to the composite profile through **macroscopically visible facies change** at 1.0 cm composite depth and through a laminated section from 45.0-41.0 cm composite depth (Suppl. Fig.1). The CIC and CRS-model based chronologies are broadly consistent and particularly date sediments at 8.75 cm (SC17_7) or 7.5 cm composite depth to AD 1945/46 (Fig. 5c and Fig. 6b, Suppl. Table 3.). This **coincides** with the onset of increased ^{137}Cs activity concentrations (Fig. 5c, 6c) marking the onset of nuclear weapon testing in AD 1945 (Ferm, 2000; Kudo et al., 1998; Norris and Arkin, 1998). Therefore, we applied the date of AD 1945 (8.75 cm in core SC17_7) as an anchor point for the chronology of the uppermost 63.0 cm of the composite profile. According to micro-facies based sedimentological correlation, this anchor point is located at 7.5 cm composite depth. The section of homogeneous sediments from this point down to 63.0 cm depth was interpolated. This interpolation is based on sedimentation rate calculations obtained by lead-210 dating and varve thickness measurements in adjacent sediment intervals. Calculations include sedimentation rates from the upper 7.5 cm (1.12 mm/yr), from 15 varves (1.9 mm/yr) between 41.0-63.0 cm depth and from 100 varves (1.66 mm/yr) below 63.0 cm depth and result in a mean SR of 1.56 mm/yr corresponding to 356 interpolated years. Adding the number of 356 interpolated years to the radiometric date of AD 1945 results in an age of AD 1589 (360 years BP) at 63.0 cm depth. We assume a **conservative** uncertainty of ca. 10% **as a maximum error** for our interpolation. The anchor point thus has an age of 360 ± 40 years BP. The uncertainty of this anchor point is added to the varve counting uncertainty.

4.4.3 Radiocarbon dating

In total, we dated 36 samples of bulk organic carbon, daphnia remains, aquatic plant remains and *Ruppia maritima* seeds. Only two samples were terrestrial plant remains (wood fragments) and sufficiently large to be used for AMS ^{14}C dating (Tab. 2, Fig. 5 & 7). Except the two ages from terrestrial plant remains (Poz-54302 with 9988 ± 203 and Poz-63307 with 6140 ± 137 cal yr BP), all other ages deviate from the varve chronology between 155 years at 0.0 cm depth and 6,150 years at 585.0 cm depth (Fig. 7). We observe a general trend of decreasing deviations up core with the maximum deviation of ~6,150 years at 585.0 cm depth in LZ I. Looking at more detail, the deviations between radiocarbon and varve ages exhibit a prominent step-wise increase particularly at the boundary between lithozone LZ IV and LZ V when it abruptly decreases from ~3,000 years to ~1,000 years. Modern aquatic plants collected during the field campaign in 2012 showed large modern reservoir ages of 330 ± 30 and 2425 ± 25 ^{14}C years and living daphnia yielded ages of 225 ± 30 ^{14}C years.

5. Discussion

5.1 Interpretation of fine laminations as varves

The construction of varve chronologies relies on the proof of seasonal origin of fine laminations (Brauer et al., 2014; Ojala et al., 2012; Zolitschka et al., 2015). **Laminations are absent in the upper 63.0 cm of the Lake Chatyr Kol sediment core and cyclic successions of mixed-clastic laminations are only observed below this depth. Therefore, the seasonal origin of the Chatyr**

Kol sediments cannot be proofed through modern observation in sediment traps because no varves are formed at present day. Instead, we applied process-related deposition models (Fig. 4.1) based on detailed micro-facies analyses obtained from petrographic thin sections and compared our observations with varve types described in literature (Brauer, 2004; Zolitschka et al., 2015). We associate the observed successions of different types of mixed-clastic laminations with the formation of different seasonal sublayers that are known from lakes with carbonaceous catchments (Brauer and Casanova, 2001; Kelts and Hsü, 1978; Lauterbach et al., 2011; Lauterbach et al., 2019) and high-altitude glacial environments (Guyard et al., 2007; Leemann and Niessen, 1994). The observed succession of sublayers are interpreted as mixed varve types (clastic, -organic and – endogenic) as defined by Zolitschka et al. (2015).

Varve formation at Lake Chatyr Kol is related to the high seasonality of the local climate with an ice cover during winter as well as strong annual variations of the temperature and precipitation affecting productivity, endogenic carbonate formation and local runoff. Varve preservation is promoted by the unique morphology of the deep western lake basin, where anoxic bottom water conditions can be maintained even under relatively low lake levels (Fig.2).

We interpret the annual sedimentary cycle to always start with the deposition of a basal detrital sublayer with a sharp lower boundary which results from winter/spring snow and/or glacial melt (Guyard et al., 2007; Leemann and Niessen, 1994; Zolitschka et al., 2015) after the ice break-up in ~April (Shnitnikov, 1978). Runoff with suspended sediment load is then likely directed through the Kegagyry River in the east but may also be the result of surface runoff through the activation of several widely distributed smaller tributaries in the catchment (Fig. 1).

Basal detrital sublayers are generally overlain by blooms of chrysophytes and/or diatoms within clastic-organic, organic-clastic, clastic-diatom and clastic-calcitic varve types. Chrysophytes and/or diatom blooms develop in consequence of available nutrients provided by runoff and spring overturn in combination with rising temperatures during the summer season (Zolitschka et al., 2015). The productive phase in calcitic-clastic varves is however reflected by calcite precipitation, which is the main carbonate phase (endogenic, detrital and resuspended) in the Chatyr Kol sediments. The formation of endogenic calcite in Lake Chatyr Kol is controlled by: 1) photosynthesis, when high aquatic productivity lowers the concentrations of CO₂, increases the pH of the lake water and leads to a reduced solubility of CO₃²⁻ (Hodell et al., 1998; Kelts and Hsü, 1978; Zolitschka et al., 2015), 2) evaporation leading to an oversaturation of carbonate ions, 3) sufficient supply of dissolved cations either through surface runoff or groundwater inflow (Shapley et al., 2005). Changes in weathering and hydrological conditions can lead to variations in the supply of Ca²⁺ and Mg²⁺ ions and subsequently change the Mg/Ca ratio of the lake water (Müller et al., 1972). The formation of aragonite requires high lake water Mg/Ca ratios (>12), whereas magnesium-calcite forms at lower Mg/Ca ratios (Kelts and Hsü, 1978; Müller et al., 1972). XRF element intensity maps do not provide quantitative results but do indicate, that Mg is abundant in the XRF map 1 from LZ II which results in the formation of endogenic Sr- and Mg-rich calcites (Fig. 4.2 a).

Aragonite precipitates, related to an evaporative concentration in summer, were only microscopically observed in the intervals between 600.0-605.0 and 609.0-616.0 cm composite depth suggesting that Mg/Ca ratios probably remained above Mg/Ca ratios >12.

350 After the spring to early summer lake productivity, the deposition of a mixed sublayer consisting of silt- to clay-sized detrital grains and low to medium amounts of endogenic calcite is observed in all lamination types (Fig. 4.1, Suppl. Fig. 2). In clastic-calcitic varves, the mixed sublayers appear different and include especially resuspended calcite as evidenced also in Ca and Sr intensities (Fig. 4.1c, Fig. 4.3a, Suppl. Fig. 2d). The mixed sublayer indicates resuspension of shore material (littoral calcite) to the core location due to e.g. wind induced wave activity and weak runoff during the ice-cover free season from
355 ~April to October (Shnitnikov, 1978).

The intercalation of discrete detrital layers within the mixed sublayer (Fig. 4.1e, Suppl. Fig. 2e), as observed in clastic-organic laminae in LZ V, indicates pulses of runoff of suspended material which may be caused by late rainfall events in summer (Aizen et al., 2001; Shnitnikov, 1978).

360 One annual depositional cycle usually ends with the deposition of a thin sublayer of very fine amorphous organic matter which is deposited under quiet water conditions when the lake was ice covered (Fig. 4.1). In lithozone V, an additional micritic sublayer is deposited before the amorphous organic sublayer at the end of the seasonal cycle in clastic-calcitic laminae when water turbulence is low.

5.2 Varve counting and chronology construction

The interpretation of different types of fine laminations allowed varve counting as a main tool for constructing the Chatyr Kol
365 chronology largely based on incremental methods. Around 80% of the varves in the sediment record are double-counted in petrographic thin sections while the remaining part of ca 20% had to be interpolated based on sedimentation-rate estimates due to poor varve preservation. The resulting chronology comprises 11,259 years and is anchored to the absolute time scale at 63.0 cm sediment depth supported by a combination of lead-210 dating and occasional sedimentation rate measurements as described below. The resulting age-depth model is within uncertainties in good agreement with two calibrated AMS ¹⁴C dates
370 of wood pieces at 380.5 cm depth ($6,140 \pm 137$ cal years BP; Poz-63307) and at 528.0 cm depth ($9,988 \pm 203$ cal years BP; Poz-54302) (Fig. 5b, Tab. 2). The corresponding varve-based ages are $5,905 \pm 320$ years BP and $9,611 \pm 505$ years BP, respectively. As for all chronologies, uncertainties are inherent also to varve chronologies, which are commonly assessed via replicate counts (Brauer and Casanova, 2001; Lamoureux, 2001; Lotter and Lemcke, 1999; Ojala et al., 2012; Żarczyński et al., 2018; Zolitschka et al., 2015). However, there is no standard procedure on how to calculate and present the uncertainties
375 (Ojala et al., 2012; Zolitschka et al., 2015). Commonly, mean values of replicate count differences, the difference of maximum and minimum counts or their standard deviation are reported (Brauer et al., 2014; Ojala et al., 2012; Żarczyński et al., 2018; Zolitschka et al., 2015). Despite the inevitable increase of cumulative uncertainties with age or depth, systematic uncertainties arise and are caused by changes in varve preservation, strongly and abruptly varying sedimentation rates and the challenging differentiation of varve types with complex structures (Ojala et al., 2012; Żarczyński et al., 2018; Zolitschka et al., 2015). The
380 overall very small difference between the two counts of the Chatyr Kol varved record of only -71 varves is due to the compensating effect between over- and underestimations of varve counts throughout the record. For the floating varve

chronology, we therefore compare the results for each individual thin section comprising **between a maximum of 324 (506.8-497.6 cm) and a minimum of 13 (varves)** (65.4-63.0 cm) (Fig. 5a, Fig. 8).

Counting uncertainties for individual thin sections are reported as their percentage deviation from the first count used for the
385 chronology and range between 0 and 23.7 % (Fig. 5a, Fig. 8). Largest deviations of 23.7 % in LZ I are caused by a low visibility
of varve boundaries and by coring artefacts. Deviations of ~13 % in LZ II and of ~16 % in LZ III result from the abrupt
intercalations between clastic-organic, clastic-diatom, clastic-calcitic and calcitic-clastic varve types with varying varve
thicknesses (Fig. 4 LZ II & LZ III). Deviations in LZ IV with a maximum of 19.5 % coincide with generally lowest VQI values
(Fig. 8) and result from the domination of clastic-organic and organic-clastic varves with lowest thicknesses and discontinuous
390 basal detrital sublayers (Fig. 4 LZ IV) leading to generally higher counting uncertainties. Lowest deviations of <1 % in LZ II,
LZ III, LZ IV and in LZ V represent best preservation and thus easily countable varves of different varve types. Generally,
clastic-organic and clastic-calcitic varves with higher varve thickness, especially in LZ V are most reliably countable. The
relatively high uncertainties of 7.7 % in LZ V are due to the low number of varves comprised in individual thin sections (Fig.
8). For the total uncertainty estimate for the floating varve chronology we use the mean of ± 5 % calculated from the
395 uncertainties for each 10 cm interval. This conservative estimate is more realistic than the very low difference in the two
repeated varve counts. An uncertainty of 5 % is in the range of elsewhere reported varve chronologies (Ojala et al., 2012).
Since the upmost 63.0 cm of the sediment profile are largely homogeneous, the varve chronology is floating and needs to be
anchored to an absolute chronology at this point. The interpolation with a mean SR derived from the combination of the
consistent CIC and CRS ^{210}Pb marker (AD 1945), the SR derived from discontinuously varved sequences between 41.0 and
400 63.0 cm depth and from 100 measured varves below 63.0 cm depth seems to be the best approach for constraining the
uppermost age-depth relationship within the homogenous sediments, where further chronological markers are lacking. We are
aware, that the interpolation-based “floating” anchor point at 63.0 cm depth is prone to additional uncertainties. We considered
this by assuming a higher uncertainty of 10 % for this interval, than that of ± 5 % for the floating varve chronology.

5.3 Radiocarbon reservoir effects

405 Compared to the floating varve chronology, including two terrestrial (wood) AMS ^{14}C dates, we observed a general trend of
decreasing reservoir effects of dated aquatic material up core with the maximum deviation of ~6,150 years at 585.0 cm depth
(10,930 \pm 570 years BP) in LZ I (Fig. 7). The step-wise decrease of deviations between radiocarbon and varve ages is most
pronounced at the boundary between lithozone LZ IV and LZ V, when it abruptly decreases from ~3,000 years to ~1,000 years.
The reservoir effect generally depends on the rate of atmospheric CO_2 exchange between the water column and the air, internal
410 mixing dynamics and the input of ^{14}C depleted carbonaceous material (Ascough et al., 2010; Jull et al., 2013; Keaveney and
Reimer, 2012; Lockett et al., 2016; MacDonald et al., 1991). The catchment of Lake Chatyr Kol exhibits several sources that
could be responsible for a ^{14}C -depletion of dissolved carbon species in the lake water. Highest reservoir ages in the early
Holocene are likely the result of the combined influence of these sources: 1) the input of old, ^{14}C -depleted CO_2 with glacial
meltwater (c.f. Hall and Henderson, 2001) at the onset of a warming Holocene and 2) the weathering and erosion of the northern

415 outcropping limestones, which led to the release and input of dissolved bicarbonate to the lake (c.f. Abbott and Stafford, 1996; Hutchinson et al., 2004). Both processes lead to a ^{14}C -depleted CO_2 and HCO_3^- uptake during photosynthesis by e.g. submerged aquatic plants like *Ruppia maritima* at 585.0 cm depth (Fig.7) and by phytoplankton, on which daphnia feed and which therefore also show similar reservoir effects. A high detrital input and thus a potentially high input of dissolved bicarbonate is supported by increased varve thicknesses during the early Holocene (Sect. 5.4.1, Fig. 9). Furthermore, 3) thawing of permafrost since the beginning of a warming Holocene might have released dissolved ^{14}C -depleted organic material and thus affect the ^{14}C TOC bulk measurements. Our fieldtrip observations and observations by Shnitnikov (1978) of modern permafrost reduction and the development of thermokarst in the southern part of the catchment around the neighbouring Lake Maloye (Fig. 1) support this assumption. The cause of a step-wise reservoir effect reduction is therefore likely also related to the combined effect of a generally decreasing glacial influence and a decreasing input of bicarbonate until ~AD 1150 at the boundary between LZ IV and LZ V (Fig.7, 9). The abrupt decrease of the reservoir effect after ~AD 1150, despite an increase in detrital carbonate supply (Sect. 5.4.5, Fig. 9) might be related to the silting up of the basin leading to a shallower water depth, which is more susceptible to water circulation and an enhanced atmospheric CO_2 exchange (c.f. Geyh et al., 1997).

5.4 Holocene variations in varve microfacies

The Lake Chatyr Kol sediment profile comprises six different varve types (Sect. 4.2, Fig. 4.1, Suppl. Fig. 2), which occurrences showed varying dominances in the different lithozones that are described below. The individual lithozones always comprised more than one varve type (Fig. 9) with a maximum of five different varve types occurring in LZ III and II to three varve types in LZ V.

5.4.1 Lithozone I (623.5-566.0 cm: 11,619 \pm 603 to 10,730 \pm 560 years BP)

435 Lithozone I is characterized by relatively high and varying varve thicknesses and by the presence of clastic-organic, clastic-calcitic and clastic-aragonite varves (Fig. 9 LZ I). Clastic-organic varves constitute about 57 % of the observed and counted varve types, clastic-calcitic 29 % and clastic-aragonitic 14 %. Generally thick detrital coarse-grained spring sublayers, which were observed in all varve types in this LZ, are indicative for intense runoff by winter/spring snow meltwater and/or by glacial thawing during summer (Shnitnikov, 1978) caused by highest insolation (Berger and Loutre, 1991; Chen et al., 2008; Jin et al., 2011; Li and Morrill, 2010) at the onset of a warming early Holocene. Glaciers of the inner Tian Shan started to retreat between ~12-8 ka years BP (Bondarev, 1997; Shnitnikov, 1978) causing enhanced detrital input into the lake. The low species abundancies of aquatic plants (*Ruppia Maritima* or *Potamogeton* sp.), daphnia and characeae reflect a littoral community and indicate a low aquatic productivity and a relative low lake level during this time. Clastic-calcitic varves appear at the base of the composite profile and towards the end of LZ I, whereas clastic-aragonitic varves dominate in the period from ~11,500 to 445 11,000 years BP. Most likely, idiomorphic aragonite formed due to a combination of Mg-rich water supply to the lake and strong evaporative conditions causing lake water Mg/Ca ratios of >12 (Kelts and Hsü, 1978; Müller et al., 1972).

5.4.2 Lithozone II (566.0-480.0 cm: 10,730 ± 560 to 8,040 ± 430 years BP)

In this lithozone calcitic-clastic varves constitute about 21 % of the observed varves and clastic-calcitic varves ~22 %, while
450 clastic-organic varves make up ~42 % and clastic-diatom varves ~15% (Fig. 9 LZ II). This lithozone is characterized by
intercalations of calcitic varve types (calcitic-clastic & clastic-calcitic) with clastic-organic and clastic-diatom varves (Fig.
4.2). The variations of these varve types might be related to external (climatic) forcing or lake-internal or sedimentation
variability (Turner et al., 2016 and reference herein). XRF element maps show endogenic calcite sublayers that are enriched
in Sr and Mg alternating with clastic (Si and Al) or diatom (Si) layers (Fig. 4.2) suggesting Sr- and Mg-rich calcite in this
455 lithozone which indicates evaporative concentration (Müller et al., 1972). The shift of the dominant endogenic carbonate type
from aragonite in LZ I to calcite in LZ II around ~10,730 years BP (Fig. 9 LZ II) coincides with an increase in biological and
photosynthetic activity, as inferred from the establishment of a diverse lake fauna seen in high abundancies of chrysophytes,
planktic (*Cyclotella choctawhatcheeana*) and periphytic diatoms (*Achnanthes brevipes*) as well as of aquatic plants, ostracods,
characeae and few daphnia. Identifying the main drivers controlling endogenic carbonate formation thus remains speculative.
460 Species assemblages and associated biological activity during the summer season indicates favourable warm summers and
sufficient nutrient supply through e.g. runoff. Because the species assemblages show mixed littoral and pelagic species
abundancies, these are interpreted as an indication for a low lake level.

5.4.3 Lithozone III (480.0-273.0 cm: 8,040 ± 350 to 4,140 ± 230 years BP)

465 The deposition of clastic-calcitic varves comprise ~38 % of the observed varves in this lithozone, while clastic-organic varves
make up ~27 % and clastic-diatom varves ~34 % (Fig. 9 LZ III). Clastic-calcitic varves are generally thicker than the other
varve types of this LZ mainly because of exceptionally thick summer sublayers. These summer layers consist of endogenic
calcite mixed with resuspended calcites and fine-grained detrital grains (Fig. 4.1c, Fig. 4.3 a). This is confirmed by elevated
Sr values of the XRF element mapping, indicating the presence of Sr-rich carbonates. These varves likely reflect increased
470 resuspension of carbonates from the littoral zone due to wind induced wave activity. As in LZ II, alternations of clastic-calcitic
(Ca, Sr) (Fig. 4.3 a) and clastic-diatom, clastic-organic (Si, Al) (Fig. 4.3 b) varves are characteristic for this lithozone as well.
At ~8,040 years BP the deposition of calcitic-clastic varves ceased and is replaced by the deposition of clastic-calcitic, clastic-
organic and clastic-diatom varves probably due to decreasing summer insolation and lower summer temperatures (Berger and
Loutre, 1991; Chen et al., 2008; Jin et al., 2011; Li and Morrill, 2010). In addition, higher lake levels since that time are
475 indicated by the dominance of planktic diatoms and the occurrence of lake deposits at the eastern and southern shore which
have been dated from 6,688 ± 473 to 4,621 ± 594 cal years BP (¹⁴C ages published by Shnitnikov (1978) calibrated with
OxCal4.3 & IntCal13). During our field work we found lake sediments on a shallow terrace ~7 m above the current lake level
east of the lake which also revealed a mid-Holocene age of 5,786 ± 122 cal years BP (Poz-109830 Tab.2). Higher lake levels
at that time have been explained by the preceding early Holocene glacier retreat in the catchment (Bondarev, 1997; Shnitnikov,

480 1978). However, more recently even minor glacial advances in the Aksai Basin east of the Chatyr Kol catchment ^{10}Be exposure dated between 7.5 and ~ 4.5 ka were reported (Koppes et al., 2008).

5.4.4 Lithozone IV (273.0-130.0 cm: $4,140 \pm 230$ to 800 ± 60 years BP/ AD 1150 ± 60)

485 This Lithozone contains clastic-organic (58 %), organic-clastic (18 %), clastic-diatom (17 %) and clastic-calcitic varves (6 %) (Fig. 9 LZ IV). High abundancies of planktic diatoms as well as clastic-diatom varves prevail until $\sim 2,900$ years BP, whereas clastic-organic and organic-clastic varves with abundant aquatic plant remains and periphytic diatoms occur afterwards and dominate the sediments particularly after $\sim 2,200$ years BP. Abundant aquatic plant remains and periphytic diatoms can be explained by reworking due to wave activity and water column mixing during low lake levels. Low lake levels are inferred from the modern observation that aquatic plants occupy the shallow parts of the Lake from ~ 15.0 to 0.5 m. A decreased lake level combined with the shallow bathymetry of the lake basin (Fig. 1) further promotes large impacts on the species communities, which supports the changing abundancies from a planktic to a littoral dominated fauna and flora after $\sim 2,200$ years BP. An increased mixing of the water body also caused a clear decline in varve preservation. The rate of detrital input as observed in clastic-organic, organic-clastic varves (Fig. 4.1d, Suppl. Fig. 2e) is rather constant and mainly appears within the spring sublayer suggesting stable snow melt runoff from 4,140 years BP to AD 1150. At $\sim 2,600$ and at 2,300 years BP 495 clastic-calcite varves with thickened summer sublayers appear for a few decades indicating enhanced resuspension.

5.4.5 Lithozone V (130.0-41.0 cm: AD 1150 ± 60 to AD 1730 ± 30)

500 Clastic-organic varves constitute 59 % of the observed varves in LZ V, clastic-calcitic varves 26 % and organic-clastic varves 15 % (Fig. 9 LZ V), the latter ceasing at 110.5 cm (AD 1260 ± 50). Varve microfacies changes abruptly at 130 cm depth or AD 1150 from the dominance of organic-clastic varves to dominating clastic-organic and clastic-calcitic varves. Within 5 years, varve thickness drastically increased from $\varnothing 0.43$ mm in LZ IV to $\varnothing 1.52$ mm in LZ V due to thicker summer sublayers. Thicker summer sublayers result from thicker mixed sublayers rich in algae remains (*Botryococcus*, chrysophytes, diatoms) and additional late summer detrital sublayers (Fig. 4.1.e, Suppl. Fig. 2f). Hence, the increase in summer layer thickness suggest both, higher lacustrine productivity and an increase in summer runoff events. However, the reasons for these changes remain 505 elusive and a relation to known climatic periods like the Medieval Climate Anomaly and the Little Ice Age is not found. One might speculate that the frequent occurrence of late summer runoff layers either reflects convective rainfall events due to recycling of local moisture sources (Aizen et al., 2001) or changing atmospheric circulation regimes. Changes in boundary conditions in the catchment of the lake are unlikely since microfacies analyses does not show pronounced changes in grain size distribution of the detrital material. Human impact cannot fully be excluded but low indices of human and livestock fecal 510 biomarkers (Schroeter et al., 2020) are an argument against major human impact. The presence of lake deposits at the northern and southern shores ca 1.5 - 1 m above present day lake level dated at AD 1420 ± 204 , AD 1044 ± 160 and AD 858 ± 166 (Shnitnikov, 1978) suggests that increased summer runoff events might have resulted in a more positive water budget and lake level rise.

515 **5.4.6 Lithozone VI (41.0-0.0 cm: AD 1730 ± 30 to AD 2012)**

At around AD 1730 ± 30 varve formation and/or preservation ceased and sediments became predominantly homogeneous. The cessation of varves might be related to enhanced mixing of the water column resulting in a loss of the oxygen minimum zone (Fig. 2a) caused by decreasing water depth due to silting-up, which accelerated with the abrupt increase in sedimentation rate at AD 1150 and/or due to strengthening of the wind conditions and wave activity.

520 **6. Conclusion**

We present the first varved lake sediment record in arid Central Asia that covers almost the entire Holocene. The established floating varve chronology provides an independent dating for a setting with scarce material for radiocarbon dating. In particular, our varve chronology allows a quantification of changes in radiocarbon reservoir ages throughout the Holocene. The largest reservoir effect of ~6150 years in the early Holocene is likely caused by glacial melt and enhanced local erosion resulting in a surplus of dead carbon. [Lowest reservoir ages of ~1,000 years and less in the late Holocene might be related to enhanced atmospheric CO₂ exchange when the lake was shallower due to silting-up of the lake basin and/or increased windiness inducing increased water column mixing and CO₂ exchange with the atmosphere.](#) The construction of the varve-based chronology was only possible through detailed micro-facies analyses of the entire sediment sequence in overlapping thin sections [that allowed the development of](#) seasonal deposition models for all observed types of fine laminations. Based on these models and their comparison with published varve micro-facies data, we interpret all six Chatyr Kol lamination types as varves. Compared to many other varved lake sediment records, the Chatyr Kol varves are very heterogeneous and a complex pattern of six different micro-facies types developed throughout the Holocene. All varve types are predominantly clastic and comprise variations of their summer sublayers with changing dominances of organic, diatom, calcitic, aragonitic and additional detrital sublayers. Varve thickness changed accordingly with the varve micro-facies types, whereby the most conspicuous increase of varve thickness occurred at AD 1150 which is caused by increased erosion and runoff. The increase in detrital input into the lake further caused an acceleration of the silting-up processes.

[XRF element mapping results support our microfacies analysis and provide additional information on the composition of carbonate sublayers and detrital carbonate. Microfacies analysis and XRF element mapping show variations between Mg and Sr rich calcitic sublayers of calcitic-clastic and clastic-calcitic varves. Additionally, XRF element mapping results of the elements Al and Si clearly distinguish clastic-organic and clastic diatom varves.](#) Nevertheless, the complex succession and variations of varve types throughout the Holocene including major change points still requires further detailed investigations and interpretation together with other proxy data.

Data availability

The presented data is provided through PANGAEA: <https://doi.pangaea.de/10.1594/PANGAEA.909981>
545 and at <https://varve.gfz-potsdam.de>.

Competing interest

The authors declare that they have no conflict of interest.

Authors contributions

JK performed the microfacies analysis, ^{210}Pb and ^{137}Cs gamma spectrometry and wrote the manuscript with contributions from
550 all co-authors. JM designed the project and organised field work. SL carried out sediment coring and was responsible for ^{14}C
dating. RU provided information about the catchment geology and lake level changes. RT carried out the XRF element
mapping. AB supervised analyses and manuscript writing.

Acknowledgements

We are grateful for the suggestions of three anonymous reviewers which helped to improve the manuscript. We thank S.
555 Lauterbach, S. Pinkerneil, M. Köhler, R. Schedel, and D. Henning for retrieving the long piston cores from Lake Chatyr Kol
in 2012 in the framework of the project CADY (Central Asian Climate Dynamics, BMBF grant 03G0813). S. Pinkerneil and
Y. Beutlich are thanked for help with the geochemical analyses. S. Orunbaev, M. Daiyrov, S. Kalmuratov, G. Omurova and
K. Jusupova are acknowledged for their support during field trips. We further thank T Goslar for AMS ^{14}C dating and D.
Berger, G. Arnold and B. Brademann for thin section preparation and G. Schettler for his help with lead-210 dating. We also
560 want to thank Rik Tjallingii for conducting μXRF element mapping and his help with the revision of the manuscript. This
paper is a contribution to Topic 8 'Rapid Climate Change from Proxy data' within the climate initiative REKLIM of the
Helmholtz-Association.

Funding

This study was conducted in the framework of CAHOL (Central Asian HOlocene), a subproject of the joint project CAME
565 II (Central Asia Climate Tipping Points and their Consequences) funded by the German Federal Ministry of Education
(BMBF) through grant 03G0864B.

References

- 570 Abbott, M. B. and Stafford, T. W.: Radiocarbon geochemistry of modern and ancient Arctic lake systems, Baffin Island, Canada, *Quaternary Research*, 45, 300-311, <https://doi.org/10.1006/qres.1996.0031>, 1996.
- Academy of Science of the Kyrgyz SSR: Atlas of the Kyrgyz Soviet Socialistic Republic: Natural conditions and resources, State Agency of Cartography and Geodesy, Central Directorate for Geodesy and Cartography, Council of Minister of the USSR, Moscow, 1987
- 575 Aizen, E. M., B., A. V., M., M. J., Nakamura, T., and Ohta, T.: Precipitation and atmospheric circulation patterns at mid-latitudes of Asia, *International Journal of Climatology*, 21, 21, <https://doi.org/10.1002/joc.626>, 2001.
- Aizen, V. B.: Association between atmospheric circulation patterns and firn-ice core records from the Inilchek glacierized area, central Tien Shan, Asia, *Journal of Geophysical Research*, 109, <https://doi.org/10.1029/2003JD003894> <https://doi.org/10.1029/2003JD003894>, 2004.
- Appleby, P.: Chronostratigraphic techniques in recent sediments. In: *Tracking environmental change using lake sediments*, Springer, https://doi.org/10.1007/0-306-47669-X_9, 2002.
- 580 Appleby, P. and Oldfield, F.: The calculation of lead-210 dates assuming a constant rate of supply of unsupported 210Pb to the sediment, *Catena*, 5, 1-8, [https://doi.org/10.1016/S0341-8162\(78\)80002-2](https://doi.org/10.1016/S0341-8162(78)80002-2), 1978.
- Ascough, P., Cook, G., Church, M., Dunbar, E., Einarsson, Á., McGovern, T., Dugmore, A., Perdikaris, S., Hastie, H., and Friðriksson, A.: Temporal and spatial variations in freshwater 14 C reservoir effects: Lake Mývatn, northern Iceland, *Radiocarbon*, 52, 1098-1112, <https://doi.org/10.1017/S003382220004618X>, 2010.
- 585 Berger, A. and Loutre, M.-F.: Insolation values for the climate of the last 10 million years, *Quaternary Science Reviews*, 10, 297-317, [https://doi.org/10.1016/0277-3791\(91\)90033-Q](https://doi.org/10.1016/0277-3791(91)90033-Q), 1991.
- Bondarev, L. G.: Fluctuations of local glaciers in the southern ranges of the former USSR: 18,000-8000 BP. [https://doi.org/10.1016/S1040-6182\(96\)00023-7](https://doi.org/10.1016/S1040-6182(96)00023-7), 1997.
- 590 Boomer, I., Aladin, N., Plotnikov, I., and Whatley, R.: The palaeolimnology of the Aral Sea: a review, *Quaternary Science Reviews*, 19, 1259-1278, [https://doi.org/10.1016/S0277-3791\(00\)00002-0](https://doi.org/10.1016/S0277-3791(00)00002-0), 2000.
- Brauer, A.: Annually Laminated Lake Sediments and Their Palaeoclimatic Relevance. In: *The Climate in Historical Times: Towards a Synthesis of Holocene Proxy Data and Climate Models*, Fischer, H., Kumke, T., Lohmann, G., Flöser, G., Miller, H., von Storch, H., and Negendank, J. F. W. (Eds.), Springer Berlin Heidelberg, Berlin, Heidelberg, https://doi.org/10.1007/978-3-662-10313-5_7, 2004.
- 595 Brauer, A. and Casanova, J.: Chronology and depositional processes of the laminated sediment record from Lac d'Annecy, French Alps, *Journal of Paleolimnology*, 25, 163-177, <https://doi.org/10.1023/A:1008136029735>, 2001.
- Brauer, A., Hajdas, I., Blockley, S. P. E., Bronk Ramsey, C., Christl, M., Ivy-Ochs, S., Moseley, G. E., Nowaczyk, N. N., Rasmussen, S. O., Roberts, H. M., Spötl, C., Staff, R. A., and Svensson, A.: The importance of independent chronology in integrating records of past climate change for the 60–8 ka INTIMATE time interval, *Quaternary Science Reviews*, 106, 47-66, <https://doi.org/10.1016/j.quascirev.2014.07.006>, 2014.
- 600 Chen, F., Yu, Z., Yang, M., Ito, E., Wang, S., Madsen, D. B., Huang, X., Zhao, Y., Sato, T., John B. Birks, H., Boomer, I., Chen, J., An, C., and Wünnemann, B.: Holocene moisture evolution in arid central Asia and its out-of-phase relationship with Asian monsoon history, *Quaternary Science Reviews*, 27, 351-364, <https://doi.org/10.1016/j.quascirev.2007.10.017>, 2008.
- 605 De Grave, J., Glorie, S., Buslov, M. M., Izmer, A., Fournier-Carrie, A., Batalev, V. Y., Vanhaecke, F., Elburg, M., and Van den haute, P.: The thermo-tectonic history of the Song-Kul plateau, Kyrgyz Tien Shan: Constraints by apatite and titanite thermochronometry and zircon U/Pb dating, *Gondwana Research*, 20, 745-763, <https://doi.org/10.1016/j.gr.2011.03.011>, 2011.
- Esper, J., Shiyatov, S. G., Mazepa, V. S., Wilson, R. J. S., Graybill, D. A., and Funkhouser, G.: Temperature-sensitive Tien Shan tree ring chronologies show multi-centennial growth trends, *Climate Dynamics*, 21, 699-706, <https://doi.org/10.1007/s00382-003-0356-y>, 2003.
- 610 Ferm, B.: *Nuclear Explosions 1945-1998*, 2000.
- Fohlmeister, J., Plessen, B., Dudashvili, A. S., Tjallingii, R., Wolff, C., Gafurov, A., and Cheng, H.: Winter precipitation changes during the Medieval Climate Anomaly and the Little Ice Age in arid Central Asia, *Quaternary Science Reviews*, 178, 24-36, <https://doi.org/10.1016/j.quascirev.2017.10.026>, 2017.
- 615 Geyh, M. A., Schotterer, U., and Grosjean, M.: Temporal changes of the 14 C reservoir effect in lakes, *Radiocarbon*, 40, 921-931, <https://doi.org/10.1017/S0033822200018890>, 1997.
- Guyard, H., Chapron, E., St-Onge, G., Anselmetti, F. S., Arnaud, F., Magand, O., Francus, P., and Mélières, M.-A.: High-altitude varve records of abrupt environmental changes and mining activity over the last 4000 years in the Western French Alps (Lake Bramant, Grandes Rousses Massif), *Quaternary Science Reviews*, 26, 2644-2660, <https://doi.org/10.1016/j.quascirev.2007.07.007>, 2007.

- 620 Hall, B. L. and Henderson, G. M.: Use of uranium–thorium dating to determine past 14C reservoir effects in lakes: examples from Antarctica, *Earth and Planetary Science Letters*, 193, 565-577, [https://doi.org/10.1016/S0012-821X\(01\)00524-6](https://doi.org/10.1016/S0012-821X(01)00524-6), 2001.
- Heinecke, L., Mischke, S., Adler, K., Barth, A., Biskaborn, B. K., Plessen, B., Nitze, I., Kuhn, G., Rajabov, I., and Herzschuh, U.: Climatic and limnological changes at Lake Karakul (Tajikistan) during the last ~29 cal ka, *Journal of Paleolimnology*, 58, 317-334, <https://doi.org/10.1007/s10933-017-9980-0>, 2017.
- 625 Herzschuh, U.: Palaeo-moisture evolution in monsoonal Central Asia during the last 50,000 years, *Quaternary Science Reviews*, 25, 163-178, <https://doi.org/10.1016/j.quascirev.2005.02.006>, 2006.
- Hodell, D. A., Schelske, C. L., Fahnenstiel, G. L., and Robbins, L. L.: Biologically induced calcite and its isotopic composition in Lake Ontario, *Limnology and Oceanography*, 43, 187-199, <https://doi.org/10.4319/lo.1998.43.2.0187>, 1998.
- Hou, J., D'Andrea, W. J., and Liu, Z.: The influence of 14C reservoir age on interpretation of paleolimnological records from the Tibetan Plateau, *Quaternary Science Reviews*, 48, 67-79, <https://doi.org/10.1016/j.quascirev.2012.06.008>, 2012.
- 630 Huayu, L., Cunfa, Z., Mason, J., Shuangwen, Y., Hua, Z., Yali, Z., Junfeng, J., Swinehart, J., and Chengmin, W.: Holocene climatic changes revealed by aeolian deposits from the Qinghai Lake area (northeastern Qinghai-Tibetan Plateau) and possible forcing mechanisms, *The Holocene*, 21, 297-304, <https://doi.org/10.1177/0959683610378884>, 2010.
- Hutchinson, I., James, T. S., Reimer, P. J., Bornhold, B. D., and Clague, J. J.: Marine and limnic radiocarbon reservoir corrections for studies of late- and postglacial environments in Georgia Basin and Puget Lowland, British Columbia, Canada and Washington, USA ☆, *Quaternary Research*, 61, 193-203, <https://doi.org/10.1016/j.yqres.2003.10.004>, 2004.
- Jarvis, A.: Hole-field seamless SRTM data, International Centre for Tropical Agriculture (CIAT), <http://srtm.csi.cgiar.org>, 2008.
- Jin, L., Chen, F., Morrill, C., Otto-Bliesner, B. L., and Rosenbloom, N.: Causes of early Holocene desertification in arid central Asia, *Climate Dynamics*, 38, 1577-1591, <https://doi.org/10.1007/s00382-011-1086-1>, 2011.
- 640 Jull, A. J. T., Burr, G. S., and Hodgins, G. W. L.: Radiocarbon dating, reservoir effects, and calibration, *Quaternary International*, 299, 64-71, <https://doi.org/10.1016/j.quaint.2012.10.028>, 2013.
- Keaveney, E. M. and Reimer, P. J.: Understanding the variability in freshwater radiocarbon reservoir offsets: a cautionary tale, *Journal of Archaeological Science*, 39, 1306-1316, <https://doi.org/10.1016/j.jas.2011.12.025>, 2012.
- Kelts, K. and Hsü, K.: Freshwater carbonate sedimentation. In: *Lakes*, Springer, https://doi.org/10.1007/978-1-4757-1152-3_9, 1978.
- 645 Koppes, M., Gillespie, A. R., Burke, R. M., Thompson, S. C., and Stone, J.: Late Quaternary glaciation in the Kyrgyz Tien Shan, *Quaternary Science Reviews*, 27, 846-866, <https://doi.org/10.1016/j.quascirev.2008.01.009>, 2008.
- Kudo, A., Zheng, J., Koerner, R., Fisher, D., Santry, D., Mahara, Y., and Sugahara, M.: Global transport rates of 137Cs and 239+ 240Pu originating from the Nagasaki A-bomb in 1945 as determined from analysis of Canadian Arctic ice cores, *Journal of environmental radioactivity*, 40, 289-298, [https://doi.org/10.1016/S0265-931X\(97\)00023-4](https://doi.org/10.1016/S0265-931X(97)00023-4), 1998.
- 650 Lamoureux, S.: Varve Chronology Techniques. In: *Tracking Environmental Change Using Lake Sediments: Basin Analysis, Coring, and Chronological Techniques*, Last, W. M. and Smol, J. P. (Eds.), Springer Netherlands, Dordrecht, https://doi.org/10.1007/0-306-47669-X_11, 2001.
- Lauterbach, S., Brauer, A., Andersen, N., Danielopol, D. L., Dulski, P., Hüls, M., Milecka, K., Namiotko, T., Obremaska, M., and Von Grafenstein, U.: Environmental responses to Lateglacial climatic fluctuations recorded in the sediments of pre-Alpine Lake Mondsee (northeastern Alps), *Journal of Quaternary Science*, 26, 253-267, <https://doi.org/10.1002/jqs.1448>, 2011.
- 655 Lauterbach, S., Mingram, J., Schettler, G., and Orunbaev, S.: Two twentieth-century MLH = 7.5 earthquakes recorded in annually laminated lake sediments from Sary Chelek, western Tian Shan, Kyrgyzstan, *Quaternary Research*, 92, 288-303, <https://doi.org/10.1017/qua.2019.21>, 2019.
- Lauterbach, S., Witt, R., Plessen, B., Dulski, P., Prasad, S., Mingram, J., Gleixner, G., Hettler-Riedel, S., Stebich, M., and Schnetger, B.: Climatic imprint of the mid-latitude Westerlies in the Central Tian Shan of Kyrgyzstan and teleconnections to North Atlantic climate variability during the last 6000 years, *The Holocene*, 24, 970-984, <https://doi.org/10.1177/0959683614534741>, 2014.
- 660 Leemann, A. and Niessen, F.: Holocene glacial activity and climatic variations in the Swiss Alps: reconstructing a continuous record from proglacial lake sediments, *The Holocene*, 4, 259-268, <https://doi.org/10.1177/095968369400400305>, 1994.
- Li, Y. and Morrill, C.: Multiple factors causing Holocene lake-level change in monsoonal and arid central Asia as identified by model experiments, *Climate Dynamics*, 35, 1119-1132, <https://doi.org/10.1007/s00382-010-0861-8>, 2010.
- 665 Locket, G., Ramisch, A., Wünnemann, B., Hartmann, K., Haberzettl, T., Chen, H., and Diekmann, B.: A Process- and Provenance-Based Attempt to Unravel Inconsistent Radiocarbon Chronologies in Lake Sediments: An Example from Lake Heihai, North Tibetan Plateau (China), *Radiocarbon*, 57, 1003-1019, https://doi.org/10.2458/azu_rc.57.18221, 2016.
- Lotter, A. F. and Lemcke, G.: Methods for preparing and counting biochemical varves, *Boreas*, 28, 243-252, <https://doi.org/10.1111/j.1502-3885.1999.tb00218.x>, 1999.
- 670 MacDonald, G. M., Beukens, R. P., and Kieser, W.: Radiocarbon dating of limnic sediments: a comparative analysis and discussion, *Ecology*, 72, 1150-1155, <https://doi.org/10.2307/1940612>, 1991.
- Mathis, M., Sorrel, P., Klotz, S., Huang, X., and Oberhänsli, H.: Regional vegetation patterns at lake Son Kul reveal Holocene climatic variability in central Tien Shan (Kyrgyzstan, Central Asia), *Quaternary Science Reviews*, 89, 169-185, <https://doi.org/10.1016/j.quascirev.2014.01.023>, 2014.
- 675

- Mischke, S., Liu, C., Zhang, J., Zhang, C., Zhang, H., Jiao, P., and Plessen, B.: The world's earliest Aral-Sea type disaster: the decline of the Loulan Kingdom in the Tarim Basin, *Scientific Reports*, 7, 43102, <https://doi.org/10.1038/srep43102>, 2017.
- Mischke, S., Weynell, M., Zhang, C., and Wiechert, U.: Spatial variability of ^{14}C reservoir effects in Tibetan Plateau lakes, *Quaternary International*, 313-314, 147-155, <https://doi.org/10.1016/j.quaint.2013.01.030>, 2013.
- 680 Müller, G., Irion, G., and Förstner, U.: Formation and diagenesis of inorganic Ca-Mg carbonates in the lacustrine environment, *Naturwissenschaften*, 59, 158-164, 1972.
- Norris, R. S. and Arkin, W. M.: Known nuclear tests worldwide, 1945-98, *Bulletin of the Atomic Scientists*, 54, 65-68, 1998.
- Ojala, A. E. K., Francus, P., Zolitschka, B., Besonen, M., and Lamoureux, S. F.: Characteristics of sedimentary varve chronologies – A review, *Quaternary Science Reviews*, 43, 45-60, <https://doi.org/10.1016/j.quascirev.2012.04.006>, 2012.
- 685 Peck, J.: Mid to Late Holocene climate change in north central Mongolia as recorded in the sediments of Lake Telmen, [https://doi.org/10.1016/S0031-0182\(01\)00465-5](https://doi.org/10.1016/S0031-0182(01)00465-5), 2002.
- Potts, P. J., Thompson, M., Chenery, S. R., Webb, P. C., and Kasper, H. U.: GeoPT13-An international proficiency test for analytical geochemistry laboratories-report on round 13/July 2003 (Köln Loess), International Association of Geoanalysts, 2003.
- Ramsey, C. B.: Bayesian analysis of radiocarbon dates, *Radiocarbon*, 51, 337-360, <https://doi.org/10.1017/S0033822200033865>, 2009.
- 690 Rasmussen, K., Ricketts, R., Johnson, T., Romanovsky, V., and Grigina, O.: An 11,000-year history of Central Asian paleoclimate change recorded in deep sediments of Lake Issyk-Kul, *Kyrgyzstan, Eos Trans. AGU*, 81, 2000.
- Reference Book of Climate USSR, Kyrgyz SSR, Hydrometeo Publishing, Leningrad, 1988.
- Reimer, P. J., Bard, E., Bayliss, A., Beck, J. W., Blackwell, P. G., Ramsey, C. B., Buck, C. E., Cheng, H., Edwards, R. L., and Friedrich, M.: IntCal13 and Marine13 radiocarbon age calibration curves 0–50,000 years cal BP, *Radiocarbon*, 55, 1869-1887, https://doi.org/10.2458/azu_js_rc.55.16947, 2013.
- 695 Ricketts, R. D., Johnson, T. C., Brown, E. T., Rasmussen, K. A., and Romanovsky, V. V.: The Holocene paleolimnology of Lake Issyk-Kul, *Kyrgyzstan: trace element and stable isotope composition of ostracodes*, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 176, 207-227, [https://doi.org/10.1016/S0031-0182\(01\)00339-X](https://doi.org/10.1016/S0031-0182(01)00339-X), 2001.
- Romanovsky, V. V.: Climate, Glaciers, Lakes of Tien Shan: Journey of the past, 2007.
- 700 Romanovsky, V. V. and Shatravin, V. I.: Reconstruction of glacial and climate changes in the late Pleistocene and Holocene based on studies of closed alpine lakes in the Tien Shan and Pamir, 2007.
- Schettler, G., Mingram, J., Negendank, J. F. W., and Jiaqi, L.: Palaeovariations in the East-Asian Monsoon Regime Geochemically Recorded in Varved Sediments of Lake Sihailongwan (Northeast China, Jilin Province). Part 2: a 200-Year Record of Atmospheric Lead-210 Flux Variations and its Palaeoclimatic Implications, *Journal of Paleolimnology*, 35, 271-288, <https://doi.org/10.1007/s10933-005-0102-z>, 2006.
- 705 Schroeter, N., Lauterbach, S., Stebich, M., Kalanke, J., Mingram, J., Yildiz, C., Schouten, S., and Gleixner, G.: Biomolecular Evidence of Early Human Occupation of a High-Altitude Site in Western Central Asia During the Holocene, *Frontiers in Earth Science*, 8, <https://doi.org/10.3389/feart.2020.00020>, 2020.
- Schwarz, A., Turner, F., Lauterbach, S., Plessen, B., Krahn, K. J., Glodniok, S., Mischke, S., Stebich, M., Witt, R., and Mingram, J.: Mid-to late Holocene climate-driven regime shifts inferred from diatom, ostracod and stable isotope records from Lake Son Kol (Central Tian Shan, Kyrgyzstan), *Quaternary Science Reviews*, 177, 340-356, <https://doi.org/10.1016/j.quascirev.2017.10.009>, 2017.
- Shapley, M. D., Ito, E., & Donovan, J. J.: Authigenic calcium carbonate flux in groundwater-controlled lakes: implications for lacustrine paleoclimate records. *Geochimica et Cosmochimica Acta*, 69(10), 2517-2533, <https://doi.org/10.1016/j.gca.2004.12.001>, 2005
- Shnitnikov, A. V.: Paleolimnology of Chatyrkel Lake (Tien-Shan), *Polskie Archiwum Hydrobiologii*, 25, 1978.
- 715 Sidorov, D. A.: Two new species of freshwater amphipods (Crustacea: Gammaridae) from Central Asia, with comments on the unusual upper lip morphology, *Zootaxa*, 3317, 1-24, <http://dx.doi.org/10.11646/zootaxa.3317.1.1>, 2012.
- Taft, J. B., Phillippe, L. R., Dietrich, C. H., and Robertson, K. R.: Grassland composition, structure, and diversity patterns along major environmental gradients in the Central Tien Shan, *Plant Ecology*, 212, 1349-1361, <https://doi.org/10.1007/s11258-011-9911-5>, 2011.
- 720 Turner, T. E., Swindles, G. T., Charman, D. J., Langdon, P. G., Morris, P. J., Booth, R. K., Parry, L. E., and Nichols, J. E.: Solar cycles or random processes? Evaluating solar variability in Holocene climate records, *Sci Rep*, 6, 23961, <https://doi.org/10.1038/srep23961>, 2016.
- Wetzel, R. G.: Limnology: lake and river ecosystems, Gulf Professional Publishing, 2001.
- Williams, M. and Konovalov, V.: Central Asia temperature and precipitation data, 1879–2003, Boulder, Colorado: USA National Snow and Ice Data Center, 2008. 2008.
- 725 Wolff, C., Plessen, B., Dudashvili, A. S., Breitenbach, S. F., Cheng, H., Edwards, L. R., and Strecker, M. R.: Precipitation evolution of Central Asia during the last 5000 years, *The Holocene*, 27, 142-154, <https://doi.org/10.1177/0959683616652711>, 2017.
- Żarczyński, M., Tylmann, W., and Goslar, T.: Multiple varve chronologies for the last 2000 years from the sediments of Lake Żabińskie (northeastern Poland) – Comparison of strategies for varve counting and uncertainty estimations, *Quaternary Geochronology*, 47, 107-119, <https://doi.org/10.1016/j.quageo.2018.06.001>, 2018.

- 730 Zhou, A., Chen, F., Qiang, M., Yang, M., and Zhang, J.: The discovery of annually laminated sediments (varves) from shallow Sugan Lake
in inland arid China and their paleoclimatic significance, *Science in China Series D: Earth Sciences*, 50, 1218-1224,
<https://doi.org/10.1007/s11430-007-0081-1>, 2007.
- Zolitschka, B., Francus, P., Ojala, A. E. K., and Schimmelmann, A.: Varves in lake sediments – a review, *Quaternary Science Reviews*, 117,
1-41, <https://doi.org/10.1016/j.quascirev.2015.03.019>, 2015.

735

740

745

750

755

Tables

760

Core ID	Latitude	Longitude	Depth (m)
CHAT12	40°36.370	75°14.020	20
SC17_1	40°36'756	75°14'481	15.05
SC17_2	40°36'587	75°15'138	17.25
SC17_3	40°36'315	75°14'577	18.25
SC17_4	40°39'124	75°19'891	5-10
SC17_5	40°36'363	75°14'079	19.5-20
SC17_6	40°36'213	75°13'939	19.2
SC17_7	40°36'147	75°14'062	18.5

Table 1: Coordinates of long and short cores.

765

770

775

780


Depth (cm)	LabID	¹⁴ C	Error	cal. years BP (2σ range)	cal a BP (midpoint + span)	material
0	Poz-109830	5050	40	5908-5664	5786±122	leaves
0	Poz-54280	330	30	308-473	391±83	recent aquatic plant
0	Poz-54281	2425	25	2354 - 2692	2523±169	recent aquatic plant
0	Poz-54279	225	30	-4 - 310	155±155	recent <i>Daphnia</i>
38.5	Poz-54282	755	35	660 - 735	697±36	aquatic plant
41.5	Poz-56609	1265	30	1088 - 1285	1187±98	bulk TOC
59.5	Poz-54283	955	30	798 - 964	881±83	aquatic plant
88	Poz-54286	1595	30	1410 - 1549	1480±69	aquatic plant
98	Poz-54284	1715	35	1552 - 1706	1629±77	aquatic plant
98.7	Poz-56614	1730	30	1564 - 1708	1636±72	aquatic plant
98.7	Poz-556592	2220	30	2152 - 2324	2238±86	bulk TOC
111	Poz-54287	1925	30	1817 - 1947	1882±65	<i>Daphnia</i> remain
115.5	Poz-54288	1960	30	1830 - 1989	1910±79	<i>Daphnia</i> remain
179.5	Poz-56596	4150	35	4572 - 4827	4700±128	bulk TOC
209.7	Poz-56610	4930	35	5597 - 5727	5662±65	bulk TOC
229.5	Poz-54289	5790	50	6415 - 6665	6540±125	<i>Daphnia</i> remain
252	Poz-56595	5840	40	6533 - 6747	6640±107	bulk TOC
255.7	Poz-54290	5880	35	6637 - 6785	6659±126	<i>Daphnia</i> remain
299.7	Poz-56593	6840	40	7591 - 7757	7674±83	bulk TOC
345	Poz-56594	7610	40	8025 - 8180	8103±78	bulk TOC
345	Poz-54292	7305	35	8350 - 8514	8432±82	bulk TOC
370.1	Poz-56613	8200	50	9015 - 9300	9158±143	bulk TOC
380.5	Poz-63307	5360	40	6003 - 6277	6140±137	wood
391.4	Poz-54294	8550	50	9465 - 9604	9535±70	<i>Daphnia</i> remain
391.7	Poz-54293	8710	50	9546 - 9887	9717±171	<i>Daphnia</i> remain
437.2	Poz-54296	9160	50	10230 - 10487	10359±129	<i>Daphnia</i> remain
439.9	Poz-56611	9360	50	10427 - 10713	10570±143	bulk TOC
466	Poz-54297	9670	50	10789 - 11211	11000±212	<i>Daphnia</i> remain
469	Poz-54298	9690	50	10795 - 11226	11011±216	<i>Daphnia</i> remain
508	Poz-54299	10840	50	12681 - 12804	12743±62	<i>Ruppia maritima</i>
510	Poz-54300	11060	50	12790 - 13062	12926±136	bulk TOC
528	Poz-54301	12150	50	13831 - 14175	14003±172	bulk TOC
528	Poz-54302	8890	50	9785 - 10191	9988±203	deciduous wood
549.5	Poz-56591	12820	60	15105 - 15550	15328±223	bulk TOC
571	Poz-56608	13220	70	15660 - 16125	15893±233	bulk TOC
585	Poz-63308	14060	90	16759 - 17419	17089±330	<i>Ruppia maritima</i>
620.5	Poz56590	13190	70	15612- 16092	15852±240	bulk TOC

Table 2: ¹⁴C dates (calibrated with OxCal 4.3, IntCal13).

785


790

Figures



800

Figure 1: Location of Lake Chatyr Kul, the composite profile (red dot) and the gravity cores (yellow dots). The orange square marks the location of ^{14}C dated leaves (Poz-109830, Tab. 2) found in the top of a mid-Holocene-shoreline at ~3540 m a.s.l. The relief map of Kyrgyzstan relies on the CGIAR-CSI SRTM 90m (3 arcsec) digital elevation data (Version 4) of the NASA Shuttle Radar Topography Mission (Jarvis, 2008). The figure was modified from Lauterbach et al. (2014). Photo of instable shores (white arrow) of Maloye lake.




805 **Figure 2: Oxygen concentration (YSI Pro 6600 V2), Temperature and specific conductivity measured with a CTD sensor during the field trip in 2012 at the core's location (N 40°36.371', E 75°14.006').**

810

815

820




825 **Figure 3: The composite profile CHAT12 (dark grey: piston cores A5o, A5u, B3o, B1u and A4u, used depth sections are displayed within). Additional gravity cores taken in 2017 (SC17_1 to SC17_7) were only partly used for thin section preparation and gamma spectrometry dating. The gravity cores cover approximately the upper first meter of the composite profile.**

830

835

varve deposition models


thin section pictures of dominant varve types in the different lithozones



homogenous sediments


with faint laminac in the upper 1.5 cm depth (grey bar)

clastic-organic laminations




Winter amorphous organic layer
Summer runoff event (clastic varves in LZ V)
Summer mixed layer containing fine detrital grains and endogenic calcite
Spring coarse detrital runoff layer often with pennate diatoms and chrysophytes

organic-clastic laminations



Winter amorphous organic layer
Summer mixed layer containing detrital grains, rich in aquatic plant remains & periphytic diatom *Achnanthes brevipes*, daphnia, ostracods, characeae
Spring coarse detrital runoff layer often with pennate diatoms and chrysophytes

clastic-calcitic laminations



Winter amorphous organic layer
Summer mixed layer of endogenic calcite formation and fine detrital grains
Spring coarse detrital runoff layer often with pennate diatoms and chrysophytes

clastic-diatom laminations



Summer/Autumn *Cyclotella choctawhatcheeana* blooms
Summer mixed layer
Spring coarse detrital runoff layer often with pennate diatoms and chrysophytes

calcitic-clastic laminations



Winter amorphous organic layer
Summer layer of intensive endogenic calcite formation with scattered detrital grains
Spring coarse detrital runoff layer

elastic-aragonitic laminations



Summer mixed layer of fine detritus and endogenic calcite with occasionally aragonite formation
Spring coarse detrital runoff layer rich in pyrite

◆ Pyrite ◊ Aragonite ■ endogenic calcite ■ amorphous organic

● Coarse and fine detrital grains (qz, fsp, ca, mica) ■ mixture of detrital and endogenic calcite ■ mixture of fine detritus and organics

● planktic diatom *Cyclotella choctawhatcheeana* ■ periphytic diatom *Achnanthes brevipes* ○ chrysophytes ~ aquatic plant remains





Figure 4:1 Thin section pictures of different lamination (varve) types in cross-polarized (left) and plane polarized (right) light of the different lithozones LZ I to LZ V. **a)** Clastic-organic laminations; **b)** Intercalation of clastic-organic, clastic-diatom and calcitic-clastic laminations; **c)** Intercalation of clastic-calcitic, clastic-organic and clastic-diatom laminations (upper part); **d)** Organic-clastic laminations; **e)** Clastic-organic laminations; **f)** homogenous sediments. Black/white/grey bars alongside the thin section pictures indicate one individual varve, but the reliability of grey bars (varves) is being generally lower due to high amounts of aquatic plant remains and low preservation. Process-related deposition Models of the observed lamination/varve types illustrate the seasonal depositional successions (a-e).




840 Fig. 4.2 XRF-map 1: Varve types and XRF element mapping of a thin section from LZII from 507 to 497.5 cm depth. Element maps show an alternation of siliciclastic sediments with high amounts of Si and Al with calcite layers (Ca) with high amounts of Sr and Mg according to the presence of clastic-organic, clastic-diatom and calcitic-clastic and calcitic-clastic varves respectively. Microfacies analyses show endogenic calcite within calcitic-clastic varves within the summer sublayer (image a). Clastic-organic and clastic-diatom varves are indicated by high amounts of Si and Al with only individual calcite (arrow image b) and Mg-calcite grains.


845 Note, that coinciding occurrences of individual elements results in colour mixing e.g. Al (red) and Si (blue) becomes pink. Endmembers of the Al-Si map are indicative for the presence of Al-rich clays and diatomaceous Si.

850


138 ± 6
counted varves



870 Fig. 4.3 XRF-map 2: Varve types and XRF element mapping of a thin section from LZIII from 357 to 338.5 cm depth. Element maps
 875 show an alternation of siliciclastic sediments with high amounts of Si and Al with calcitic layers (Ca) with high amounts of Sr according to the presence of clastic-organic, clastic-diatom and clastic-calcitic varves respectively. Micro-facies analyses show mixed calcitic (resuspended and endogenic) summer sublayers within clastic-calcitic varves (image a). Clastic-organic and clastic-diatom varves are indicated by high amounts of Si and Al with only individual calcite grains (arrow image b). Note, that coinciding occurrences of individual elements results in colour mixing e.g. Al (red) and Si (blue) becomes pink. Endmembers of the Al-Si map are indicative for the presence of Al-rich clays and diatomaceous Si (blue).



880 **Figure 5:** a) Varve counting uncertainty estimates (mean of 5% =green) and VQI distribution. b) Age model of the floating varve chronology (Chatvd19) from 63.0-623.5 cm depth with a basal age of 11619 ± 603 years BP (light green). The black line shows the measured varve thickness, black dots mark the distribution of calibrated AMS ¹⁴C ages (with 2σ uncertainty) of wood pieces (Tab.2). b) ²¹⁰Pb CIC (grey squares) and CRS (black dots) age models, ¹³⁷Cs activity concentration profile and constrained age model for the uppermost part of the composite profile (green triangles).



890 **Figure 6: Gamma spectrometry results of the gravity core SC17_7. Light grey intervals indicate uncorrelated sequences of the $\ln(^{210}\text{Pb}_{\text{unsupp}})$ vs. depth profile which affected the CIC model calculations (Suppl. Fig.2, Suppl. Tab. 2, 3). Core pictures of the upper part of the composite profile CHAT12 (right) and the gravity core SC17_7 (left) illustrate the facies change to calcite-enriched sediments in the uppermost centimeter.**

895

900




Figure 7: Radiocarbon reservoir effect (grey step plot). The reservoir effect was determined by the difference of aquatic cal. a BP (colored symbols) and varve ages (green). Floating varve chronology (green) and distribution of calibrated AMS ¹⁴C ages (with 2σ uncertainty) of wood pieces, TOC bulk, aquatic plant remains, daphnia and *Ruppia maritima* remains (Tab.2).




Figure 8: VQI, Counting differences for individual thin sections (green= 1st count, black = 2nd count, and their percentage deviation).






Figure 9: Holocene seasonal deposition patterns, semi-quantitative species assemblages and calibrated ^{14}C dates of paleo-shore deposits (black dots from Shnitnikov (1978), yellow square from own sample taken in 2017 (Table 2). Percentage distribution of varve types in the different lithozones.