

Towards *in-situ* U–Pb dating of dolomites

Bar Elisha^{1,3}, Perach Nuriel¹, Andrew Kylander-Clark², Ram Weinberger^{1,3}

¹Geological Survey of Israel, Jerusalem, Israel

²Department of Earth Sciences, University of California, Santa Barbara, CA, USA

5 ³Department of Earth and Environmental Sciences, Ben-Gurion University, Be'er Sheva, Israel

Correspondence to: Bar Elisha (brelisha@bgu.ac.il)

Abstract. Recent U–Pb dating by laser ablation ICP-MS has demonstrated that reasonable precision (3–10%, 2 σ) can be achieved for high-resolution dating of texturally distinct calcite phases. Absolute dating of dolomite, for which biostratigraphy and traditional dating techniques are very limited, remains challenging although it may resolve many fundamental questions related to the timing of mineral-rock formation by syngenetic, diagenesis, hydrothermal, and epigenetic processes. In this study we explore the possibility of dating dolomitic rocks via recent LA-ICP-MS dating techniques developed for calcite. The *in-situ* U–Pb dating was tested on a range of dolomitic rocks of various origins (i.e., syngenetic, early diagenetic and epigenetic) from the Cambrian to Pliocene age—all of which from well-constrained stratigraphic sections in Israel. We present *in-situ* U–Pb results of dolomitic rock samples, together with imaging techniques and chemical characterizations. **We show the complexity of in-situ dolomite dating and discuss variables such as crater morphology, textural features, chemical and age zoning and detrital impurities that may affect the interpretation of the resulted ages.** Textural examination indicates zonation and mixing of different phases at the sub-millimetre scale (<1 μm), and thus Tera-Wasserburg ages may represent mixing dates of early diagenesis and some later epigenetic dolomitization event(s). **We conclude that age mixing at the sub-millimetre scale is a major challenge in dolomite dating that need to be further studied. We also note the importance of matrix-matched standards for reducing uncertainties of the dated material.**

10
15
20

1 Introduction

Dolomite is vastly abundant in exposed stratigraphic sequences, and its manifestation in the geological record increases towards older sedimentary strata (Warren, 2000). Nonetheless, it is very rare in modern environments and has seldom been successfully grown in laboratory experiments at near-surface conditions (Machel, 2004, and references therein). Although the conditions and kinetics promoting dolomite growth are not well understood, its formation is considered as a by-product of a chemical reaction between Mg-rich fluids and calcite-bearing rocks. Previous studies suggested that dolomite is being formed either by diagenetic replacement of limestone during deposition (syngenetic; Sass, 1969), soon after deposition (early diagenetic; Ahm et al., 2018; Frisia et al., 2018), or at a later stage (epigenetic; Sibley and Gregg, 1984). Distinguishing between different dolomitization processes is challenging, yet critical for resolving some of the issues and ambiguities related to the formation of dolomitic rocks. Accurate U–Pb absolute dating of dolomite by laser ablation inductively coupled plasma

25
30

mass spectrometry (LA-ICP-MS) could contribute to better understanding of dolomitization process by placing these event(s) in the proper geological context. Previous U–Pb dating of dolomites on whole-rock samples of U-rich dolostones, conducted in the highest level of cleanroom standards, yielded scattered results along the isochron (Winter and Johnson, 1995; Hoff et al., 1995; Ovchinnikova et al., 2007; Polyak et al., 2016). These studies suggested that *in-situ* dating of dolomites should be
35 feasible, and indeed several studies recently reported on successful *in-situ* age determination of dolomites using the LA-ICP-MS methodology (Burisch et al., 2018; Salih et al., 2019; Hu et al., 2020; Incepri et al., 2020; Mueller et al., 2020).

Recent developments of LA-ICP-MS has opened a new avenue for measuring absolute ages of carbonates, thus improving the understanding of many fundamental geological processes, such as fossilization (Li et al., 2014), tectonic faulting (Ring et al., 2016; Roberts and Walker., 2016; Nuriel et al., 2017; Parrish et al., 2018), duration of sedimentation, and diagenesis (Hodson
40 et al., 2016; Godeau et al., 2018). Despite the low concentrations of U and radiogenic Pb in carbonates (<10 ppm and <2 ppm, respectively) and the considerable amounts of common Pb (up to 100 ppm), a reliable age determination of calcite is obtained via isochron regression on a Tera-Wasserburg inverse concordia diagram (Tera and Wasserburg, 1972). By this method, the common-Pb composition and the age are determined by the upper and lower intercept of the regression isochron with the concordia curve. While LA-ICP-MS analyses on calcite evolved to be a conventional method of dating (Roberts et al., this
45 issue), a thorough methodology for dating other carbonates, such as dolomite, is still needed (Guillong et al., this issue).

Dating dolomitic rocks is more challenging than calcite, particularly because their complicated growth history is often characterized by the formation of multi-phase microcrystalline grains (e.g. partial replacement, zoning). Growth-zones cannot be separated physically, and their size is often smaller than the diameter of the laser spot (usually >50 μm). In addition, well-characterized dolomite reference materials (RM) are currently unavailable for the LA community and differences between
50 calcite and dolomite in terms of matrix-effect and plasma efficiency are not well understood (Guillong et al., this issue). In order to examine the effect of using common RM and the suitability of conventional LA-ICP-MS calcite procedure for dolomite geochronology, we studied dolomitic rock samples with well-defined stratigraphic ages. We show differences in texture, crater morphology, impurities, and down-hole fractionation trends, between RMs and dolomite and discuss textural characteristics and chemical properties of successful and unsuccessful dolomite dating. Finally, we consider the age results in
55 the geological context of the studied rocks.

2 Methods

For LA-ICP-MS analyses of dolomites we prepared 40 μm thick thin sections polished to 1 μm . U–Pb LA-ICP-MS analyses were performed at the Department of Earth Science, University of California, Santa Barbara, following the analytical procedure described in Nuriel et al., (2017) for calcite-bearing rocks. Samples were ablated using a *Photon Machines* 193 nm ArF
60 Excimer laser equipped with a HelEx ablation cell and coupled to a *Nu Instruments Plasma 3D* multi-collector ICP-MS. Both RMs and unknowns were ablated with similar spot size of 85 μm and fluence of $\sim 1 \text{ J/cm}^2$. In order to remove any contaminants, and especially common Pb from the sample surface, all samples were cleaned with methanol and pre-ablated (4 pulses) prior

to a 20 s baseline. Material was then ablated for 15 seconds at 10 Hz, resulting in a pit depth of ~15 μm . On the MC-ICP-MS, masses ^{202}Hg , ^{204}Pb , ^{206}Pb , ^{207}Pb , and ^{208}Pb were measured on Daly detectors, and masses ^{232}Th and ^{238}U were measured on Faraday detectors at low resolution (300, 10% valley definition) using an integration time of 100 ms. We used a two-steps standardization technique using NIST614 glass and the WC-1 calcite reference material (Roberts et al., 2017) following the procedure outlined in Nuriel et al. (2017). Data were reduced using Iolite v. 2.5 (Paton et al., 2010) and the $^{238}\text{U}/^{206}\text{Pb}$ and $^{207}\text{Pb}/^{206}\text{Pb}$ isotopic ratios for each analysis were plotted on Tera–Wasserburg diagrams using Isoplot and IsoplotR (Ludwig, 2012; Vermeesch, 2018); U and Pb concentrations were calculated semi-quantitatively, using NIST614 as the primary reference material (RM). **Uncertainties were propagated on individual unknown ratios such that $^{207}\text{Pb}/^{206}\text{Pb}$ (2%) and $^{206}\text{Pb}/^{238}\text{U}$ (4%) ratios of a zircon standard, run throughout the session (Mud Tank; Black and Gulson, 1978), yielded a single population; this resulted in reasonable mean square weighted deviations (MSWDs) for the calculated ages of calcite RMs.** Secondary calcite RMs—ASH-15 (2.9646 ± 0.011 Ma; Nuriel et al., this issue) and Duff Brown (64 ± 0.67 Ma; Hill et al., 2016)—yielded dates within uncertainty of their accepted values (ASH-15: 2.973 ± 0.090 MSWD = 1.3, n = 107; Duff Brown: 63.2 ± 2.3 Ma, MSWD = 1.9, n = 106). Uncertainty correlations are calculated following Schimdtz and Schoene, 2007. The Pb concentration for each spot analysis was calculated by the total counts of Pb isotopes, compared to the NIST glass value (2.32 ppm). **The ^{204}Pb concentration was calculated using the ^{206}Pb concentration and assuming a Stacy-Kramers $^{206}\text{Pb}/^{204}\text{Pb}$ ratio to avoid difficulties related to the Hg interference on ^{204}Pb .**

Following LA analyses, we used several techniques to characterize the studied dolomite samples in detail. Whole-rock analyses of Rare Earth Element (REE) composition was done on Perkin Elmer NexION 300D ICP-MS instrument. Dolomite powders were dissolved, evaporated, and diluted ~3000 in 0.1N nitric acid solution before mixed with Rh/Rn internal standards. The raw data were corrected for blank, drift and isobaric interferences and converted into concentrations in ppm using USGS RM. The overall uncertainties are estimated to be less than 5%.

Imaging of the LA craters and identifying major phases in the samples was performed by using a field-emission FEI Scanning Electron Microscope (SEM) at the Ilse Katz Institute for Nanoscale Science & Technology at Ben-Gurion University of the Negev, Israel, with 3 kV acceleration voltage, 0.1 nA current and 30° stage tilt. This device is equipped with EDS detector and ‘Oxford’ EBSD (Electron backscatter diffraction) sensor, used for producing crystallographic phase maps. For EBSD mapping, the instrument was setup to 15 kV accelerating voltage and 26 nA current, 70° tilt, 2x2 binning and 0.1 μm step size. Wave Dispersion Spectroscopy (WDS) maps were preformed using a JEOL microprobe at the Hebrew University, Israel, with accelerating voltage of 15–25 kV, beam current of 80 nA, step size of 0.5 μm and dwell time of 0.35 s.

X-ray diffraction (XRD) patterns were acquired in Bragg-Brentano geometry at the Geological Survey of Israel using a PANalytical X’Pert diffractometer with $\text{CuK}\alpha$ radiation operated at 45 kV and 40 mA. Samples were scanned from 3 to 70° 2 θ at a step size of 0.013° 2 θ , using a PIXcel detector in continuous scanning line (1D) mode with an active length of 3.35°. The equivalent time per step was ~30 sec, resulting in a total measurement time of about 10 min per scan. Mineral phase identification and semi-quantification was performed using HighScore Plus® software based on ICSD database.

2.1 Studied dolomites

Dolomitic rocks in Israel and environs include syngenetic to early diagenetic dolomites, epigenetic dolomites, hydrothermal dolomites and mixed/hybrid ones, whereas their ages are well constrained by field relations and dates of adjacent geological units (Fig. 1). Thin section scans and representative photomicrographs of each studied sample are provided in Fig. 1 and are described in the following sections. Cathodoluminescence images of representative carbonate material, used to infer slight changes in fluid composition (e.g. Mn^{2+} , Fe^{2+} content), and/or precipitation conditions, are presented in Fig. 2.




Figure 1: Thin-sections scans and representative photomicrographs of each dolomitic sample from this study. Red dots on thin-section scans showing the locations of LA-ICP-MS analyses. Width of thin-sections are 27 mm. Sample locations along the stratigraphic column of Israel is also provided in the right panel.

2.1.1 Syngenetic Cambrian dolomites and hydrothermal dolomites (Timna Valley)

Cambrian sediments are exposed in southern Israel and unconformably overlie Precambrian crystalline basement rocks of the Arabian-Nubian Shield (Fig. 1; Beyth et al., 1999). In the Timna Valley, southern Israel, Cambrian dolomitic rocks of the Timna Formation are well-known for their copper deposits and ancient to present-day mining and are considered to be formed as early diagenetic in a marine environment at 25-50 °C (Segev, 2016). Based on fluid inclusions and petrographic studies,

Eliyahu et al. (2017) suggested that the Timna formation dolomites were formed in high temperatures and the dolomites are epigenetic in nature. Dolomitic rocks of the Timna Formation (sample Tm-MU-2; Table 1) represent the earliest oceanic transgression in the area, constrained by trilobite burrowing to upper Georgian (~520 Ma; Parnes, 1971) and by a dike intrusion dated to ~532 Ma (Beyth and Heimann, 1999). Sample Tm-MU-2 (Fig. 1) is composed of reddish sparry dolomite of fine grains <10 μm in size, with minor iron-oxides scattered within the sample. Dolomite veins of later epigenetic diagenesis (Sample Tm-DV-1; Fig. 1) are found in the crystalline basement rock and sandstones in Timna Valley, in association with copper, quartz, calcite and Mn and Cu-carbonates. Sample Tm-DV-1 is composed of euhedral zoned dolomite crystals of up to 200 μm , with opaque cores and transparent rims. It was previously suggested that these euhedral dolomite crystals of epigenetic open-space filling cements, associated with Cu mineralization, are related to low-temperature (~260 $^{\circ}\text{C}$; Beyth et al., 1997) hydrothermal activity and mineralization that assumed to occur during Neogene Times (Kohn et al., 2019). On the other hand, Eliyahu et al. (2017) suggested that all Cu mineralization in the Timna valley is associated with epigenetic hydrothermal dolomite mineralization, driven by basinal fluids. The zoned hydrothermal dolomite grains of sample Tm-DV-1 are slightly zoned under CL (Fig. 2) with very similar luminescence, suggesting minimal changes in fluid composition and/or precipitation conditions.


2.1.2 Syngenetic and early diagenetic dolomites (Mount Carmel and Umm el Fahm Ridge)

Dolomitic rocks dominate the exposed Cretaceous sequence of Mount Carmel, Umm el Fahm Ridge and Judean Mountains, which were part of an extensive shallow carbonate platform. The studied Cenomanian dolomitic rocks of the Deir Hanna Formation (Fig. 1) are exposed on the SE flank of the Umm el Fahm anticline near the village of Mei-Ami (Sass et al., 2013). These rocks are underlain and overlain by volcanic flows that are dated to 99 ± 0.5 and 95 ± 0.5 Ma, respectively (Ar–Ar; Segev et al., 2002). They were described as syngenetic dolomites based on preferred orientations of dolomite grains, with a c-axis maximum perpendicular to the bedding planes (Sass, 1969). Samples MAM-3 and MAM-7 (Fig. 1) are composed of fine-grained (<10 μm) micritic dolomite, which reflect continuity of reefs along fine-grained, well-bedded shelf basin rocks (Sass and Bein, 1978).

Dolomitic rocks of the Zikhron Formation from Mount Carmel are considered ‘early diagenetic’ (Sass and Bein, 1978; Segev and Sass, 2009; Fig. 1) and crop out between two volcanic flows of 97 ± 0.5 and 95 ± 0.5 Ma (Segev, 2009). Samples MU-1 and MU-2 (Fig. 1) are composed of ~40 μm dolomite grains and represent sparry dolomite mosaic of similar ages as MAM-3 and MAM-7. Dolomitic rocks from the Albian Yagur Formation crop out near the Kerem Maharal village and are overlain by the oldest (99 Ma) volcanic flow known in Mount Carmel. Samples KM-1 (Fig. 1) is a sparry dolomite with ~60 μm dolomite grains and is considered as an ‘early diagenetic’ dolomite. The non-homogenized luminescence of the sparry sample KM-1 (Fig. 2) may indicate a possible mixture of phases that precipitated under different conditions.

2.1.3 Fault-related Epigenetic dolomitization of early diagenetic dolomites – (Judean Desert)

145 Strata of dolomitic rocks are abundant at the western margin of the Dead Sea basin and include the Cenomanian Hevion, Zafit and Tamar formations (Sneh and Avni, 2016). These dolomitic rocks are considered 'early diagenetic' dolomites that were later faulted and cemented by epigenetic dolomite during the activity along the Dead Sea fault. Dolomite-cemented breccias were sampled along one of the major faults of the Dead Sea western margin fault zone (En Feshkha Fault; sample EFN-1; Fig. 1) and preserve microstructures of mosaic (sparry) dolomite fragments bounded by sparry dolomite cement. The bright luminescence of the cement material in sample EFN-1 suggest a single phase of precipitation that is distinctively different from precipitation conditions of the fragment material.



150 **Figure 2.** PPL images (left panels) and Cathodoluminescence (CL) images (right panels) of representative studied samples. Note the differences in CL colors of breccia fragments and cement in Sample EFN-1, the non-homogenize CL response in sample KM-1 and the slight zoned dolomite crystals in sample Tm-DV-1.

3 Results

We present U–Pb ages of eight dolomite samples (Table 1), and a ‘Tera-Wasserburg’ inverse concordia diagrams as $^{207}\text{Pb}/^{206}\text{Pb}$ and $^{238}\text{U}/^{206}\text{Pb}$ linear regression isochrons of these samples (Fig. 3). The $^{207}\text{Pb}/^{206}\text{Pb}$ (Common Pb) values were not anchored to specific values and range between 0.7862 ± 0.0033 and 0.9683 ± 0.0071 . MSWD values are between 0.27 and 41. U–Pb isotopic ratios analyses of the syngenetic Cambrian dolomite Tm-MU-2 forms an isochron that intercept at 277 ± 59 (MSWD = 0.53; n = 80; Fig. 3A), with a common Pb value of 0.8664 ± 0.0061 . The U–Pb data of the hydrothermal dolomite sample Tm-DV-1 shows lower intercept age of 37 ± 75 (MSWD = 0.27; n = 70; Fig. 3B) and common Pb value of 0.8385 ± 0.0038 with similar pattern to sample Tm-MU-2. Syngenetic Cretaceous dolomite samples MAM-3 and MAM-7 yielded lower intercept age of 137 ± 14 (MSWD = 0.63; n = 80) and 170 ± 11 (MSWD = 1.6; n = 80) and common Pb value of 0.7899 ± 0.0047 and 0.8427 ± 0.0032 , respectively (Fig. 3C–D). Sample MU-1 of Cenomanian stratigraphic age yielded intercept age of 58 ± 5 Ma and common Pb value of 0.8046 ± 0.0036 (MSWD = 5.4; n = 80; Fig. 3E). The U–Pb ages of the sparry sample MU-2 yielded a lower intercept age of 93 ± 7 and common Pb value of 0.8064 ± 0.004 (MSWD = 2.4; n = 70; Fig. 3F). Sparry sample KM-1 shows lower intercept age of 55 ± 6 (MSWD = 1.3; n = 80; Fig. 3G) and common Pb value of 0.7862 ± 0.033 . Analyses of the fault related dolomite sample EFN-1 was performed on both the homogeneous sparry dolomitic fragments (n = 30) and on the fine grained enclosing dolomitic cement between fragments (n = 50). All 80 spot analyses in this sample yielded age of 51 ± 1 Ma (MSWD = 41; n = 80) and upper intercept value of 0.9683 ± 0.0071 (Fig. 3H).

Sample	Latitude	Longitude	Petrographic description	Stratigraphic age	U–Pb age	MSWD	Common Pb value
Tm-MU-2	29.471561	34.591990	Micritic	Cambrian	277 ± 59	0.53	0.8664 ± 0.0061
Tm-DV-1	29.778567	34.977663	Hydrothermal	Miocene	37 ± 75	0.27	0.8385 ± 0.0038
MAM-3	32.504359	35.141783	Micritic	Cretaceous	137 ± 14	0.63	0.7899 ± 0.0047
MAM-7	32.496742	35.150912	Micritic	Cretaceous	173 ± 11	1.6	0.8427 ± 0.0032
MU-1	32.671247	35.092221	Sparry	Cretaceous	58 ± 5	5.4	0.8046 ± 0.0036
MU-2	32.671516	35.092353	Sparry	Cretaceous	93 ± 7	2.4	0.8064 ± 0.004
KM-1	32.641733	34.981380	Sparry	Cretaceous	55 ± 6	1.3	0.7862 ± 0.33
EFN-1 fragments	31.716563	35.448523	Sparry	Unknown	74 ± 3	0.99	0.8140 ± 0.021
EFN-1 cement	31.716563	35.448523	Sparry	Unknown	6.5 ± 1	1.5	0.7853 ± 0.0047

170 **Table 1: Sample description and stratigraphic age and their corresponding LA results.**




Figure 3: Tera-Wasserburg concordia plots of studied dolomites: syngenetic Cambrian dolomites (A), hydrothermal dolomites (B), syngenetic Cretaceous dolomites (C–D), early diagenetic dolomites (E–G) and epigenetic dolomite (H). All diagrams have similar axes. Isochrons and uncertainties are presented as black lines and gray areas, respectively. Insets show enlargements of ellipse concentration areas. Uncertainty ellipsoids of spot analysis are plotted in green and represent 2σ uncertainties.

4 Discussion

4.1 Significance of the U–Pb ages

180 The expected stratigraphic ages of most studied dolomites in this study are inconsistent with their U–Pb ages. In order to further study the significance of the resulted ages we discuss the geochemistry of each sample such as the U and Pb concentration of each spot analysis, together with the total U and Pb content of each sample (Fig. 4A). We also present whole-rock REE content and discuss the relation between its trace elements signature and the resulted U–Pb ages (Fig. 4B).

185 Sample Tm-MU-2 was assumed to produce Cambrian age (~520 Ma) but yielded 277 ± 59 Ma, ~180 Ma younger than expected. Uncertainty ellipses of this sample are plotted near the common Pb value, therefore the lower intercept is far projected and poorly constrained (Fig. 3A). U and Pb concentration of this sample are plotted in the upper-left quadrant of figure 4A, with low U (~0.2 ppm) and high Pb contents (~5 ppm). The common Pb value in this sample (0.8664 ± 0.0061) may represent incorporation of radiogenic Pb derived from the surrounding crystalline rocks, as expected for carbonates associated with hydrothermal activity (Stacey and Kramers 1975). This is also supported by the REE signature of sample Tm-MU-2, showing elevated LREE, depleted HREE and positive Gd anomaly (Figure 4B). The REE pattern of this sample is similar to other dolomites in this study, although one order of magnitude higher. Sample Tm-DV-1 display similar pattern to those of sample Tm-MU-2, with uncertainty ellipses near the common-Pb intercept (Fig. 3B) and with low U (~0.2 ppm) and high Pb contents (~5 ppm) of individual spot analyses plotted in the upper-left quadrant in figure 4A. These patterns suggest that dolomitic rocks associated with hydrothermal activity are most likely to contain high common Pb concentrations and are specified here as dolomites with low-chances for successful dating.

195 The stratigraphic age of samples MAM-3 and MAM-7 was constrained to 99 and 95.4 Ma (Segev et al., 2002). However, their U–Pb ages yielded ‘small scale isochron’ (Ring and Gerdes, 2016) with 137 ± 14 and 170 ± 11 intercepts, respectively, 40% to 70% older than expected. Although the low $^{207}\text{Pb}/^{206}\text{Pb}$ value of 0.7899 in sample MAM-3 indicate higher incorporation of radiogenic-Pb during dolomitization compared to sample MAM-7 (0.8427 ± 0.0032), MAM-7 displays much larger age offset than MAM-3. In these samples U and Pb contents plot close to 1 ppm U but their total Pb content is up to 20 ppm, forming a cluster above the center of the diagram in figure 4A. We suggest that dolomites with similar U and Pb contents can also be classified as low-chances for successful dating.

205 The isochrone of sample MU-1 was expected to produce Cenomanian age, but its isochrone intercepts at 58 ± 5 Ma, ~40 Ma younger than expected. On the other hand, sample MU-2 was collected several meters away and produced age of 93 ± 7 Ma. This age is within the uncertainty of the 95–97 Ma Ar–Ar ages of the constraining volcanic layers. The U content of these samples is between 0.5 to 2 ppm and Pb content is between <0.1 and 4 ppm, forming a cluster around the center of the diagram in figure 4A. Sample KM-1 is constrained stratigraphically to 99 Ma, however yielded ~50% younger age than expected. Its isochrone shows similar age pattern to sample MU-1, with lower intercept age of 55 ± 6 . The REE signature of the above three samples are rather similar, with slightly elevated LREE and similar Gd anomaly (Fig. 4B). In sample EFN-1 the spot analyses

210 are clearly a mix of two different phases as the ellipses are arrayed along two isochrons. The results of this sample are further discussed in more detail.





Figure 4: (A) U vs. Pb [ppm] of single spots analyses by LA-ICP-MS of studied dolomite samples, together with whole rock U and Pb content of each sample (large circles). (B) Corresponding whole-rock REE patterns normalized to chondrite values.

215 4.2 Textural characteristics of analyzed dolomites

It was previously suggested that differences in crater morphology between calcite and dolomite may cause mass fractionation due to uneven mass removal between dolomites and the calcite standard WC-1 and that 160% difference in ablation efficiency in micritic dolomite may result in age offset of 4–8% (Guillong et al., this issue). In order to test whether the discrepancy in ages of the studied dolomites is due to imperfections in crater rims and bottom we imaged individual laser pits and point out and discuss few differences in their crater morphologies (Fig. 5). Although we did find some differences, none of them are sufficient enough to explain the large offsets between expected and obtained dates by differences in ablation efficiency.

220 Crater morphology in sapric dolomite samples with crystal size >10 μm is consistently similar, with only minor roughness on crater bottom and rims (Fig. 5). In sample MU-2 the age is consistent with its stratigraphic appearance, but the morphology of its crater suggests that the age of MU-2 and the younger age of MU-1 probably represent actual diagenesis/dolomitization processes. Laser craters of samples Tm-MU-2 and Tm-DV-1 has moderate roughness along the bottom with minor imperfections along crater rims (Fig. 5). The poorly constrained ages of these samples, therefore, seem to relate more to their trace-element chemistry rather than differences in crater morphology. On the other hand, the morphology of laser pits in micritic dolomites samples (MAM-3 and MAM-7) display multiple imperfections along crater bottom and rims compared to sparry dolomites (Fig. 5). It might be possible that these morphological differences effect at some extent the 40–70% age offset obtained. In conclusion, we suggest that differences in ablation efficiency can have only minor effect on the resulted ages and other parameters should be considered.



235 **Figure 5: Ablation craters of studied samples, arranged by crater geometry and bottom roughness, from smooth (MU-2, sparic dolomite), via moderate (TM-MU-2, sparic dolomite), to rough (MAM-7, micritic dolomite). Imperfections along crater rims are marked by white arrows.**

Panchromatic back-scattered electron (BSE) images of representative samples show that intracrystalline porosity, distribution of grain size, tiling pattern and type of mineral zoning of dolomite rhombs are much more significant parameters to be considered (Fig. 6). Intracrystalline porosity are usually smaller than spot size of 85 µm, and may include other phases beside dolomite, such as k-feldspar, pyrite, oxides and bituminous minerals (Fig. 6A; Olanipekun and Azmy, 2017). These phases may include detrital contaminations with inherited U–Pb ages, which might lead to mixed results and ages older than expected. Aside from external impurities, crystals smaller than the spot size (85 µm), along with zoning in the crystals can also lead to mixed results. Except for sample Tm-DV-1, where dolomite grains reach 200 µm, analyses of a single crystal in the studied samples is impossible. In sample MU-1 the longest diagonal of dolomite crystals is ~60 µm. Dolomite cores of MU-1 appear much brighter in BSE compared to their concentric enclosing rims, probably due to higher Mg/Ca ratio and minor concentration of Fe (Fig. 6A; Olanipekun and Azmy, 2017). Dolomite crystals from sample KM-1 display mainly a concentric zoning pattern with a very thin lamina separating the core from the rim. Abundant disseminated calcite inclusions are found in the cores but have relatively homogeneous rim sections (Fig. 6B). Such signature is likely to be associated with the mechanism of epigenetic dolomitization governed by diagenetic replacement of pore fluids and re-precipitation of dolomite (Putnis & Putnis, 2007; Olanipekun & Azmy, 2017). Within the large fragments of breccia sample EFN-1, a mixture of different zoning patterns can be seen: dolomite crystals that lack distinctive core to rim zones and crystals with two zones of bright cores and dark rims (Fig. 6C). The cement between the large fragments in this sample contain <50 µm isolated fragments of broken dolomite crystals embedded in homogeneous cement with bright BSE response (Fig. 6D). The BSE images highlight complexities in the chemical zoning of dolomite at sub-millimeter scale, i.e. distinct core and rim, semi-homogenized grains, or mixture of different grains. It is therefore important to acknowledge that spot analysis of these textures may result in age-mixing or averaging of the different phases.

240
245
250
255





Figure 6: Panchromatic BSE images of samples MU-1 (A), KM-1 (B) and fragments and cement of sample EFN-1 (C and D, respectively). BSE images are efficient in revealing the grain size of the sample, as well as porosity and additional intracrystalline phases. LA craters are marked by circles of 85 μm diameter. Representative grain boundaries are marked by black polygons.

260 4.3 Early phases and purity of dolomite

The fact that dolomite recrystallization may preserve former remnants of calcite is an important aspect to consider in dolomite geochronology. X-ray diffraction (XRD) analyses on rock powders can help resolving this issue and were applied on the studied samples. Two samples were identified as pure dolomite (MU-1 and MU-2), three samples contain minor calcite component (MAM-3, MAM-7 and KM-1) and one sample encompass minor quartz component along with the dolomite
 265 (Tm-MU-2; Fig. 7A). As a complimentary, EBSD maps combined with EDS analyses can further distinguish between dolomite and high-Mg calcite. For example, EBSD phase mapping identified $\sim 45\%$ dolomite, $\sim 48\%$ calcite and $\sim 7\%$ zero solution on sample KM-1. In samples MU-1 and MU-2 dolomite is much more abundant, with average of 67% dolomite, 25% calcite and 8% zero solution (Fig. 7B and C). Although calcite phase is relatively abundant in these samples, EDS has identified more than 2:3 Mg/Ca ratio, indicating it is a high-Mg calcite. This support previous interpretations of replacement of calcite by
 270 dolomite. The difference between XRD and EBSD analyses imply that pseudosymmetry of high-Mg calcite and dolomite can be unambiguously detected by *in-situ* EBSD phase mapping and not by XRD powder analyses. While the labor-intensive EBSD analysis is more sensitive in detecting calcite replacement than XRD, both methods are recommended for detecting impurities. In this study, less successful samples for dating (e.g. MAM-3 and MAM-7) are with higher calcite percentage relative to successfully dated samples (e.g. MU-1 and MU-2; Fig. 7C–D).



275

Figure 7: (A) XRD results of the studied samples: all samples are composed entirely of dolomite (peaks above black vertical lines), while some samples show minor calcite contribution (gray vertical lines). Sample Tm-MU-2 shows additional minor peaks of quartz. EBSD phase maps of samples KM-1 (B), MU-1 (C) and MU-2 (D). Dolomite is marked in purple, calcite in orange and zero solutions and grain boundaries are marked in black.

280

The WDS elemental maps of Fe, Mg and Ca were performed on sample KM-1 and are presented aside BSE image of the same location. The zonings in dolomite grains seen in the BSE are visible in the Fe map (Fig. 8B). Under the resolution of the scan (<0.01 wt.%), Mg and Ca maps do not show chemical zoning, but Ca-rich and Mg-depleted zones can be seen within grain boundaries. These clusters are probably remnants of primary calcite that was later replaced by dolomite (Fig. 8C, D). The WDS mapping could be therefore used for detecting zoning and remnant calcite impurities in the dolomite sample, which in

285

case of late dolomitization event(s), might shift the determined age towards the stratigraphic age of the sample. It is therefore highly recommended to use WDS elemental mapping for samples with sparry grains.




Figure 8: BSE image of LA crater on sample KM-1 (A) compared with WDS elemental maps of the same location (B-D). Zoning in dolomite rhomb is highlighted by Fe elemental map and absent on Mg and Ca. Mg-depleted and Ca-enriched clusters can be seen within the Fe rims of the dolomite crystals.

290

4.4 Average down-hole fractionation of RMs and selected unknowns


Results from samples MAM-3 and MAM-7 may be the most enigmatic of the sample set, as their ages are considerably older than expected, whereas other samples in this suite yield reasonably acceptable ages. One explanation might be that these samples had a different laser-induced elemental fraction (LIEF) than that of the rest of the sample suite and the calcite reference materials. Although similar in chemistry, these samples have a different texture from other samples, as they are micritic, rather than crystalline. Figure 9 shows stacked integration plots of the down-hole raw ^{207}Pb -corrected $^{206}\text{Pb}/^{238}\text{U}$ ratio of unknowns and RMs from each of two sessions in which a sample of either expected age (MU-2; session 1) or unexpected age (MAM-7; session 2) was analyzed. In both sessions, WC-1 (primary calcite RM), Duff Brown Tank (secondary calcite RM), NIST614 glass, and a zircon RM, Mud Tank (Black and Gulson, 1978), yielded consistent down-hole patterns, with zircon being the steepest, NIST614 with a minor negative slope and the calcite RMs in between. The down-hole pattern in MU-2 (run 1) was very similar to that of the primary calcite RM (WC-1) and it is therefore not surprising that it yielded the expected age. MAM-7 (run 2), however, yielded a negative down-hole fractionation pattern, beyond that of any of the standards. Using NIST614 as a primary standard for calcite yields an age that is too old for calcite reference materials, and long-term correction factors typically range between 10–20% for $^{206}\text{Pb}/^{238}\text{U}$. This is expected for the calcite vs. NIST glass fractionation patterns; the higher $^{206}\text{Pb}/^{238}\text{U}$ ratios of the calcite RMs down-hole would yield older ages relative to NIST. Interestingly, however, MAM-7 is older than expected, even though its $^{206}\text{Pb}/^{238}\text{U}$ ratio becomes smaller down-hole. This may indicate that the difference in $^{206}\text{Pb}/^{238}\text{U}$ ratios between measured and expected are caused by plasma-ionization differences between particles of MAM-7

295

300

305

and those of the reference materials and crystalline dolomite. A similar offset is seen in the zircon data; the steeper down-hole fractionation of Mud Tank zircon would expect an age that is older than the reference value. Instead, the recovered age was typically ca. 20% younger than its accepted value. This further indicates the importance of analyzing samples of similar chemical and textural makeup when standardizing unknowns, and that drill rate is only one component of age offset.



315 **Figure 9. Average down-hole fractionation of RMs and select unknowns. Raw ^{207}Pb -corrected values (corrected for baseline) are normalized to the average value and a linear fit shows different fractionation trends between glass, zircon, calcite and dolomite. Lower panel showing the difference in average down-hole fractionation between unknown samples and reference materials in two different analytical runs.**

4.5 Reevaluation of U–Pb results and interpretation

In sample EFN-1 fragments and cement arrange along two different isochrons, forming a wedge with mixed ages between isochrons (Fig. 10). The $^{207}\text{Pb}/^{206}\text{Pb}$ interception occurs to the right of the concordia curve, resulting in higher common-Pb

320 values for the ~~older isochron~~. The stratigraphic age of this faulted unit is considered Cenomanian and cropped out in other regions as limestones rather than dolomite. If dolomitization occurred after brecciation and cementation during a faulting event, a single age for both fragments and the cement is expected. However, fragments and cement yielded two distinct linear trends, indicating that dolomitization of the host rock occurred before brecciation and dolomitization of the cement occurred during or after the faulting event at 6.5 ± 1 Ma (MSWD = 1.5; n = 32). Along the fragments two isochrons of acceptable ages can be
325 identified, at 74 ± 3 and 58 ± 3 . The different ages within the fragments may represent two separated diagenesis and dolomitization events of the rock before faulting, whereas cementation and epigenetic dolomitization of the cement occurred much later, at ~ 6 Ma.

The age of sample MU-2 (93 ± 7) is correspond to the expected stratigraphic age for this unit and probably represent early diagenesis. In sample MU-1, on the other hand, a wedge pattern similar to the fragments in sample EFN-1 can be identified.
330 Out of 80 spot analyses, the older 13 dates form a reasonable isochron with age of 91 ± 6 Ma, with MSWD of 1.8. This ~~age~~ falls within the expected stratigraphic age and probably represents ~~early diagenetic age for that sample~~. The youngest 38 spot analyses yield an age of 53 ± 2 Ma, with MSWD of 2. The older isochron corresponds to the expected stratigraphic age of this sample, while the younger isochron is ~ 30 Ma younger and may reflect either the time of closure during late-stage dolomitization, or mixed age between stratigraphic age and a much younger dolomitization event (Fig. 10).

335 Despite its low-resolution isochron, a similar wedge pattern to MU-1 can be seen in sample KM-1, with an older age of 101 ± 11 Ma (MSWD = 0.46; n = 15) and younger age of 56 ± 3 Ma (MSWD = 0.94; n=50). This repeating pattern may represent actual dolomitization event at ~ 55 Ma in these localities. An Early Eocene dolomitization event is, however, not familiar in the local geological record. Hence, the age of 55 Ma may reflect mixed ages of stratigraphic age (early diagenesis) and some younger event(s), similar to sample EFN-1, whereas a young event correspond to the age of 6.5 Ma and association with
340 faulting along the Dead Sea Fault. The use of CL imaging can help to establish how homogeneous the samples are in terms of precipitation conditions. Micritic material are very hard to study by simple microscopy and slight differences in luminescence may suggest superimposed precipitation events. In such cases, early events that left very small remnant material, but with high U content, and a later dominant event with low U-content, can easily produce mixed age that is shifted towards and old ages. In such cases, it might be useful to implement the methodology described in Drost et al. (2018), in which 2-D elemental and
345 isotopic ratio maps are used for targeting subdomains in carbonate samples with complex geological histories, such as diagenetic overprinting.




Figure 10: Age interpretation of U–Pb results for samples EFN-1, MU-1 and KM-1.

Conclusions

- 350
- Accurate U–Pb dating of dolomite by laser ablation inductively coupled plasma mass spectrometry (LA-ICP-MS) contribute to better understanding of dolomitization process.
 - CL and BSE images highlight complexities in the chemical zoning of dolomite at sub-millimetre scale, including distinct core and rim, semi-homogenized grains, or mixture of different grains. Pre-analysis screening by these methods are recommended.
- 355
- Labor-intensive EBSD analysis is more sensitive in detecting calcite replacement than XRD, but both methods are recommended for detecting impurities.
 - A comparison of down-hole fractionation between RMs and unknowns, even those of similar chemical makeup, can be a valuable tool in estimating true uncertainty and inaccuracy of unknowns.
 - Textural characteristics such as micritic vs. well-crystalized grains have minor effect on ablation efficiency and can
- 360
- have only minor effect on the resulted ages.
 - Differences between obtained and stratigraphic ages, in particularly within micritic material, suggest for superimposed dolomitization events at the sub-millimetre scale. A detailed study by CL, EBSD, SEM or 2-D elemental and isotopic ratio maps are recommended in addition to U–Pb analysis.

Acknowledgments

365 We thank Dr. Omri Dvir from the Hebrew University, Israel for the help with WDS maps and to Dr. Navot Morag from the Geological Survey of Israel for the help with XRD analyses and interpretation. We thank Dr. Einat Nativ-Roth and Mrs. Roxana Golan from the Ilse Katz Institute for Nanoscale Sciences and Technology, Ben-Gurion University of the Negev for

their help with SEM imaging and EBSD mapping. This study was supported by the Israeli Government (under the Geological Survey of Israel, Dead Sea, project 40709).

370 **References**

- Ahm, A. S. C., Bjerrum, C. J., Blättler, C. L., Swart, P. K., & Higgins, J. A.: Quantifying early marine diagenesis in shallow-water carbonate sediments, *Geochimica et Cosmochimica Acta*, 236, 140-159, 2018.
- Beyth, M., Longstaffe, F. J., Ayalon, A., & Matthews, A.: Epigenetic alteration of the Precambrian igneous complex at Mount Timna, southern Israel: Oxygen-isotope studies, *Israel Journal of Earth Sciences*, 46, 1-11, 1997.
- 375 Beyth, M., & Heimann, A.: The youngest igneous event in the crystalline basement of the Arabian-Nubian Shield, Timna Igneous Complex, *Isr. J. Earth Sci*, 48, 113-120, 1999.
- Beyth, M., Segev, A., Bartov, Y.: Geological Map of Israel. Sheet 25-IV Be'er Ora, Scale 1:50,000, Israel Geological Survey, Jerusalem, 1999.
- Black, L. P. and Gulson BL.: The age of the mud tank carbonatite, strangways range, northern territory. *BMR Journal of*
- 380 *Australian Geology & Geophysics*, 3, 227-232, 1978.
- Burisch, M., Walter, B.F., Gerdes, A., Lanz, M. and Markl, G.: Late-stage anhydrite-gypsum-siderite-dolomite-calcite assemblages record the transition from a deep to a shallow hydrothermal system in the Schwarzwald mining district, SW Germany. *Geochimica et Cosmochimica Acta*, 223, 259-278, 2018.
- Drost, K., Chew, D., Petrus, J.A., Scholze, F., Woodhead, J.D., Schneider, J.W. and Harper, D.A.: An Image Mapping
- 385 Approach to U-Pb LA-ICP-MS Carbonate Dating and Applications to Direct Dating of Carbonate Sedimentation. *Geochemistry, geophysics, geosystems*, 19, 4631-4648, 2018.
- Eliyahu A., Katzir Y., Ginat H.: A new metallogenic model for sediment-hosted stratiform copper mineralization at the Timna-Faynan southern basin margin, northern Arabian Nubian Shield, in 6th Kaplan Workshop: Partitioning of trace elements and stable isotopes from the core to the oceans: thermodynamics and applications, Kibbutz Eilat, Israel, 2017.
- 390 Frisia, S., Borsato, A., & Hellstrom, J.: High spatial resolution investigation of nucleation, growth and early diagenesis in speleothems as exemplar for sedimentary carbonates, *Earth-Science Reviews*, 178, 68-91, 2018.
- Godeau, N., Deschamps, P., Guihou, A., Leonide, P., Tendil, A., Gerdes, A., Hamelin, B. and Girard, J.P.: U-Pb dating of calcite cement and diagenetic history in microporous carbonate reservoirs: Case of the Urgonian Limestone, France, *Geology*, 46, 247-250, 2018.
- 395 Guillong, M., Wotzlaw, J.F., Looser, N. and Laurent, O.: Evaluating the reliability of U-Pb laser ablation inductively coupled plasma mass spectrometry (LA-ICP-MS) carbonate geochronology: matrix issues and a potential calcite validation reference material. *Geochronology*, 2, 155-167, 2020.
- Gregg, J. M., & Sibley, D. F.: Epigenetic dolomitization and the origin of xenotopic dolomite texture, *Journal of Sedimentary Research*, 54, 908-931, 1984.

- 400 Hill, C. A., Polyak, V. J., Asmerom, Y., & P. Provencio, P.: Constraints on a Late Cretaceous uplift, denudation, and incision
of the Grand Canyon region, southwestern Colorado Plateau, USA, from U-Pb dating of lacustrine
limestone, *Tectonics*, 35, 896-906, 2016.
- Hodson, K.R., Crider, J.G. and Huntington, K.W.: Temperature and composition of carbonate cements record early structural
control on cementation in a nascent deformation band fault zone: Moab Fault, Utah, USA, *Tectonophysics*, 690, 240-252,
405 2016.
- Hoff, J. A., Jameson, J., & Hanson, G. N.: Application of Pb isotopes to the absolute timing of regional exposure events in
carbonate rocks; an example from U-rich dolostones from the Wahoo Formation (Pennsylvanian), Prudhoe Bay,
Alaska. *Journal of Sedimentary Research*, 65, 225-233, 1995.
- Hu, Y., Cai, C., Liu, D., Pederson, C.L., Jiang, L., Shen, A. and Immenhauser, A.: Formation, diagenesis and
410 palaeoenvironmental significance of upper Ediacaran fibrous dolomite cements. *Sedimentology*, 67, 1161-1187, 2020.
- Kohn, B, Weissbrod, T, Chung, L, Farley, K, Bodorkos, S.: Low-temperature thermochronology of francolite: Insights into
timing of Dead Sea Transform motion, *Terra Nova*, 31: 205– 219, 2019.
- Li, Q., Parrish, R. R., Horstwood, M. S. A., & McArthur, J. M.: U–Pb dating of cements in Mesozoic ammonites, *Chemical
Geology*, 376, 76-83, 2014.
- 415 Ludwig, K. R.: Isoplot/Ex, v. 3.75, *Berkeley Geochronology Center Special Publication*, 5, 75, 2012.
- Machel, H. G.: Concepts and models of dolomitization: a critical reappraisal, *Geological Society, London, Special
Publications*, 235, 7-63, 2004.
- Mueller, M., Igbokwe, O.A., Walter, B., Pederson, C.L., Riechelmann, S., Richter, D.K., Albert, R., Gerdes, A., Buhl, D.,
Neuser, R.D. and Bertotti, G.: Testing the preservation potential of early diagenetic dolomites as geochemical archives.
420 *Sedimentology*, 67, 849-881, 2020.
- Nuriel, P., Weinberger, R., Kylander-Clark, A.R.C., Hacker, B.R. and Craddock, J.P.: The onset of the Dead Sea transform
based on calcite age-strain
analyses, *Geology*, 45, 587-590, 2017.
- Nuriel, P., Wotzlaw, J., Ovtcharova, M., Vaks, A., Stremtan, C., Šala, M., Roberts, N., and Kylander-Clark, A.R.C.: The use
425 of ASH-15 flowstone as matrix-matched standard for laser-ablation geochronology of calcite, *Geochronology*, 2021.
- Olanipekun, B. J., & Azmy, K.: *In situ* characterization of dolomite crystals: Evaluation of dolomitization process and its effect
on zoning, *Sedimentology*, 64, 1708-1730, 2017.
- Ovchinnikova, G.V., Kuznetsov, A.B., Melezhhik, V.A., Gorokhov, I.M., Vasil'eva, I.M. and Gorokhovskii, B.M.: Pb–Pb age
of Jatulian carbonate rocks: the Tulomozero Formation of southeast Karelia. *Stratigraphy and Geological Correlation*, 15,
430 359-372, 2007.
- Parnes, A.: Late Lower Cambrian trilobites from the Timna area and Har Amram (southern Negev, Israel), *Israel Journal of
Earth Sciences*, 20, 179-205, 1971.

- Parrish, R.R., Parrish, C.M. and Lasalle, S.: Vein calcite dating reveals Pyrenean orogen as cause of Paleogene deformation in southern England, *Journal of the Geological Society*, 175, 425-442, 2018.
- 435 Paton, C., Woodhead, J. D., Hellstrom, J. C., Hergt, J. M., Greig, A., and Maas, R.: Improved laser ablation U-Pb zircon geochronology through robust downhole fractionation correction, *Geochemistry Geophysics Geosystems*, 11, 2010.
- Polyak, V.J., Provencio, P.P. and Asmerom, Y.: U–Pb dating of speleogenetic dolomite: A new sulfuric acid speleogenesis chronometer. *International Journal of Speleology*, 45, 2016.
- Putnis, A. and Putnis, C. V.: The mechanism of re- equilibration of solids in the presence of a fluid phase, *Journal of Solid-*
 440 *State Chem.*, 180, 1783–1786, 2007.
- Ring, U. and Gerdes, A.: Kinematics of the Alpenrhein-Bodensee graben system in the Central Alps: Oligocene/Miocene transtension due to formation of the Western Alps arc, *Tectonics*, 35, 1367-1391, 2016.
- Roberts, N.M.W. and Walker, R.J.: U-Pb geochronology of calcite-mineralized faults: Absolute timing of rift-related fault events on the northeast Atlantic margin, *Geology*, 44, 531-534, 2016.
- 445 Roberts, N. M. W., Rasbury, E. T., Parrish, R. R., Smith, C. J., Horstwood, M. S. A., and Condon, D. J.: A calcite reference material for LA- ICP-MS U–Pb geochronology, *Geochemistry Geophysics Geosystems*, 18, 2807-2814, 2017.
- Roberts, N.M., Drost, K., Horstwood, M.S., Condon, D.J., Chew, D., Drake, H., Milodowski, A.E., McLean, N.M., Smye, A.J., Walker, R.J. and Haslam, R.: Laser ablation inductively coupled plasma mass spectrometry (LA-ICP-MS) U-Pb carbonate geochronology: strategies, progress, and limitations, *Geochronology discuss*, 2, 33-61, 2020.
- 450 Salih, N., Mansurbeg, H., Kolo, K., Gerdes, A. and Pr eat, A.: *In situ* U–Pb dating of hydrothermal diagenesis in tectonically controlled fracturing in the Upper Cretaceous Bekhme Formation, Kurdistan Region-Iraq. *International Geology Review*, 1-19, 2019.
- Sass, E.: Microphotometric determination of preferred orientation in undeformed dolomites, *Science*, 165, 802-803, 1969.
- Sass, E., and Bein, A.: Platform carbonates and reefs in the Judean Hills, Carmel and Galilee, *in* Proceedings Tenth
 455 International Congress on Sedimentology. International Association of sedimentologists, Jerusalem, Field guidebook, 239-274, 1978.
- Sass, E., Dekel, A., Sneh, A.: Geological Map of Israel. Sheet 5-II Umm El Fahm, Scale 1:50,000. Israel Geological Survey, Jerusalem, 2013.
- Schmitz, M.D. and Schoene, B.: Derivation of isotope ratios, errors, and error correlations for U–Pb geochronology using
 460 ^{205}Pb - ^{235}U -(^{233}U)-spiked isotope dilution thermal ionization mass spectrometric data. *Geochemistry, Geophysics, Geosystems*, 2007.
- Segev, A., Sass, E., Ron, H., Lang, B., Kolodny, Y., and McWilliams, M.: Stratigraphic, geochronologic, and paleomagnetic constraints on Late Cretaceous volcanism in northern Israel, *Israel Journal of Earth Sciences*, 51, 2002.
- Segev, A., Sass, E.: Geological Map of Israel. Sheet 3- III Atlit, Scale 1:50,000. Israel Geological Survey, Jerusalem, 2009.
- 465 Segev, A.: $^{40}\text{Ar}/^{39}\text{Ar}$ and K–Ar geochronology of Berriasian–Hauterivian and Cenomanian tectonomagmatic events in northern Israel: implications for regional stratigraphy, *Cretaceous Research*, 30, 810-828, 2009.

- Segev, A.: Major unconformities in the stratigraphic succession in the Timna-Eilat region and their relation to copper and manganese mineralization cycles, *Isr. Geol. Soc. Field Trips Guide. Elat*, 154-188, 2016.
- 470 Sneh, A., Avni Y.: Geological Map of Israel. Sheet 11- II Jerusalem, Scale 1:50,000. Israel Geological Survey, Jerusalem, 2016.
- Stacey, J.T. and Kramers, J.: Approximation of terrestrial lead isotope evolution by a two-stage model. *Earth and planetary science letters*, 26, 207-221, 1975.
- Tera, F., and Wasserburg, G. J.: U-Th-Pb systematics in three Apollo 14 basalts and the problem of initial Pb in lunar rocks, *Earth and Planetary Science Letters*, 14, 281-304, 1972.
- 475 Vermeesch, P.: IsoplotR: A free and open toolbox for geochronology, *Geoscience Frontiers*, 9, 1479-1493, 2018.
- Warren, J.: Dolomite: occurrence, evolution and economically important associations, *Earth-Science Reviews*, 52, 1-81, 2000.
- Winter, B.L. and Johnson, C.M.: U-Pb dating of a carbonate subaerial exposure event. *Earth and Planetary Science Letters*, 131(3-4), pp.177-187, 1995.