

Robust Isochron Calculation

Roger Powell¹, Eleanor CR Green¹, Estephany Marillo Sialer¹, and Jon Woodhead¹

¹School Earth Sciences, The University of Melbourne, Vic 3010, Australia

Correspondence: Roger Powell (powell@unimelb.edu.au)

Abstract.

The standard classical statistics approach to isochron calculation assumes that the distribution of uncertainties on the data arising from isotopic analysis is strictly Gaussian. This effectively excludes from consideration datasets that have more scatter, even though many appear to have age significance. A new approach to isochron calculations is developed in order to circumvent this problem requiring only that the central part of the uncertainty distribution of the data defines a “spine” in the trend of the data. This central spine can be Gaussian but this is not a requirement. This approach significantly increases the range of datasets from which age information can be extracted but also provides seamless integration with well-behaved datasets, and thus all legacy age determinations. The approach is built on the robust statistics of Huber (1981), but using the data uncertainties for the scale of data scatter around the spine, rather than a scale derived from the scatter itself, ignoring the data uncertainties. This robust data-fitting reliably determines the position of the spine when applied to data with outliers, but converges on the classical statistics approach for datasets without outliers. The spine width is determined by a robust measure, the normalised median absolute deviation of the distances of the data points to the centre of the spine, divided by the uncertainties on the distances. A test is provided to ascertain that there is a spine in the data, requiring that the spine width is consistent with the uncertainties expected for Gaussian-distributed data. An iteratively-reweighted least squares algorithm is presented to calculate the position of the robust line and its uncertainty, accompanied by an implementation in Python.

1 Introduction

The ability to fit a straight line through a body of isotope ratio data in order to form an isochron is the cornerstone of many geochronological methods. In detail, however, this is a non-trivial task, since uncertainties are usually associated with all variables, and these are often correlated, precluding simple “least squares” line-fitting techniques. Most of the research in this area was conducted in the late 1960’s and early 1970’s, being dominated by a classical statistics approach in which data uncertainties, derived from the analytical methods, are taken to be strictly Gaussian-distributed (e.g. York, 1969; York et al., 2004, and references therein). This approach, referred to here as YORK, became entrenched in the geochemical community, particularly in the last two decades as the essential component of the very widely-used software, ISOPLOT, e.g. Ludwig (2012).

In this contribution we examine some of the problems inherent in these techniques and suggest an alternative approach. Our primary focus here will be on general-purpose isochron calculations that determine the age of an “event” that established the isotopic compositions of samples in a dataset. This involves what are called model 1 and 2 calculations in ISOPLOT - as

described below. Approaches that try to extract detail within events, including ISOPLOT model 3 calculations, are not considered (but see e.g. Vermeesch, 2018).

1.1 On ISOPLOT

30 In order to show that there are significant problems in using ISOPLOT for general-purpose isochron calculations, and then to see how they can be addressed, it is first necessary to outline the ISOPLOT protocol, some details of which may not be apparent to the end user. Central to the ISOPLOT workflow, the main tool for considering data scatter is m_{swd} , the mean standard weighted deviates (also called the reduced chi-squared statistic), see eq. 1. For strictly Gaussian distributed uncertainties $(n-2)m_{\text{swd}}$ is distributed as chi-squared (χ_{n-2}^2), meaning that if uncertainties are correctly assigned, a strong statistical statement can be made
35 about whether the scatter of a particular dataset is solely consistent with the associated data uncertainties (i.e. with no geological scatter), for example in the form of a 95% confidence interval on m_{swd} , Wendt and Carl (1991). Such a confidence interval is not fixed, but depends on the number of datapoints under consideration, so for example for $n = 10$, $m_{\text{swd}} < 1.94$ (meaning that m_{swd} extends from less than 1 through to 1.94), while for $n = 50$, $m_{\text{swd}} < 1.36$. A dataset with m_{swd} in the chosen range for the number of datapoints has data scatter that is consistent with the data uncertainties. This situation is commonly referred to as
40 m_{swd} “passes”; otherwise m_{swd} “fails”, in relation to χ_{n-2}^2 . The situation where m_{swd} passes provides a “pure” interpretation of YORK, and, in ISOPLOT is referred to as a model 1 isochron fit. This is depicted as the horizontal line in Fig. 1, indicating that in this range of m_{swd} , corresponding to a confidence interval, the calculated uncertainty on an isochron age does not vary with m_{swd} . Such a figure is drawn by taking an actual dataset and progressively modifying it to show what happens as m_{swd} varies, as described in Appendix A.

45 What if m_{swd} is greater than the upper limit of the chosen confidence interval? Then the data are considered to have *excess* scatter, in addition to that accounted for by the data uncertainties (assuming that they are strictly Gaussian)¹. At this point, ISOPLOT, asks the user whether an alternative—model 2—calculation should be undertaken. This decision point is indicated at A in Fig. 1. If the user declines, ISOPLOT gives results that are referred to here as model 1x, as shown in Fig. 1. The model 1 age uncertainty is multiplied by $\sqrt{m_{\text{swd}}}$, to reflect the data scatter being more than expected from the data uncertainties alone.
50 With further scatter, the switch is made to model 2, at B in Fig. 1. Alternatively, if the user accepts the switch to model 2, then model 1x is not used, indicated by the vertical line at A extending up to the model 2 line in Fig. 1. The model 2 calculation in ISOPLOT is unrelated to YORK. The data uncertainties are discarded and the slope of the line through the linear trend is calculated as the geometric mean of the lines calculated by unweighted least squares of y on x and of x on y (see Appendix A).

55 In summary, then, in ISOPLOT the calculation of ages and their uncertainties involves a number of decision points based around the concept of m_{swd} that impart significant (and in our view unwelcome) step-changes in the way that the data are handled, and algorithms applied. To assist in further discussion of these matters we depart from the language of ISOPLOT at this point, reintroducing the term *errorchron*, counterposed to *isochron*, following Brooks et al. (1972). The idea is that

¹excess scatter occurs when the data distribution has higher variability in the tails than is indicated by the variability of the central part of the distribution, for example if the distribution is Gaussian in the centre but having “fat” tails




Figure 1. Age uncertainty ($\text{age} \pm$) plotted against $\sqrt{\text{mswd}}$ under the ISOPLOT protocol for a progressively modified dataset (see text, and Appendix A). Under the condition of a model 1 fit, the age uncertainty is constant with increasing data scatter (reflected in increasing mswd), until there is a step change in the data treatment at A when the age uncertainty is multiplied by $\sqrt{\text{mswd}}$. Then at B there is another step change in age uncertainty calculation with increasing data scatter forming ISOPLOT model 2 (see text)

isochrons have a higher chance of having age significance, while errorchrons have a lower chance. In particular, it seems to be unhelpful for the results of model 2 calculations to be called isochrons as is the case in ISOPLOT, given that there is excess scatter in the data.

1.2 Replacing ISOPLOT

Given that ISOPLOT's implementation of model 1+2 line fits is the gold standard of isochron calculations presently, where are the problems, and then what can be done to address them? A shortcoming in YORK stems from the assumption that data uncertainties are *strictly* Gaussian-distributed. In real-world application this appears to be too restrictive, with datasets that are likely to have age significance being labelled as errorchrons because mswd is too large. While using YORK guided by mswd is optimal statistically if data uncertainties are strictly Gaussian, this logic fails once uncertainties are even slightly non-Gaussian (originally Tukey, 1960). In such circumstances, both mswd and least squares methods themselves, like YORK, become unreliable (e.g. Hampel et al., 1986; Huber, 1981).

Rather than being truly Gaussian, data uncertainties may well be Gaussian-distributed in their centres, but slightly fat-tailed distant from the centres. An isotopic dataset looks intuitively acceptable if the data has a central linear "spine", in which scatter is commensurate with stated analytical uncertainty, but this spine is flanked by data of somewhat larger scatter (i.e. excess scatter, from the "fat tail"). This excess scatter may originate in the isotopic analysis or as a result of geological disturbance.

Age-significance in such data manifests primarily via the position of the spine. In the following, we focus on this spine in the
75 data.

Adopting this spine approach, a successful calculation method for a dataset that may not have strictly Gaussian-distributed uncertainties must, firstly, ascertain whether or not such a spine exists in the data—and hence whether calculations yield an isochron or an errorchron. Secondly, in the case of an isochron calculation, the successful method must reliably locate the spine without being perturbed by vagaries in the more scattered data. Classical statistical methods can do neither of these
80 things, tending to be excessively influenced by the data at the extremes of the scatter. However, the field of robust statistics offers calculation methods that can. When a dataset has no excess scatter, reflected in `mswd` lying within an appropriate χ^2 -constrained confidence interval, such methods can be devised to retrieve identical (or nearly identical) results to classical statistical methods, but, in addition, provide reliable age and age-uncertainty estimates in the presence of excess scatter around a spine. This continuity of operation with increasing `mswd` contrasts with previous approaches and means that the steps in the
85 ISOPLOT line in Fig. 1, which are certainly undesirable, are circumvented. Moreover the involvement of potentially unreliable least squares-based methods, like ISOPLOT model 2, are avoided when the data show excess scatter.

2 An Algorithm for Isochron Calculations

An algorithm is sought that finds a robust straight line through a 2-dimensional linear data trend, while converging with the classical statistical approach of YORK for datasets with consistent scatter (i.e. `mswd` passes). This section describes the nature
90 of the problem and the theoretical basis for the robust statistical approach that will be adopted. The algorithm adopted

1. determines a preliminary fit of the data, not dependent on vagaries of the data scatter
2. determines the spine width in relation to this preliminary fit
 - if the spine width is in an acceptable range: isochron
 - if the spine width is not in an acceptable range: errorochron
- 95 3. determines a robust fit of the data, starting from the preliminary fit, this fit converging with YORK for “good” data

This algorithm is fleshed out below, and then evaluated via simulated datasets and applied to a natural dataset. The central calculation in the algorithm is detailed in Appendix B, and a `python` implementation is provided in Appendix C.

2.1 Uncertainty distributions and data fitting

Geochronological datasets are collected on the presumption that the isotopic compositions were established via an “event”
100 the age of which is to be estimated. Given the focus here on data with linear trends, even if the effect of the event is recorded perfectly by the samples analysed—the isotopic compositions lying on a line—the actual data are measured with finite precision and so the data inevitably scatter about the trend. An uncertainty probability distribution can be used to describe the form of the data scatter.

Classical statistical methods assume that the underlying uncertainty distribution of a dataset is known, typically taken to be Gaussian. Under the Gaussian assumption, if the analytical uncertainty on the measurements have been appropriately inferred, m_{swd} , the classical statistics parameter used in YORK to validate an isochron, tests that the scatter of datapoints is consistent with the inferred uncertainties. But, in general, there is no reason to suppose that a given analytical technique generates a truly Gaussian uncertainty distribution. Even small amounts of geological disturbance destroy the optimality of YORK. If the uncertainty distribution is not strictly Gaussian then classical methods of data fitting become sub-optimal or worse.

While there are many possible non-Gaussian uncertainty distributions, this paper is concerned with a situation commonly occurring in datasets, in which the datapoints form a linear spine with Gaussian-like scatter, but additional scatter is seen in the tails of the distribution. Such a dataset still encodes meaningful age information in its spine, yet it will typically fail an m_{swd} test owing to its departure from a Gaussian distribution. In this work, datasets of this nature are modelled using a contaminated Gaussian uncertainty distribution (Gaussian mixture) (e.g. Tukey, 1960). Such distributions are written $c\%dN$, meaning that with a probability $(100 - c)\%$ the distribution involves a standard deviation, σ , but with a probability $c\%$ the distribution has a standard deviation, $d\sigma$, both with a mean of zero (see Powell et al., 2002; Maronna et al., 2019, Sect. 2.1). An example is 25%3N, with $c = 25$ and $d = 3$, so that with 25% probability the uncertainty is drawn from $N(0, 3\sigma)$, and 75% probability drawn from $N(0, \sigma)$, with the $N(0, s)$ notation indicating a Gaussian distribution with a mean of zero and a standard deviation of s . Such distributions provide excess scatter suitable for developing and evaluating a robust line-fitting calculation. It does not matter if excess scatter in real data is drawn from a different distribution. Note that with the sample sizes provided by most modern geochronological techniques, it is not necessarily possible to test for Gaussian behaviour, or such departures from Gaussian behaviour, although they may be evident on quantile-quantile plots (see below).

2.2 Isochrons and errorchrons

In YORK, assuming that the data uncertainties are strictly Gaussian distributed, the probability distribution of m_{swd} provides bounds that can be used to distinguish isochrons from errorchrons (e.g. Wendt and Carl, 1991). These bounds come from a 95% confidence interval on m_{swd} , as discussed in Appendix A. Datasets whose scatter give m_{swd} outside the bounds are deemed to be errorchrons, not isochrons. The focus in this paper is on m_{swd} that is too large, indicating excess scatter. M_{swd} is defined with the residuals, r_k , the distance in y of the datapoint, k , to the line, e_k , weighted by the uncertainty on this distance, σ_{e_k}

$$m_{swd} = \frac{1}{n-2} \sum_{k=1}^n r_k^2 \quad \text{with} \quad r_k = \frac{e_k}{\sigma_{e_k}} = \frac{a + bx_k - y_k}{b^2 \sigma_{x_k}^2 + \sigma_{y_k}^2 - 2b \sigma_{x_k} \sigma_{y_k} \rho_{x_k y_k}} \quad (1)$$

The line being fitted is $y = a + bx$; datapoint, k , is $\{x_k, y_k\}$; the analytical uncertainty on x_k , σ_{x_k} ; the analytical uncertainty on y_k , σ_{y_k} ; and the correlation between x_k and y_k , $\rho_{x_k y_k}$ (see derivation of eq. B4 in Appendix B). Note that the slope, b , appears in the denominator of r_k , as well as the numerator.

If, instead, data uncertainties are $c\%dN$, with unknown c and d , or some other contaminated Gaussian distribution then there is no equivalent of the m_{swd} argument to say which datasets should give isochrons rather than errorchrons. The approach advocated here is to use a measure that reflects whether the dataset has a linear spine of “good” data within it. The measure

suggested, s , coined the spine width, is robust, and is defined as

$$s = \text{nmad}(r) = 1.4826 \text{ median}(|r_k - \text{median}(r)|) \quad (2)$$

with a normalisation constant², 1.4826. Given that s is based on a median, its magnitude depends on that half of the data that have the smallest absolute values of centred r , in other words those that would define a spine. If the data were in fact Gaussian-distributed, it is expected that s should be in a range about 1 in the same way that mswd is, given that r already involves the analytical uncertainties. The larger is s , greater than 1, the less pronounced is the linear spine in the data (or the uncertainties have been underestimated). Whereas the 95% confidence interval (95%ci) on mswd for Gaussian-distributed uncertainties comes from a well-established probability distribution, with $(n-2)\text{mswd} \sim \chi_{n-2}^2$ (e.g. Wendt and Carl, 1991), the confidence interval on s needs to be found by simulation (see Appendix D), with the simulated datasets just involving Gaussian-distributed uncertainties. The intervals are given in this table, Table 1:

n	95%ci		$\sqrt{\text{mswd}}$	95%ci		s
	low	high	*	low	high	*
5	0.268	1.765	1.614	0.09	1.64	1.48
6	0.348	1.669	1.540	0.17	1.62	1.47
8	0.454	1.552	1.449	0.26	1.58	1.45
10	0.522	1.480	1.392	0.31	1.55	1.43
15	0.621	1.379	1.312	0.40	1.50	1.40
30	0.739	1.260	1.215	0.58	1.39	1.33
60	0.818	1.181	1.151	0.71	1.28	1.23

Whereas 1-sided confidence intervals are advocated in Appendix A—columns marked with an asterisk in the Table—2-sided confidence intervals are also given in the table. Using the asterixed column, for example for a dataset with 10 datapoints ($n = 10$), the dataset is deemed to yield an isochron if the observed s is less than 1.43. If s is larger, the dataset gives an errorchron. For isochrons, the age uncertainty is calculated as in Appendix B. For errorchrons, the age uncertainty is not calculated because the data scatter is not consistent with the analytical uncertainties at 95% confidence, s being larger than the bound.

2.3 A robust statistics approach to isochron calculation

We seek a statistical approach to isochron calculation that is *robust* (e.g. Huber, 1981; Hampel et al., 1986), meaning that it is not excessively affected by outliers in the data, while having desirable statistical properties, for example good efficiency (see below). In addition, we require the approach to converge to YORK for a “good” dataset, one with a near-Gaussian uncertainty distribution, allowing seamless compatibility with classical data interpretation. The overall approach adopted will be referred

²the normalisation constant makes $\text{nmad}(r)$ an unbiased estimator of the standard deviation when the data are Gaussian distributed as the sample size becomes large (e.g. Maronna et al., 2019, Sect. 2.4)

to as SPINE, involving the use of spine width for isochron-errorchron distinction, combined with robust line-fitting. The line-fitting is based on the approach of Huber (1981), as outlined in Maronna et al. (2019), Sect. 2.2.2. Whereas most robust
 160 line-fitting methods use the scatter of the data as a scale, data uncertainties having been discarded (e.g. Powell et al., 2002),
 here the data uncertainties are used. This is necessary in order to preserve continuity of the results with YORK, in which the
 data uncertainties are an integral part of the calculation.

In both the Huber approach in SPINE, and in YORK, a straight line is fitted to a dataset by minimising a function of the
 residuals, r_k . In the case of YORK this is just the mswd , (1). Since isochron data are generally bivariate with correlated
 165 analytical uncertainties in x and y , the analytical uncertainty in datapoint k can be represented as an ellipse as in Fig. 2. The
 absolute value of the residual for datapoint, k , r_k , is in fact the scaling factor on the size of the ellipse required to expand it or
 reduce it until it touches the best-fit line (Fig. 2).

The function that is minimised to find the best-fit line can be written $\sum \rho(r_k)$ for both YORK and SPINE. Whereas in YORK,
 $\rho(r_k) = r_k^2$ for all r_k , in SPINE $\rho(r_k) = r_k^2$ near the centre of the uncertainty distribution (as in YORK), but downweights
 170 datapoints for which the absolute value of the residual is greater than a cut-off value, h . Thus, in SPINE, and in Fig. 3, the
 Huber ρ is

$$\rho(r_k) = \begin{cases} 2hr_k - h^2 & r_k < -h \\ r_k^2 & \text{if } -h < r_k < h \\ 2hr_k - h^2 & r_k > h \end{cases} \quad (3)$$




Figure 2. For an example datapoint, $\{x_k, y_k\}$, the inner ellipse is calculated with the analytical uncertainties, V_k , at the 1σ level (in black). Given a line, $y = a + bx$ (in blue), the ellipse must be drawn at the $|r_k|\sigma$ level (in red) to touch the line, in this case $|r_k| = 5.73$. The data point is $x_k = 529.14$, $y_k = 0.5614$, and $\sigma_{x_k} = 1.870$, $\sigma_{y_k} = 0.00127$ and $\rho_{x_k y_k} = -0.967$. The line is $y = 0.8108 - 0.0004764x$.




Figure 3. Plots of $\rho(r)$ against r for YORK in red (r^2), and for Huber in SPINE (eq. 3) in blue. The two curves are coincident for $|r| < h$, with $h = 1.4$ the vertical green lines. See text.

In SPINE, for residuals that have an absolute value less than an adjustable constant, h , the contribution to the sum being minimised is the same as for YORK, but it is linear in the residual for larger absolute value. Note that as h becomes larger and larger, SPINE converges to YORK. The value to use for h is discussed in Maronna et al. (2019)³.

The iteration developed in Appendix B minimises $\sum_k \rho(r_k)$ with respect to the unknown, θ , a two-element column vector, $\{a, b\}^T$ in the line equation, $y = a + bx$. The iteration is applicable to SPINE and also YORK. As a starting point of the iteration, data are fit for θ with SIEGEL (Siegel, 1982), which is highly robust toward excess scatter. However SIEGEL is much less efficient than SPINE (see below), so SPINE is a better ultimate estimator. A full iteration is envisaged in Appendix B. Once θ is calculated, the measure of scatter used to distinguish an isochron from an errorchron can be calculated using Table 1. If an isochron is deemed to have been calculated, the uncertainty on θ , \mathbf{V}_θ , can be found, as outlined in Appendix B.

The SPINE algorithm can be summarised:

1. determine a preliminary fit of the data using SIEGEL
2. determine the spine width using `nmad`
 - if the spine width is in an acceptable range, from col. 6 in Table 1: isochron
 - if the spine width is not in the acceptable range: errorchron
3. determine a robust fit of the data, minimising $\sum \rho(r_k)$, with the Huber $\rho(r_k)$, (3), starting from the SIEGEL estimate of θ

³The value used here, $h = 1.4$, involves a particular tradeoff between robustness and efficiency (Maronna et al., 2019, Sect. 2.2.2 and 3.4)

2.4 Application of SPINE to simulated datasets

Assessing algorithms for data fitting is best done using simulated datasets, so that the true “age” represented by the data is known. In this case, datasets were generated by drawing data points from a range of uncertainty distributions, all centred on a linear trend reflecting an age of 4 Ma. Full details are provided in Appendix D. Two features of the datasets are varied: the number of datapoints in the dataset, and the uncertainty structure adopted, the latter via varying c and d in $c\%dN$. The algorithm is assessed in terms of its ability to retrieve the specified age of the linear trend on which the simulated datasets are built, and on the uncertainty in the age.

Given that the datasets investigated have fat-tailed contaminated-Gaussian uncertainty distributions, the focus is on the effect of excess scatter in the data, in other words, data scatter over and above what is expected for Gaussian data uncertainties. Nevertheless a small proportion of datasets do have small scatter, giving s which is below the lower bound for that number of datapoints.

The analysis below compares the results of YORK, applied only to those simulated datasets that lie within the m_{swd} bounds, with the results of SPINE applied to those datasets that lie within the spine width (s) bounds. The greatest majority of the former are included in the latter, e.g. $> 97\%$ for $n = 10$). Importantly, however, SPINE typically identifies reliable age information in many more datasets than YORK. In the following table, $m\%excl$ and $s\%excl$ are the percentage of simulated datasets excluded on the basis of the m_{swd} and s bounds, respectively:

n	N		5%3N		25%3N		10%10N	
	$m\%excl$	$s\%excl$	$m\%excl$	$s\%excl$	$m\%excl$	$s\%excl$	$m\%excl$	$s\%excl$
5	2.5	2.5	8.7	4.0	30.2	9.8	32.5	9.5
6	2.5	2.5	9.6	3.9	34.6	13.8	37.3	10.9
8	2.5	2.5	12.7	4.2	44.7	14.5	46.0	10.4
10	2.5	2.5	14.2	4.0	51.8	15.2	53.5	9.7
15	2.5	2.5	17.4	4.2	65.2	17.1	68.2	9.1

Note that, for example, for $n = 10$, datasets drawn from 5%3N, in fact have $100(10^{0.95}) = 59.9\%$ of the datasets having all uncertainties Gaussian, while 40.1% have at least one uncertainty drawn from 3 times Gaussian (3N). For 25%3N, 5.6% are Gaussian only, and for 10%10N, 34.9%. The leftmost columns are 2.5% by definition.

A 95% confidence interval on age can be found by ordering the list of ages calculated for the datasets, selecting the the lower limit at the 2.5% point in the list, and the upper limit at the 97.5% point. For datasets that lie within the s bound in SPINE or m_{swd} bound in YORK, the 95% confidence intervals on ages are essentially the same, even though a much larger proportion of datasets lie within bounds using SPINE than using YORK. A comparison of ages can also be made between YORK and SPINE for simulated datasets that lie *outside* the m_{swd} bounds. For $n = 10$ and Gaussian-distributed data, the 95% confidence interval is ± 0.021 Ma for both YORK and SPINE, but for 5%3N, 25%3N and 10%10N, the 95% confidence interval on the YORK ages are ± 0.035 , ± 0.040 , and ± 0.092 Ma, respectively, while the SPINE ages are ± 0.027 , ± 0.034 , and ± 0.034 Ma. The differences between the ages given by the two methods, normalised by the uncertainties on the SPINE ages, are ± 1.62 , ± 1.63 and ± 5.54 .

Comparison of the intervals shows that the m_{swd} bounds are relevant to the application of YORK, in that generally it does not work well outside the bounds, while indicating that SPINE is (much) more reliable for age determination for such datasets.

Considering age uncertainty, it might be expected that the uncertainties suffer from the excess scatter in the data in datasets that yield an errorchron with YORK but an isochron with SPINE. This appears not to be the case, but there is a small degradation
 220 in the age uncertainties retrieved caused by an unavoidable efficiency loss using SPINE. Efficiency at the Gaussian distribution is the ratio of the variance obtained by the optimal estimator (YORK), divided by the variance using the chosen robust estimator (in this case, SPINE). SPINE involves an efficiency loss at the Gaussian distribution, which is illustrated in Figure 4 via kernel density estimate (kde) plots of the age uncertainties calculated for simulated datasets with $n = 10$ and Gaussian-distributed uncertainties. KDE plots are probability distributions akin to smoothed histograms (Wand and Jones, 1995). The red curve is
 225 the kde for datasets that have all $|r_k| < h$, as would be given by YORK on datasets with Gaussian-distributed uncertainties. The blue curve is the kde for all datasets with at least one $|r_k| > h$. The overall kde, in black, is the kde of all of the datasets in the red and blue kde, in observed proportion, about 30% to 70%. The efficiency loss of SPINE is reflected in the displacement of the black curve to slightly higher age uncertainty than the red curve. The relationships shown in Figure 4 for $n = 10$ can be seen for other n in Figure 5. The pairs of red and black lines, correspond to, and have the same meaning as the red and black
 230 lines in Figure 4. As expected, the distribution of age uncertainties moves towards larger values as the sample size n decreases.




Figure 4. Kernel density estimates (kde) for age uncertainty calculated with SPINE on 10,000 simulated datasets with $n=10$ and Gaussian-distributed uncertainties. (a) Those datasets for which all $|r_k| < h$ (in red); (b) those datasets for which at least one $|r_k| > h$ (in blue), and (c) overall result combining a and b in observed proportion (in black).

Whereas SPINE involves an efficiency loss at the Gaussian distribution, it performs better—i.e. it is much more robust—than YORK for data from contaminated Gaussian distributions (e.g. Maronna et al., 2019, Table 2.2). The ability of SPINE to retrieve age uncertainty information for such data varies with the probability and scale of the contamination, as shown in Figure 6. Not unexpectedly the more seriously contaminated distributions (25%3N and 10%10N) involve a greater displacement of the kde




Figure 5. Kernel density estimates for age uncertainty calculated with SPINE on 10,000 simulated datasets with a range of n values and Gaussian-distributed uncertainties. In each case, the kde for those datasets for which all $|r_k| < h$ is in red and the overall kde is in black.

235 to higher age uncertainty than the more weakly contaminated 5%3N distribution. Although the displacement of the blue curves from the black curve is real, nevertheless the ability of SPINE to retrieve age uncertainties from datasets with contaminated distributions is good.




Figure 6. Kernel density estimates for age uncertainty calculated with SPINE on 10,000 simulated datasets with $n=10$ and several uncertainty structures. The kde for those datasets for which all $|r_k| < h$ is in red, the kde for datasets with Gaussian-distributed uncertainties is in black, and the kde for all of the datasets for 5%3N, 25%3N and 10%10N respectively are in blue.

2.5 Application of SPINE to a natural dataset

In order to show the real-world utility of SPINE, we consider data for a carbonate flowstone from the Riversleigh World
240 Heritage fossil site in Queensland, Australia (Sample 0708). Isotope dilution U-Pb data for the bulk sample were previously
published by Woodhead et al. (2016) providing a Model 2 isochron with an age of 13.72 ± 0.12 Ma and with a m_{swd} of 3.7.
The new data presented here were obtained by laser ablation ICPMS on the same sample using methods outlined and published
in Woodhead and Petrus (2019). Such datasets are typically larger with little error correlation (rounder error ellipses), but with
larger uncertainties than isotope dilution data. These new data define an errorchron under the YORK assumptions, with $m_{swd} =$
245 1.68, and a model 2 age of 13.68 ± 0.31 Ma. These data might therefore be rejected under the m_{swd} criterion despite exhibiting
a well-developed linear trend. With SPINE, $s = 1.24$, within the s range for an isochron, the age is 13.69 ± 0.26 Ma (\pm is
1.96 σ). The data for 0708 are plotted in Fig. 7, with 95% confidence ellipses on the datapoints. Further calculations with this
sample, comparing the results of our new algorithm with existing approaches are presented in Appendix A.




Figure 7. Laser ablation example 0708. See text.

With 0708, the SPINE and ISOPLOT ages are very similar, but m_{swd} suggests an errorchron, so there is excess scatter on this
250 basis. Quantile-quantile plots can be used to show the nature of the distribution of the residuals that contribute to such excess
scatter. Here, Fig. 8, the ordered residuals normalised to $n_{mad}(r)$ are plotted against the quantiles of the standard Gaussian
distribution, $N(0, 1)$ (see Fox, 2016, Sect 3.1.3). The normalisation means that if the residuals are Gaussian-distributed, they
should plot close to a line of unit slope through the origin. The figure also includes 95% pointwise confidence intervals (blue
dashed lines). The figure suggests that the residuals are in fact effectively Gaussian-distributed, lying between the blue-dashed
255 lines (and m_{swd} has been too sensitive in flagging that the residuals are not Gaussian-distributed).

Generally, geochronological datasets do not have obvious outliers—they might be “cleaned” of isolated datapoints before
an age calculation, or the dataset might even be discarded. But many of the simulated datasets from contaminated Gaussian




Figure 8. Quantile-quantile plot for sample 0708, using the SIEGEL fit of the data to calculate the residuals to plot. The dashed red line has unit slope through the origin; the blue dashed lines are the 95% pointwise confidence intervals. See text.

distributions, as used above, do contain outliers. In Appendix E an example of a simulated dataset with outliers (from 25%3N) is used to show how YORK and SPINE behave, and to show what a quantile-quantile plot looks like in those circumstances.

260 3 Discussion

This work was motivated by the belief that many isotopic datasets contain meaningful age information that cannot be identified using classical statistical methods and may therefore be discarded or discounted. In such cases, the age information is contained in a linear spine in the data, but the data also contains scatter that is inconsistent with a Gaussian uncertainty distribution, having fatter tails than Gaussian. A statistical test based on the spine width is devised, akin to using m_{swd} in classical methods, allowing an isochron-errorchron distinction to be made. Many datasets that give isochrons based on the spine width, give errorchrons under the assumption of a strict Gaussian uncertainty distribution. A statistically-robust isochron calculation method is able to retrieve this age information and to provide appropriate uncertainty estimates. Calculated ages and age uncertainties are more reliable than ISOPLOT ones for data with excess scatter.

Contaminated Gaussian distributions provide a model for a type of dataset with excess scatter relative to a strictly Gaussian-distributed one. The robust isochron method presented in this work can however be applied *in general* to data which is Gaussian-distributed only in the central spine of the uncertainty distribution, with non-Gaussian scatter occurring in the tails, arising from analytical or geological uncertainty. Indeed, it can be applied to any dataset with a well-developed spine in the data.

In most robust statistics data fitting approaches, the formal uncertainties output during data measurement are ignored. Instead, the scale used in the data fitting is derived from the scatter in the data themselves, an approach adopted by Powell et al. (2002). The new approach followed here *does* include the data measurement uncertainties, and this allows the results to converge on those of YORK when the data lack excess scatter. This provides compatibility with older “good” datasets processed

using the classical statistical approach but, going forward, allows age information to be extracted from a much wider range of datasets which might otherwise be rejected for having m_{swd} greater than the isochron cutoff.

280 A problem with SPINE, shared with YORK, is that the effect of high-leverage data is not taken into account. Such data are easily recognised in x - y plots, when a small proportion of the data—even one datapoint—is separated from the main body of the data along the trend through the data. Data fitting tends to be overly constrained to fit high-leverage data, giving them small residuals, even if the best fit of the main body of the data alone would give the high-leverage data larger residuals. In 0708, the point at highest x is relatively high leverage ($\hat{h} = 0.171$). Robust approaches have been developed to handle high leverage data, e.g. Maronna et al. (2019), ch. 5, but are not yet developed for the situation where the data uncertainties are
285 taken into account, nor for the relatively small datasets that are typical of geochronological studies (c.f. Fig. 7). Huber and Rochetti (2009), ch. 7, have a counter view advocating data assessment, rather than aiming for a black-box method to try and automatically safeguard against the potentially deleterious effect of high leverage data, an approach we suggest here. In the case of the relatively high leverage datapoint in 0708, omitting this datapoint gives 13.75 ± 0.27 Ma (compare row 11 with row 1 in the Table in Appendix A), within uncertainty of the age including this datapoint.

Here are collected some results of calculations for sample 0708, used in Fig. 7, and some related algorithmic details.

thought experiment in Fig. 1

The thought experiment sketched in Fig. 1 aimed to show the consequence for ISOPLOT behaviour of the modification of the observed data in a dataset to reduce or increase the scatter about the linear trend. The calculated equivalent of Fig. 1 for sample 295 0708 is shown in Fig. A1, including also the corresponding SPINE results. In calculating the Figure, the modification of the dataset is achieved by first taking the data points with their attendant error ellipses (i.e. covariance matrices), and moving them all in to lie on the linear trend, considered as fixed by a YORK calculation. Then the points and ellipses are considered to be displaced away from the trend. This is “move” in the Table below, going from -1, when the points lie on the trend, through 0, with the points as in the original data, to positive when displaced further away from the trend. “Move” varies more or less 300 linearly with $\sqrt{m_{swd}}$ from -0.133 at the left-edge of the Figure, to 0.064 on the right-edge; the spine width changes from 1.08 to 1.32 across the Figure. For these calculations the last line in the dataset is omitted as it is relatively high leverage ($\hat{h} = 0.171$), not wishing this datapoint to affect the results.




Figure A1. Age uncertainty ($\text{age} \pm$) plotted against $\sqrt{m_{swd}}$ under the ISOPLOT protocol for the progressively modified dataset, 0708 (see text). In model 1, the age uncertainty is constant with increasing data scatter (reflected in increasing m_{swd}), until there is a step change in age uncertainty at A when the \pm is multiplied by $\sqrt{m_{swd}}$. Then at B there is another step change with further increase of data scatter to model 2 (see text). The location of the step at A is based on a 95% confidence interval for m_{swd} (discussed below), whereas the $\sqrt{m_{swd}}$ position of the step at B is arbitrary. The y -axis is drawn at the $\sqrt{m_{swd}}$ of the actual data, 1.27 (i.e. no modification of the data). See text.

In the Figure, extending from the left, through $m_{swd} = 1$, to A, the ISOPLOT age uncertainty (model 1, i.e. YORK) is constant because the data scatter is consistent with the data uncertainties. Through this range the SPINE age uncertainty is above the

305 model 1 line because of the efficiency loss embodied in SPINE, as shown in Figs 4–5. However, after the age uncertainty steps with increasing m_{swd} , to the right of the diagram, the SPINE age uncertainty is smaller than the model 1x and model 2 age uncertainty. This is because SPINE gives an isochron on the basis of spine width up to C at $\sqrt{m_{swd}} \approx 1.3$, whereas model 2 is an errorchron, on the basis of the assumption of strictly Gaussian data uncertainties. The small steps in the SPINE age uncertainty line are an artefact of the approximation used in the calculation of the age uncertainty, see Appendix B.

310 In the top part of the following Table, the results for SPINE and for ISOPLOT are summarised. The Δ column gives the change to the age from the SPINE age, normalised by the uncertainty on the SPINE age. Below the double line in the Table, are some results from the above thought experiment, Fig. A1.

		move	$\sqrt{m_{swd}}$	age	age \pm	Δ	notes
1	SPINE			13.685	0.257	–	$s = 1.24$
2	YORK		1.296	13.733	0.216	0.37	outside 95% c.l.
3	ISOPLOT model 1x			13.733	0.280		
4	ISOPLOT model 2			13.679	0.306	–0.05	
5	siegel			13.803		–0.95	
6	SPINE	–0.133	1.1	13.769	0.252		$s = 1.08$
7	YORK			13.800	0.223	–0.39	inside 95% c.l.
8	SPINE	–0.055	1.2	13.756	0.262		$s = 1.18$
9	ISOPLOT model 1x			13.800	0.268	–0.39	
10	ISOPLOT model 2			13.832	0.300	–0.62	
11	SPINE	0	1.27	13.747	0.267	–	$s = 1.25$ $\Delta = 0.366$
12	ISOPLOT model 2			13.836	0.317	–0.65	
13	SPINE	0.024	1.3	13.743	0.267		$s = 1.26$
14	ISOPLOT model 2			13.837	0.325	–0.66	

Additional algorithmic details for ISOPLOT follow next, in part related to the above thought experiment.

315 ISOPLOT model 1

In the ISOPLOT *model 1* calculation, i.e. in YORK, a decision has to be made about the confidence interval on m_{swd} that is used to denote the range of data scatter (i.e. m_{swd}) that is considered to be accounted for by the data uncertainties, without need to either multiply the age uncertainty by $\sqrt{m_{swd}}$ (i.e. model 1x), or switch directly to an alternative calculation (which is model 2 in ISOPLOT). On the understanding that data uncertainties are correctly assigned, a one-sided confidence interval on m_{swd} can be adopted, acknowledging that m_{swd} is not being used to identify the case where assigned data uncertainties are too large or too small. The upper end of the confidence interval on m_{swd} is where excess scatter is considered to start, a conventional choice being derived from a 95% confidence interval. Note that there is no argument that this should be at $m_{swd} = 1$ (c.f.

320

Dickin, 2005, p37), unless the number of datapoints is huge. Even for a dataset of 50 datapoints the 95% confidence interval on m_{swd} extends to 1.36. In terms of the *probability of fit* measure used in ISOPLOT, this is $100 - 95 = 5\%$. It might be noted
 325 that the naming of probability of fit seems unhelpful - it is clearer to focus on m_{swd} .

ISOPLOT model 2

The so-called *error-in-variables* (e_{iv}) or measurement-error problem is avoided in YORK because the uncertainty in the x variable is taken into account explicitly. If it was not, then e_{iv} results in the calculated slope being biased downwards and the approach being inconsistent (e.g. Fuller, 1987).

330 In the ISOPLOT *model 2* calculation, e_{iv} is avoided, even though the data uncertainties are discarded, by making the slope of the line through the data be the geometric mean of the slopes of ordinary least squares of y on x , b_{yx} , and x on y , $1/b_{xy}$. These are

$$b_{yx} = \frac{\sum(x_k - \bar{x})(y_k - \bar{y})}{\sum(x_k - \bar{x})^2} \quad \text{and} \quad \frac{1}{b_{xy}} = \frac{\sum(x_k - \bar{x})(y_k - \bar{y})}{\sum(y_k - \bar{y})^2}$$

with $\bar{x} = \frac{1}{n} \sum x_k$ and $\bar{y} = \frac{1}{n} \sum y_k$. Then

335
$$b = \pm \sqrt{b_{yx} b_{xy}} = \pm \sqrt{\frac{\sum(y_k - \bar{y})^2}{\sum(x_k - \bar{x})^2}} \quad \text{and} \quad a = \bar{y} - b\bar{x}$$

with, in this case, the sign of the square root being negative. The calculation in ISOPLOT does not use this explicit formula, instead adopting an algebraic equivalent that allows the YORK iteration to be used.

ISOPLOT robust

In ISOPLOT, there is an option to use a *robust* isochron calculation method. The two available have high breakdown point but
 340 low efficiency (e.g. Huber, 1981, Sect. 1.2.3). The second method—(Siegel, 1982)—can be considered to supercede the first. In fact, here, SIEGEL is used in the implementation of SPINE as a starting point for the iteration (see Appendix C). The fit of the data for sample 0708 with SIEGEL is given in line 5 of the Table.

Appendix B: Iteration in SPINE

The SPINE algorithm involves minimising $\sum_k \rho(r_k)$ with respect to the unknown, θ , a two-element column vector, $\{a, b\}^T$ in
 345 the line equation, $y = a + bx$, in order to fit the data. The residual, r_k on datapoint k is defined below, and the function ρ is defined in (3) in the main text. The SPINE algorithm subsumes YORK.

Writing the k th datapoint as $\{x_k, y_k\}$, generally the isotopic data used in isochron calculations involve uncertainties in both x_k and y_k , and commonly the x_k and y_k are also correlated. These can be represented by a covariance matrix, V_k ,

$$V_k = \begin{bmatrix} \sigma_{x_k}^2 & \sigma_{x_k} \sigma_{y_k} \rho_{x_k y_k} \\ \sigma_{x_k} \sigma_{y_k} \rho_{x_k y_k} & \sigma_{y_k}^2 \end{bmatrix} \quad (\text{B1})$$

350 in which σ_{x_k} is the standard deviation on x_k , σ_{y_k} the standard deviation on y_k , $\sigma_{x_k}\sigma_{y_k}\rho_{x_k y_k}$ the covariance between x_k and y_k , and $\rho_{x_k y_k}$ the correlation coefficient between x_k and y_k . The covariance matrix can be represented by an ellipse around the data point in an x - y diagram, as illustrated in Fig. 2. The residual, r_k , a measure of the distance of the point $\{x_k, y_k\}$ to the line, is calculated from the coordinates of the data points, $\{x_k, y_k\}$, and their uncertainties in V_k , by

$$r_k = \frac{e_k}{\sigma_{e_k}} \quad (\text{B2})$$

355 in which e_k is the distance of the datapoint from the line, $e_k = a + bx_k - y_k$, and σ_{e_k} is the standard deviation on e_k . The standard deviation, σ_{e_k} , is calculated by error propagation using V_k :

$$\sigma_{e_k}^2 = \left\{ \frac{\partial e_k}{\partial x_k}, \frac{\partial e_k}{\partial y_k} \right\} V_k \left\{ \frac{\partial e_k}{\partial x_k}, \frac{\partial e_k}{\partial y_k} \right\}^T = b^2 \sigma_{x_k}^2 + \sigma_{y_k}^2 - 2b \sigma_{x_k} \sigma_{y_k} \rho_{x_k y_k} \quad (\text{B3})$$

with the term in curly brackets evaluating to $\{b, -1\}$. The residual is then

$$r_k = \frac{e_k}{\sigma_{e_k}} = \frac{a + bx_k - y_k}{\sqrt{b^2 \sigma_{x_k}^2 + \sigma_{y_k}^2 - 2b \sigma_{x_k} \sigma_{y_k} \rho_{x_k y_k}}} \quad (\text{B4})$$

360 In matrix form, the residuals can be written as a column vector, \mathbf{r}

$$\mathbf{r} = \mathbf{W}_e \mathbf{e} = \mathbf{W}_e (\mathbf{X}\boldsymbol{\theta} - \mathbf{y}) \quad (\text{B5})$$

with \mathbf{W}_e a diagonal matrix with kk th element, $1/\sigma_{e_k}$.

The minimisation of $\sum_k \rho(r_k)$ is iterative, starting from a robust but low efficiency estimate of the line, for example using Siegel (1982). The minimisation of $\sum \rho(r_k)$ is undertaken using the fact that, at the minimum, the derivative of $\sum \rho(r_k)$ with respect to θ is zero. Defining

$$2\psi(r_k) = \frac{\partial \rho(r_k)}{\partial r_k} \quad (\text{B6})$$

this function, for the Huber (1981) approach, from (3), is

$$\psi(r_k) = \begin{cases} -h & r_k < -h \\ r_k & \text{if } -h < r_k < h \\ h & r_k > h \end{cases} \quad (\text{B7})$$

For YORK, $\psi(r_k) = r_k$, equivalent to SPINE with large h .

370 At the minimum of $\sum \rho(r_k)$

$$\sum_k \frac{\partial \rho(r_k)}{\partial \theta} = 0 = \sum_k \left(\frac{\partial \rho(r_k)}{\partial r_k} \right) \left(\frac{\partial r_k}{\partial \theta} \right) = \sum_k \psi(r_k) \left(\frac{\partial r_k}{\partial \theta} \right) = \sum_k \psi(r_k) \frac{1}{\sigma_{e_k}} \left\{ \left(\frac{\partial e_k}{\partial \theta} \right) - r_k \left(\frac{\partial \sigma_{e_k}}{\partial \theta} \right) \right\} \quad (\text{B8})$$

In the curly braces, writing the first derivative, $\{1, x_k\}$, as the k th row of a matrix \mathbf{X} , and writing the first and second derivatives together as the k th row of a matrix \mathbf{X}' , then the k th row of \mathbf{X}' is

$$X'_k = \{1, x'_k\} = \left\{ 1, x_k - \frac{r_k}{\sigma_{e_k}} (b \sigma_{x_k}^2 - \sigma_{x_k} \sigma_{y_k} \rho_{x_k y_k}) \right\}$$

375 with b the slope of the line. The second column of \mathbf{X} is simply the data x values, while the second column of \mathbf{X}' is the x values on the line where the uncertainty ellipses around each datapoint would touch it, \mathbf{x}' . In matrix form, (B8) can be written as

$$\mathbf{X}'^T \mathbf{W}_e \psi(\mathbf{r}) = 0 \quad (\text{B9})$$

in which $\psi(\mathbf{r})$ is a column vector whose k th element is $\psi(r_k)$. B9 constitutes two non-linear equations requiring iteration to solve. Two iteration schemes are proposed. The first iteration, in fact used for all the simulations, proceeds directly from (B9),
380 while the second iteration is in iteratively reweighted least squares form (e.g. Maronna et al., 2019, Sect. 4.5.2).

In the first scheme, at iteration i , progressing towards the minimum with $\theta_i = \theta_{i-1} + \Delta\theta$, for $|r_k| < h$, $\psi_i(r_k) = \psi_{i-1}(r_k) + X_k \Delta\theta / \sigma_{e_k}$ and $\psi_i(r_k) = \psi_{i-1}(r_k)$ otherwise. This can be written $\psi_i(r_k) = \psi_{i-1}(r_k) + \dot{\psi}_{i-1}(r_k) X_k \Delta\theta / \sigma_{e_k}$, in which $\dot{\psi}(r_k) = \partial\psi(r_k) / \partial r_k$, which is 1 for $|r_k| < h$, and 0 otherwise, from (B7). Substituting into (B9) gives

$$\mathbf{X}'^T \mathbf{W}_e (\mathbf{W}_e \mathbf{I}' \mathbf{B} \Delta\theta + \psi(\mathbf{r})) = 0 \quad (\text{B10})$$

385 dropping iteration subscripts, with $\mathbf{I}' = \text{diag}(\dot{\psi}(\mathbf{r}))$ a modified identity matrix with its kk th element equal to $\dot{\psi}(r_k)$. Equation (B10) can then be rearranged to give $\Delta\theta$ at the current iteration

$$\Delta\theta = -(\mathbf{X}'^T \mathbf{W}_e^2 \mathbf{I}' \mathbf{X})^{-1} \mathbf{X}'^T \mathbf{W}_e \psi(\mathbf{r}) \quad (\text{B11})$$

Iteration proceeds until $\Delta\theta$ approaches 0.

The second iteration scheme is the iteration implemented in the python code. Such iterations are known to be stable, e.g.
390 Maronna et al. (2019). However the logic of the first scheme is needed to obtain the covariance matrix of θ , below. In the second scheme, a weight function, $w(r_k)$, is introduced so that the resulting equations have a simple least squares form. For (B9), this can be done by defining $w(r_k) = \psi(r_k) / r_k$, so that $\psi(r_k) = r_k w(r_k)$, introducing \mathbf{r} explicitly into (B9). Then

$$\mathbf{X}'^T \mathbf{W} (\mathbf{X}\theta - y) = 0 \quad (\text{B12})$$

in which the kk th element of the diagonal matrix, \mathbf{W} , is $w(r_k) / \sigma_{e_k}^2$, combining the weighting from w with the weighting from
395 the data uncertainties, \mathbf{W}_e . By rearranging (B12), the repeated substitution solution to (B9) is given by

$$\theta = (\mathbf{X}'^T \mathbf{W} \mathbf{X})^{-1} \mathbf{X}'^T \mathbf{W} y \quad (\text{B13})$$

Iteration is needed as \mathbf{X}' and \mathbf{W} depend slightly on θ , given a sensible starting point for the iteration as provided by SIEGEL.

Accepting that an isochron has been calculated, the covariance matrix of θ , \mathbf{V}_θ , can be calculated by error propagation of the data to θ using the logic of the first iteration scheme. First it is convenient to transform the problem solved by (B11) to an
400 identical one in which the data are represented by \mathbf{e} rather than $\{\mathbf{x}, \mathbf{y}\}$. First note that (B9) involves \mathbf{r} , a scalar for each data point, the uncertainty on \mathbf{e} in \mathbf{W}_e , and the x position on the best-fit line where the ellipse around each datapoint touches the line, at \mathbf{x}' , the second column of \mathbf{X}' . Given the best-fit θ , the y corresponding to \mathbf{x}' , is $y' = \mathbf{X}'\theta$. Defining $\mathbf{y}'' = \mathbf{y}' - \mathbf{e}$, data involving $\{\mathbf{x}', \mathbf{y}''\}$, rather than $\{\mathbf{x}, \mathbf{y}\}$, is an identical problem via (B9). In this identical problem, \mathbf{x}' and \mathbf{y}' are fixed (have no uncertainty). With this, the change in θ corresponding to (B9), becomes

$$405 \quad \Delta\theta = -(\mathbf{X}'^T \mathbf{W}_e^2 \mathbf{I}' \mathbf{X}')^{-1} \mathbf{X}'^T \mathbf{W}_e \psi(\mathbf{r}) \quad (\text{B14})$$

in terms only of \mathbf{X}' and not involving \mathbf{X} . This is akin to the transform in York et al. (2004) to convert the \mathbf{V}_θ of York (1969) to that of the \mathbf{V}_θ of Titterton and Halliday (1979).

Assuming that θ is approximately linear in each e_k around the minimum in $\sum \rho(r_k)$, then

$$\mathbf{V}_\theta = \left(\frac{\partial \theta}{\partial \mathbf{e}} \right) \mathbf{V}_e \left(\frac{\partial \theta}{\partial \mathbf{e}} \right)^T \quad (\text{B15})$$

410 At the solution of (B9), $\Delta\theta = 0$, so, by the chain rule using (B14)

$$\left(\frac{\partial \theta}{\partial \mathbf{e}} \right) = \left(\frac{\partial \Delta\theta}{\partial \theta} \right)^{-1} \left(\frac{\partial \Delta\theta}{\partial \mathbf{e}} \right) = (\mathbf{X}'^T \mathbf{W}_e^2 \mathbf{I}' \mathbf{X}')^{-1} \mathbf{X}'^T \mathbf{W}_e^2 \mathbf{I}' \quad (\text{B16})$$

Substituting into (B15), cancelling and using $\mathbf{W}_e^2 \mathbf{I}' \mathbf{W}_e^{-2} \mathbf{I}' \mathbf{W}_e^2 = \mathbf{W}_e^2 \mathbf{I}'$

$$\mathbf{V}_\theta = (\mathbf{X}'^T \mathbf{W}_e^2 \mathbf{I}' \mathbf{X}')^{-1} \quad (\text{B17})$$

The small steps in the age uncertainty (age \pm) curve in Fig A1 arise when diagonal elements in \mathbf{I}' change from one to zero as
415 mswd increases.

In YORK (or if all $|r_k| < h$ in SPINE), then $\mathbf{I}' = \mathbf{I}$ and $\psi(\mathbf{r}) = \mathbf{r}$. So

$$\Delta\theta = -(\mathbf{X}'^T \mathbf{W}_e^2 \mathbf{X})^{-1} \mathbf{X}'^T \mathbf{W}_e^2 \mathbf{r} \quad (\text{B18})$$

and the covariance matrix becomes

$$\mathbf{V}_\theta = (\mathbf{X}'^T \mathbf{W}_e^2 \mathbf{X}')^{-1} \quad (\text{B19})$$

420 If, in addition, all $\sigma_{x_k} = 0$ then $\mathbf{X}' = \mathbf{X}$, and iteration is not involved, \mathbf{e} is replaced by $-\mathbf{y}$, and

$$\theta = (\mathbf{X}^T \mathbf{W}_e^2 \mathbf{X})^{-1} \mathbf{X}^T \mathbf{W}_e^2 \mathbf{e} \quad (\text{B20})$$

with a covariance matrix of

$$\mathbf{V}_\theta = (\mathbf{X}^T \mathbf{W}_e^2 \mathbf{X})^{-1} \quad (\text{B21})$$

These are the results for fitting data by simple weighted least squares.

425 **Appendix C: SPINE Python code**

The second iteration in Appendix B is coded in the Python function, `huber2`. The starting point for the iteration is provided by the function, `siegel` (Siegel, 1982). The calling function, `recipe`, is a placeholder for a more general function to be written by the user.

```
import sys
430 import datetime
```

```

import numpy as np

out = open("out.txt", "w") # opening output file
screen = sys.stdout
435 standard = [screen, out] # default where print goes

default_h = 1.4 # default h in huber

def siegel(data): # siegel (1982)
440     n = data.shape[0];
        (x, sdx, y, sdy, cor) = np.transpose(data)
        x += 1e-8 * np.random.random(n) # naive breaking of x ties
        med = np.empty(n);
445         for i in range(n):
            col = np.empty(n);
            for j in range(n):
                if i is not j: col[j] = (y[j] - y[i])/(x[j] - x[i])
            med[i] = np.median(np.delete(col, i))
        b = np.median(med)
450     return np.array((np.median(y - x * b), b))

# -----

def calcage(theta, covtheta = None, method = 1):
455 # tera-wasserburg: focus on lower intercept
        dc238 = 1.55125e-10; dc235 = 9.8485e-10
        tc = 0; t = t1 = 1500.; tdiff = 1e-4
        (a, b) = theta
        k = 0
460     while k < 12 and abs(t - tc) > tdiff:
            tc = t; t = 1/dc238 * np.log(1 + 1/t1)
            t1 = (1/137.8 * (np.exp(dc235 * t) - 1) / (np.exp(dc238 * t) - 1) - a) / b
            k += 1
        if covtheta is None:
465             sdage = 0
        else:
            den = b * np.exp(dc238 * t) * dc238 + \
                (-np.exp(dc235 * t) * dc235 + np.exp(dc235 * t + dc238 * t) * dc235 + \
                 np.exp(dc238 * t) * dc238 - np.exp(dc235 * t + dc238 * t) * dc238)/137.8
470             jac = [(np.exp(dc238 * t) - 1)/den * (np.exp(dc238 * t) - 1), (np.exp(dc238 * t) - 1)/den]
            sdage = np.sqrt(np.dot(np.dot(jac, covtheta), np.transpose(jac)))/1e6
        return (t/1e6, sdage)

# -----

```

475

```

def huber2(data0, h = 1.4):
#   iteratively-reweighted huber line-fitter
    n = data0.shape[0]
    itmax = 20; mindel = 1e-8; mincond = 1e-12; minsump = 0.01
480   (x, sdx, y, sdy, cor) = np.transpose(data0)
    avx = np.dot(x, np.ones(n))/n;   avy = np.dot(y, np.ones(n))/n;
    div = np.array([1/avy, avx/avy])
    data = np.copy(data0)
    (x, sdx, y, sdy, cor) = np.transpose(data)
485   x /= avx; sdx /= avx; y /= avy; sdy /= avy; cov = sdx*sdy*cor
    theta = siegel(data);
    k = 0; code = 0; deltheta = (1e10, 1e10)
    while k < itmax and ( np.sqrt(np.dot(deltheta,deltheta)) > mindel):
        k += 1; (a, b) = oldtheta = theta
490   e = a + b * x - y;   sde = np.sqrt(b**2*sdx**2 - 2*b*cov + sdy**2)
        r = e/sde
        wh = [1 if abs(rk) < h else np.sqrt(h/abs(rk)) for rk in r]/sde
        xp = x - r * (b*sdx**2 - cov)/sde           # x on the line
        ypp = (y - e - r*(b*cov - sdy**2)/sde)*wh   # W^(1/2) (y'-e): y off the line
495   c = np.transpose([wh, xp*wh])                 # W^(1/2) X'
        (u, s, v) = np.linalg.svd(c, full_matrices=False)
        if mincond * s[0] > s[1]: code = -1; break  # (nearly) singular matrix
        theta = np.dot(np.dot(np.dot(np.transpose(v), np.diag(1/s)), np.transpose(u)), ypp)
        deltheta = theta - oldtheta
500   if k == itmax: code = 1 # not converged
        pc = np.dot([wh**2, xp*wh**2], e);
        sump = np.sqrt(np.dot(pc, pc));
        if sump > minsump: code = 2 # not solved the nle
        dpsi = [1 if abs(rk) < h else 0 for rk in r]
505   covtheta = np.linalg.inv(np.dot(np.dot(np.transpose(c), np.diag(dpsi)), c)) / \
        np.array([[div[0]**2, div[0] * div[1]], [div[0] * div[1], div[1]**2]])
    return (code, theta/div, covtheta, sump, k, s[1])

# -----
510   def pr(s, e="\n", printto=standard):
#   prints a string
        for pr in printto: print(s, end=e, file=pr)

515   def recipe(title, data, where = [screen]):
#   prototype calculation driver
        h = default_h
        (x, sdx, y, sdy, cor) = np.transpose(data)

```

```

n = data.shape[0]
520 today = datetime.datetime.now();
pr("=====\n" + \
    "running spine.py on "+today.ctime())
res = huber2(data)
if res[0] != 0:
525     print("sample "+title+" not calculated")
        return "bad"
(a, b) = theta = res[1]
ucovtheta = res[2]
(age, sdage) = calcage(theta, ucovtheta)
530 e = a + b * x - y
sde = np.sqrt(b**2 * sdx**2 - 2 * b * sdx * sdy * cor + sdy**2)
r = e/sde
s = nmad(r); slim = 1.92 - 0.162 * np.log(10 + n);
if s < slim:
535     pr(("sample "+title+": s = %0.2f"+ ": isochron " + "age = %0.3f +/- %0.3f Ma") % \
        (s, age, 1.96 * sdage), printto = where)
    else:
        pr(("sample "+title+": s = %0.2f"+ ": errorchron " + "age = %0.3f Ma") % \
            (s, age), printto = where)
540     return [0, age, sdage, theta]

# -----

# data rows: x sdx y sdy cor
545 data = np.loadtxt("data0708.txt", delimiter=",")

recipe("0708", data, where=standard); print()

Datafile for sample 0708, data0708.txt, see Fig. 6

73.2064, 1.12543, 0.753, 0.0075, -0.068224
550 260.417, 4.06901, 0.435, 0.01, 0.18853
169.205, 2.43357, 0.577, 0.01, 0.45644
79.1766, 0.908995, 0.751, 0.01, 0.19585
212.766, 2.94251, 0.473, 0.0085, 0.37494
154.56, 1.79165, 0.615, 0.0105, 0.22567
555 217.391, 2.83554, 0.474, 0.0095, 0.27879
209.644, 4.83455, 0.484, 0.011, 0.10969
144.092, 2.49151, 0.647, 0.0105, 0.10222
174.216, 1.97283, 0.584, 0.009, 0.21416
224.215, 3.26771, 0.477, 0.012, 0.26464
560 236.407, 3.35329, 0.461, 0.0115, 0.14718
161.551, 2.87086, 0.628, 0.011, -0.041164

```

265.252, 5.62869, 0.385, 0.0115, -0.043072
 152.439, 2.32377, 0.608, 0.0095, 0.019543
 151.745, 2.30266, 0.626, 0.0105, 0.0030752
565 101.112, 1.17572, 0.727, 0.0085, 0.038467
 265.182, 3.41058, 0.427, 0.0095, 0.070384
 286.78, 3.16634, 0.391, 0.0085, 0.24059
 287.604, 3.5568, 0.365, 0.008, -0.041298
 264.69, 2.83747, 0.419, 0.008, 0.15939
570 274.574, 2.67638, 0.378, 0.008, 0.17686
 212.314, 4.28235, 0.501, 0.013, 0.34121
 161.29, 1.69095, 0.591, 0.009, 0.14398
 140.647, 1.58253, 0.633, 0.0105, 0.18753
 183.15, 2.18036, 0.557, 0.0105, 0.49436
575 218.818, 2.39407, 0.53, 0.012, 0.21725
 312.5, 5.3711, 0.334, 0.01, -0.080507
 227.79, 2.20526, 0.49, 0.0105, 0.073412
 212.766, 2.71616, 0.507, 0.011, 0.25544
 139.276, 2.03676, 0.652, 0.01, 0.44912
580 179.533, 2.25625, 0.556, 0.0105, 0.24303
 224.215, 2.51362, 0.48, 0.0085, 0.11165
 219.78, 3.62275, 0.519, 0.0115, 0.31258
 165.017, 2.8592, 0.59, 0.0115, 0.03575
 145.773, 2.33746, 0.61, 0.0105, 0.098985
585 239.808, 4.02556, 0.442, 0.0095, -0.10478
 187.266, 2.80548, 0.538, 0.0125, 0.11864
 255.232, 2.63831, 0.443, 0.0085, 0.025538
 141.443, 4.10124, 0.616, 0.011, -0.15512
 115.34, 0.997753, 0.707, 0.01, 0.15087
590 117.509, 1.17371, 0.68, 0.008, 0.33891
 73.6377, 0.623589, 0.757, 0.0065, 0.57883
 160.256, 2.31139, 0.605, 0.012, 0.16382
 149.701, 1.56872, 0.619, 0.009, 0.24589
 245.278, 2.58694, 0.452, 0.008, 0.33748
595 251.256, 4.10343, 0.435, 0.014, 0.3589
 130.208, 2.11928, 0.666, 0.011, 0.18376
 276.243, 4.19706, 0.419, 0.01, 0.16668
 298.508, 4.90087, 0.359, 0.0105, -0.090909
 381.679, 6.40989, 0.241, 0.013, 0.0073965

600 Example output, running on the command line

```

running spine.py on Mon Sep 14 14:10:55 2020
sample 0708: s = 1.24: isochron age = 13.685 +/- 0.257 Ma
  
```


Appendix D: Simulation setup

This work was originally motivated by the dating of speleothems using the lower intercept with a U-Pb Concordia in Tera-
605 Wasserburg style plots (Woodhead et al., 2012). This paper therefore discusses $\{x, y\}$ data with the expectation that
 $x = {}^{238}\text{U}/{}^{206}\text{Pb}$ and $y = {}^{207}\text{Pb}/{}^{206}\text{Pb}$, but the logic and the algorithm are in no way restricted to this system.

10,000 simulated datasets, each containing 5, 6, 8, 10 and 15 datapoints, respectively, were used to assess SPINE. Each
dataset corresponds to an age of 4 Ma, with an underlying trend chosen to be $y = 0.811 - 0.000474737x$. For each dataset,
the x -values were drawn from a uniform probability distribution with bounds, $\{400, 1100\}$ (so the x are not equi-spaced).
610 Datapoints are assigned uncertainty with $\sigma_{x_k} = 0$ and a fixed $\sigma_{y_k} = 0.00125$, the latter representing the analytical uncertainty,
propagated from both the x and y measurement into y . In a real $\{{}^{238}\text{U}/{}^{206}\text{Pb}, {}^{207}\text{Pb}/{}^{206}\text{Pb}\}$ dataset, σ_{x_k} and σ_{y_k} would be
finite and correlated. However, this makes no difference to the calculations once data are processed into r_k form as in Fig. 2.
For a given dataset, scatter is introduced into the data by drawing the y values from an uncertainty distribution, centred on
the underlying trend, that may be either Gaussian (N) or one of three contaminated Gaussian distributions—5%3N, 25%5N,
615 or 10%10N—as in Powell et al. (2002). For $n = 10$ and gaussian-distributed uncertainties, the age uncertainty obtained is
approximately $\sigma_t = 0.01$ Ma.

Results are presented in terms of kernel density estimates using an Epanechnikov kernel (Wand and Jones, 1995). Kernel
density estimates (kde) are a way of presenting data that could otherwise be plotted as a histogram, normally normalised
so that—like a probability distribution—the area under the kde curve is 1. The smoothness of the kde is controlled by a
620 smoothing constant whose value was chosen to be just large enough for the kde to appear smooth, given that 10,000 datasets
are used in each kde.

Appendix E: SPINE handling outliers

To see the consequence of outliers stemming from residuals from a contaminated Gaussian distribution, a dataset from simu-
lations is shown in Fig. E1 and its corresponding quantile-quantile plot in Fig. E2. This simulation, with $n = 50$ and 25%3N
625 and a true age of 4.00 Ma, follows the approach in Appendix D, but the uncertainty on the y values is taken to be an order of
magnitude larger (to be comparable with sample 0708). The shape of Fig. E2 is typical of contaminated Gaussian distributions
in samples of this size, with outliers lying outside the band between the blue dashed lines. The SPINE age is 3.95 Ma, whilst
the YORK age is higher, 4.11 Ma, $\Delta = +3.0\sigma$ of the SPINE age, due to the effect of the outliers. $M_{\text{swd}} = 4.35$, so ISOPLOT
would give a model 2 age, 4.19 Ma, with $\Delta = 4.27$, being more affected by the outliers than YORK.

630 In general, with larger contaminated-Gaussian datasets, the difference between the SPINE and YORK ages are not dramatic,
even if there are obvious outliers in the quantile-quantile plots. In 10,000 simulated datasets with $n = 50$ and 25%3N, a 95%
confidence interval on Δ is ± 1.5 .




Figure E1. A 25%3N simulation with $n = 50$, with the SPINE fit in red, the YORK fit in blue. The 2σ uncertainty applied to the y value of each datapoint is indicated in the top right of the figure. See text.




Figure E2. Quantile-quantile plot for example in Fig. E1. See text.

Author contributions. Roger Powell created the approach and coded the Python script; Eleanor Green helped validate the maths/statistics and write the paper; Tephy Marillo Sialer helped with the simulations; and Jon Woodhead oversaw the applicability of the approach.

635 *Competing interests.* The authors declare that they have no conflict of interest.

Acknowledgements. We would like to thank the anonymous reviewers for their work, particularly reviewer 4 (times two). Tim Pollard has materially helped in our understanding of the ISOPLOT code functionality. JW is funded by Australian Research Council grant FL160100028.

References

- Brooks, C., Hart, S.R., and Wendt, I.: Realistic use of two-error regression treatments as applied to Rubidium-Strontium data. *Reviews of Geophysics and Space Physics*, 10, 551-577, 1972.
- Dickin, A.P.: Radiogenic isotope geology. Cambridge University Press, 492pp, 2005.
- Fox, J.: Applied regression analysis & Generalised linear models. 3rd ed. Sage, Los Angeles, 791pp: 2016.
- Fuller, W.A.: Measurement error models. John Wiley and Sons, 440pp: 1987.
- Hampel, F.R., Rousseeuw, P.J., Ronchetti, E.M., and Stahel, W.A.: Robust statistics. Wiley and Sons, New York, 502pp, 1986.
- Huber, P.J.: Robust Statistics, John Wiley and Sons, Inc., New York: 305pp, 1981.
- Huber, P.J., and Ronchetti, E.M.: Robust Statistics, John Wiley and Sons, Inc., New York: 354pp, 2009.
- Ludwig, K. R.: Isoplot/Ex Version 3.75: A Geochronological Toolkit for Microsoft Excel. Special Publication 4, Berkeley Geochronology Center, 75pp, 2012.
- Maronna, R.A., Martin, D., Yohai, V.J. and Salibián-Barrera, M.: Robust statistics. John Wiley and Sons, Chichester, 430pp, 2019.
- McLean, N.M.: Straight line regression through data with correlated uncertainties in two or more dimensions. *Geochimica et Cosmochimica Acta*, 124, 237–249, 2014.
- Powell, R., Woodhead, J., and Hergt, J.: Improving isochron calculations with robust statistics and the bootstrap. *Chemical Geology* 185, 191–204, 2002.
- Reiners, P.W., Carlson, R.W., Renne, P.R., Cooper, K.M., Granger, D.E., McLean, N.M., Schoene, B.: *Geochronology and Thermochronology*. John Wiley & Sons, 464pp, 2018.
- Siegel, A.F.: Robust regression Using repeated medians. *Biometrika*, 69, 242–244, 1982.
- Titterton, D.M., and Halliday, A.N.: On the fitting of parallel isochrons and the method of least squares. *Chemical Geology*, 26, 183–195, 1979.
- Tukey, J.W.: A survey of sampling from contaminated distributions, in: *Contributions to Probability and Statistics*, I. Olkin, Ed., Stanford University Press, 1960.
- Vermeesch, P.: IsoplotR: a free and open toolbox for geochronology. *Geoscience Frontiers*, 9, 1479–1493, 2018.
- Wand, M.P. and Jones, M.C.: Kernel smoothing. Chapman and Hall, London. 212pp, 1995.
- Wendt, I., and Carl, C.: The statistical distribution of the mean squared weighted deviation. *Chemical Geology (Isotope Geosciences Section)* 86, 275–285, 1991.
- Woodhead, J., Hand, S.J., Archer, M., Graham, I., Sniderman, K., Arena, D.A., Black, K.H., Godhelp, H., and Price, E.: Developing a radiometrically-dated chronologic sequence for Neogene biotic change in Australia, from the Riversleigh World Heritage Area of Queensland. *Gondwana Research* 29, 153–167, 2016.
- Woodhead, J., Hellstrom, J., Pickering, R., Drysdale, R., Paul, B., and Bajo, P.: U and Pb variability in older speleothems and strategies for their chronology. *Quaternary Geochronology* 14, 105–113, 2012.
- Woodhead, J. and Petrus, J., Exploring the advantages and limitations of in situ U-Pb carbonate geochronology using speleothems, *Geochronology* 1, 67–84, <https://doi.org/10.5194/gchron-1-69-2019>.
- York, D.: Least squares fitting of a straight line. *Canadian Journal of Physics*, 44, 1079–1086: 1966
- York, D.: Least squares fitting of a straight line with correlated errors. *Earth and Planetary Science Letters* 5, 320–324, 1969.

675 York, D., Evensen, N.M., Martinez, M.L., and Delgado, J.D.: Unified equations for the slope, intercept, and standard errors of the best straight line. *American Journal of Physics* 72, 367–375, 2004.