

Cosmogenic nuclide and solute flux data from central Cuban rivers emphasize the importance of both physical and chemical mass loss from tropical landscapes

Mae Kate Campbell^{1,2}, Paul R. Bierman^{2,3}, Amanda H. Schmidt⁴, Rita Sibello Hernández⁵, Alejandro
5 García-Moya⁵, Lee B. Corbett³, Alan J. Hidy⁶, Héctor Cartas Águila⁵, Aniel Guillén Arruebarrena⁵,
Greg Balco⁷, David Dethier⁸, Marc Caffee⁹

¹Department of Geology, University of Vermont, Burlington, VT 05405, USA

²Gund Institute for Environment, University of Vermont, Burlington, VT 05405

³Rubenstein School of the Environment and Natural Resources, the University of Vermont, Burlington,
10 VT 05405, USA

⁴Department of Geosciences, Oberlin College, Oberlin, OH 44074, USA

⁵Centro de Estudios Ambientales de Cienfuegos, Departamento de Estudio de la Contaminación
Ambiental. AP 5, 59350, Ciudad Nuclear, Cienfuegos, Cuba

⁶Atmospheric, Earth, and Energy Division, Lawrence Livermore National Laboratory, Livermore, CA
15 94550, USA

⁷Berkeley Geochronology Center, Berkeley, CA 94709, USA

⁸Department of Geosciences, Williams College, Williamstown, MA 01267, USA

⁹Department of Physics and Astronomy and Department of Earth, Atmospheric, and Planetary Sciences,
Purdue University, West Lafayette, IN 47907, USA

20 *Correspondence to:* Amanda H. Schmidt (aschmidt@oberlin.edu)

Abstract

We use 25 new measurements of *in situ* produced cosmogenic ^{26}Al and ^{10}Be in river sand, paired with estimates of dissolved load flux in river water, to characterize the processes and pace of landscape change in central Cuba. Long-term erosion rates inferred from ^{10}Be concentrations in quartz extracted from central Cuban river sand range from 3.4-189 $\text{Mg km}^{-2} \text{y}^{-1}$ (mean = 59, median = 45). Dissolved loads (10-176 $\text{Mg km}^{-2} \text{y}^{-1}$; mean = 92, median = 97), calculated from stream solute concentrations and modelled runoff, exceed measured cosmogenic ^{10}Be -derived erosion rates in 18 of 23 basins. This disparity mandates that landscape-scale, in this environment, mass loss is not fully represented by the cosmogenic nuclide measurements.

The $^{26}\text{Al}/^{10}\text{Be}$ ratios are lower than expected for steady-state exposure or erosion in 16 of 24 samples. Depressed $^{26}\text{Al}/^{10}\text{Be}$ ratios occur in many of the basins that have the greatest disparity between dissolved loads (high) and erosion rates inferred from cosmogenic nuclide concentrations (low). Depressed $^{26}\text{Al}/^{10}\text{Be}$ ratios are consistent with the presence of a deep, mixed, regolith layer providing extended storage times on slopes and/or burial and extended storage during fluvial transport. River water chemical analyses indicates many basins with lower $^{26}\text{Al}/^{10}\text{Be}$ ratios and high ^{10}Be ratios are underlain at least in part by evaporitic rocks that rapidly dissolve.

Our data show that when assessing mass loss in humid tropical landscapes, accounting for the contribution of rock dissolution at depth is particularly important. In such warm, wet climates, mineral dissolution can occur many meters below the surface, beyond the penetration depth of most cosmic rays, and thus the production of most cosmogenic nuclides. Our data suggest the importance of estimating solute fluxes and measuring paired cosmogenic nuclides to understand better the processes and rates of mass transfer at a basin-scale.

1 Introduction

Cosmogenic nuclide concentrations of river sand have been used to quantify rates of landscape change (often termed erosion rates) since the 1990s (Brown et al., 1995a; Granger et al., 1996; Bierman and Steig, 1996; Portenga and Bierman, 2011; Codilean et al., 2018). Accurately establishing long-term

rates of change provides an important context for understanding the effects of human activity on erosion (Reusser et al., 2015; Nearing et al., 2017), and for other common applications of cosmogenic nuclides at the basin-scale, such as quantifying the effect of tectonics (Scherler et al., 2014), climate (Marshall et al., 2017), and baselevel change (Reinhardt et al., 2007) on rates of landscape change over time.

¹⁰Be-derived rates of landscape change at a drainage-basin scale are often implicitly assumed to reflect both physical and chemical mass loss, the sum of which is termed denudation (Regard et al., 2016). However, this assumption is only valid if all mass loss from the landscape occurs within the uppermost meter or two of Earth's surface, the penetration depth of the cosmic-ray neutrons responsible for producing most cosmogenic nuclides via spallation reactions (Bierman and Steig, 1996). Deeper mass loss by rock dissolution remains largely undetected by cosmogenic nuclide analysis. Failure to account for rock dissolution at depth and the export of mass as dissolved load below the spallation-dominated nuclide production zone (~2 m) may bias cosmogenic nuclide-derived estimates of denudation (Small et al., 1999; Riebe et al., 2001a; Dixon et al., 2009a) on the low side. Incorrectly determined erosion rates can derail attempts to understand landscape evolution, soil production, and climate interaction with surface processes (Riebe et al., 2003).

Rock dissolution at depth is a major process in areas with significant groundwater-rock interactions; connecting denudation rates to landscape change requires consideration of this process. This includes any landscape where the physical removal of mass is slow, allowing for prolonged water-rock interactions, such as low-relief landscapes (Ollier, 1988). Some landscape characteristics facilitate or are the result of extensive water-rock interaction: thick saprolite (Dixon et al., 2009a), extensively jointed/fractured bedrock (Ollier, 1988), and readily soluble rocks, including carbonate (Pope, 2013) and evaporite deposits. Conditions in the humid tropics favor prolonged and extensive water rock-interaction and include the absence of recent glaciation (Modenesi-Gauttieri et al., 2011), the presence of active groundwater flow systems year-round (Ollier, 1988), and large amounts of precipitation.

Rock dissolution rates in the tropics can be among the highest globally (Pope, 2013); yet, global compilations of cosmogenic nuclide data from river sand suggest rates of landscape change in the tropics are slower than in most other climate zones (Portenga and Bierman, 2011). This dichotomy is consistent with cosmogenic rates significantly underestimating landscape denudation in areas where

deep rock dissolution is ubiquitous. Only a few studies focused in the tropics compare nuclide-derived rates to measurements of dissolved load flux in streams (e.g. Salgado et al., 2006; Hinderer et al., 2013; Regard et al., 2016). As the use of cosmogenic nuclides to measure rates of landscape change in the tropics expands (e.g. Cherem et al., 2012; Barreto et al., 2013; Derrieux et al., 2014; Mandal et al., 80 2015; Sosa Gonzalez et al., 2016a; Jonell et al., 2017), considering the potential influence of rock dissolution at depths below the production of most cosmogenic nuclides becomes more important.

Here, we present measurements of *in situ* ^{26}Al and ^{10}Be in riverine quartz, along with estimates of dissolved loads, in humid, tropical central Cuba (Bierman et al., 2020). With these data, we explore the relationships between cosmogenic nuclide concentrations, dissolved load fluxes, and landscape- 85 scale parameters at a basin scale in a humid tropical location where mass is being lost from the landscape by multiple, different processes from a variety of rock types. We characterize the rates and processes by which the Cuban landscape is changing, and place these data in a global context. Our findings illustrate the importance of considering rock dissolution when using cosmogenic nuclides to assess rates of landscape change in areas with the potential for significant mass loss by solution at 90 depth, and provide a geologic baseline for assessing the impact of human actions on the Cuban landscape.

2 Background

Terminology referring to mass loss from watersheds has been applied ambiguously in the past and can be confusing. Here, we refer to the tempo of landscape mass loss calculated from ^{26}Al and ^{10}Be 95 concentrations as erosion rates; these rates include all processes (physical and chemical) removing mass within ~ 2 m of Earth's surface. We refer to rates of landscape mass loss inferred from measurements of stream water chemistry, convolved with estimates of annual run-off volumes, as rock dissolution rates. We use the term denudation to refer to total mass loss from sampled catchments. All of these rates are expressed in terms of mass per time per area ($\text{Mg km}^{-2} \text{y}^{-1}$), which can be converted to depth over time 100 by assuming a rock density.

2.1 Quantifying basin mass loss with cosmogenic nuclides: approaches and limitations

Landscape-scale denudation occurs through both physical removal of mass (erosion) and chemical dissolution of minerals in rocks. Sediment, produced from eroding bedrock, travels downslope towards base level, whereas rock dissolution moves mass in solution from the landscape to rivers, and then to the ocean. Measurement of cosmogenic nuclides in river sediment can be used to infer the spatially averaged erosion rate of a drainage basin (Brown et al., 1995a; Granger et al., 1996; Bierman and Steig, 1996). In a basin that is steadily eroding, the concentration of cosmogenic nuclides in a sediment sample reflects the rate at which overlying mass at and near the surface was removed as the material was exhumed, through both physical mass loss and rock dissolution (Lal, 1991). Cosmogenic erosion rates are equivalent to denudation rates if, and only if, rock dissolution only occurs within a meter or two of the surface—the depth of penetration for neutrons which produce most cosmogenic nuclides. If rock dissolution occurs below the neutron penetration depth, erosion rates calculated from measured nuclide concentrations will underestimate denudation.

Measuring multiple cosmogenic nuclides with different half lives in the same sample can provide more information on the near-surface history of surface materials, such as soil mixing depth and residence time (Lal and Chen, 2005), as well as sediment storage within the watershed (Granger and Muzikar, 2001). The production ratio of $^{26}\text{Al}/^{10}\text{Be}$ at the surface at mid- and low-latitudes is constrained by measurements and nuclear physics (Nishiizumi et al., 1989; Balco et al., 2008). If sediment that has accumulated cosmogenic nuclides is buried such that production is diminished over $> 10^5$ y, the production ratio decreases because ^{26}Al decays more rapidly than ^{10}Be . Vertical soil mixing intermittently buries sediment grains, suppressing the $^{26}\text{Al}/^{10}\text{Be}$ ratio in sediment shed from the landscape surface (Makhubela et al., 2019).

Paired cosmogenic isotope concentrations are visualized using a two-isotope diagram; the y-axis is the $^{26}\text{Al}/^{10}\text{Be}$ ratio and the x-axis is the concentration of ^{10}Be with normalization based on the production rate of nuclides at the sample site (Klein et al., 1986; Granger, 2006). Sediment samples that have experienced constant exposure with no erosion, or constant exposure under steady-state erosion, will plot within an enclosed region along the top of the diagram; samples that have experienced more complex exposure histories, including burial during or after cosmic-ray exposure, will plot below this

130 region. Such complex histories could include development of a vertically mixed surface layer (Bierman, 1999; Lal and Chen, 2005) as well as extended burial during transport down slopes and in and along rivers.

Using measurements of cosmogenic nuclides to determine basin-averaged denudation rates requires the assumptions that mass loss from the basin is in steady-state, that the mineral used for cosmogenic nuclide measurements is uniformly distributed throughout the watershed, and that
135 denudation occurs within the penetration depth of most cosmic rays, the upper several meters of Earth's surface (Bierman and Steig, 1996). The grain size fraction selected for cosmogenic nuclide analysis must also be representative of the grain size distribution of sediment being produced on slopes (Lukens et al., 2016) although in many landscapes cosmogenic nuclide concentrations do not vary by sediment grainsize.

140 Erosion rates calculated from cosmogenic nuclides can be inaccurate if these assumptions are violated. Rock dissolution can leave sediment enriched in resistant mineral phases, such as zircon, titanite, and quartz—the mineral in which ^{26}Al and ^{10}Be are most commonly measured (Riebe and Granger, 2013). Such enrichment produces underestimates of long-term denudation rates unless
145 accounted for, because the enriched mineral will have a longer residence time relative to the surrounding regolith (Riebe et al., 2001a; Ferrier and Kirchner, 2008). Calculations of denudation rates from cosmogenic nuclide concentrations also rely on the assumption that mass loss is occurring primarily through surface lowering; however, some rock dissolution and, thus, some transfer of mass from rock to groundwater solutions occurs below the depth of most cosmogenic nuclide production (Fig. 1; Small et al., 1999; Dixon et al., 2009a; Riebe and Granger, 2013). In areas with significant rock
150 dissolution at depth, denudation rates inferred from cosmogenic nuclides underestimate denudation because some mass loss occurs below the depth of most nuclide production.

2.2 Chemical weathering corrections to cosmogenically-determined mass loss rates

Although the importance of accounting for loss of mass by chemical weathering (rock dissolution) when calculating cosmogenic erosion rates has been recognized (Small et al., 1999; Riebe
155 et al., 2001a; Dixon et al., 2009a; Riebe and Granger, 2013), few studies incorporate rock dissolution

information or apply correction factors to cosmogenic nuclide-derived rates. In the tropics, some studies compare export rates from dissolved loads in streams to cosmogenically-derived erosion rates, but those studies have considered these two metrics of landscape change separately (Von Blackenburg et al., 2004; Salgado et al., 2006; Hinderer et al., 2013). Other studies use the measurement of insoluble elements in bedrock, saprolite, and soil to quantify quartz enrichment through the weathering process and calculate correction factors that account for the influence of rock dissolution at and near the surface (Small et al., 1999; Riebe et al., 2001a), at depth (Dixon et al., 2009b), or both (Riebe and Granger, 2013).

Of studies that do correct for the influence of chemical weathering when calculating cosmogenic nuclide-derived rates of erosion, Riebe and Granger (2013)'s chemical erosion factor (CEF) method, or earlier quartz enrichment factor method (Riebe et al., 2001a), are often used (Regard et al., 2016). Calculating a CEF requires measurements of soil thickness and density, as well as determining the concentration of the mineral used in cosmogenic nuclide measurements (commonly quartz) and an insoluble element (commonly Zr) in numerous samples of soil, saprolite, and unweathered bedrock. The method is underpinned by the assumption that chemical mass loss is occurring exclusively in well-mixed regolith and deep saprolite (Riebe and Granger, 2013). Erosion rates calculated from cosmogenic nuclide measurements can be multiplied by the CEF to correct for the effects of chemical mass loss (Riebe and Granger, 2013). Chemical erosion factors reported in tropical environments include a CEF of 1.79 in Puerto Rico (Riebe and Granger, 2013) and 3.2 in Cameroon (Regard et al., 2016), demonstrating how significantly cosmogenic nuclide-derived estimates of erosion can underestimate total denudation rates by not accounting for the effects of deep rock dissolution.

3 Study area

Cuba is the largest Caribbean island (~110,000 km²) and is situated along the boundary between the Caribbean and North American plates. Reflecting this complex tectonic setting, Cuban geology is varied and includes silicate, carbonate, and evaporite rocks (Pardo, 2009). Lithologies include marine deposits, accreted volcanic terrains, passive-margin sediments, and obducted ophiolite, all

unconformably overlain by slightly-deformed autochthonous coarse clastic sediment and limestone (Iturralde-Vinent et al., 2016).

The Cuban landscape features a mountainous spine (600-1970 m) descending into low relief coastal plains, except along portions of the south coast where mountains meet the sea. This drainage divide parallels Cuba's east-west orientation, creating rivers that travel relatively short distances from headwaters to base level (Galford et al., 2018). Cuba's climate is tropical wet and dry, with a mean annual temperature of 24.5 °C and average annual precipitation of 1335 mm y⁻¹. The climate is highly seasonal; ~80% of this precipitation is delivered during the wet season from May-October (Llacer, 2012).

Centuries of agriculture have heavily altered the Cuban landscape (Whitbeck, 1922). Prior knowledge of mass loss at the basin scale is limited to measurements of suspended sediment discharge for short periods between 1964 and 1983 for 32 Cuban rivers (Pérez Zorrilla and Ya Karasik, 1989), and measurements of dissolved loads in five limestone basins with karst (Pulina and Fagundo, 1992). In central Cuba, underlying basin rock type is the primary control on surface water geochemistry (Betancourt et al., 2012), a finding supported by geochemical analyses of river waters from the same basins sampled in this study (Bierman et al., 2020). Dissolved load fluxes carried by Cuban rivers (Bierman et al., 2020), and rock dissolution rates inferred from these fluxes, are consistent with rates reported for other Caribbean islands [Dominica, Guadeloupe, and Martinique from Rad et al. (2013) and Puerto Rico from White and Blum (1995)], and high compared to global data compiled by Larsen et al. (2014a).

4 Methods

4.1 Field methods

We collected detrital sediment (n = 26) from the beds of active river channels in central Cuba, representing a variety of basin sizes, average slopes, and lithologies (Fig. 2; Supplement T1-2). Channel morphologies varied, but most streams were incised, and many had exposed bedrock (see Bierman et al. (2020) for photos/descriptions of select field sites). At each site we collected samples for water

chemistry analysis and measured channel parameters, including width, depth, and discharge at time of sampling.

210 4.2 Lab methods

We prepared samples for cosmogenic analysis and extracted beryllium and aluminum following the methodology of Corbett et al. (2016). We sieved bulk sediment samples in the lab and used the 250-850 μm grain size fraction for all samples, except for CU-120, which also includes finer material (63-250 μm) due to low quartz content. Sediment samples were chemically etched to purify quartz and
215 remove meteoric ^{10}Be (Kohl and Nishiizumi, 1992). Twenty-four samples yielded sufficient quartz for analysis. We measured quartz yields for all but one sample (CU-120) by recording the mass of sediment before and after dilute acid etching (Supporting Fig. 1).

We extracted ^{26}Al and ^{10}Be at the National Science Foundation/ University of Vermont Community Cosmogenic Facility, using $\sim 5\text{-}43$ g of quartz per sample (mean = 24 g). We added ~ 250
220 μg of Be to each sample using two different in-house made carriers (Supplement T5); the first batch used a low-ratio carrier made from beryl, while subsequent batches used a dilution of low-ratio commercial SPEX carrier. We added Al to samples with insufficient total Al using a commercial SPEX ICP standard in order to reach a total Al mass of ~ 1500 μg (Supplement T6). Samples were processed in
225 batches of 12, each of which included at least one blank, and two batches included one quality control standard each (Corbett et al., 2019).

$^{10}\text{Be}/^9\text{Be}$ and $^{26}\text{Al}/^{27}\text{Al}$ measurements ($n = 26$, including 2 duplicates), were made by Accelerator Mass Spectrometry (AMS) at the Purdue Rare Isotope Measurement Laboratory (PRIME). ^{10}Be ratios were normalized against standard 07KNSTD3110 with an assumed ratio of 2850×10^{-15} (Nishiizumi et al., 2007) and $^{26}\text{Al}/^{27}\text{Al}$ measurements were normalized against standard KNSTD with an
230 assumed ratio of 1818×10^{-15} (Nishiizumi, 2004). Full laboratory replicate sample preparations and measurements of ^{26}Al and ^{10}Be agree to within $< 2\%$ (Supplement T7; $n=2$). We corrected Be measurements by carrier type, since samples were prepared using different carriers; we use the average of two process blanks ($1.91 \pm 1.01 \times 10^{-15}$; 1SD) to correct 10 samples, and the average of four process blanks ($4.02 \pm 1.00 \times 10^{-15}$; 1SD) for the remaining samples (Supplement T3). We corrected Al

235 measurements using the average of 4 process blanks ($4.97 \pm 2.94 \times 10^{-15}$; Supplement T4). We subtracted
blank ratios from sample ratios and propagated uncertainties in quadrature.

4.3 Analytical methods

We extracted drainage basins and then calculated basin slopes and effective elevations (Portenga
and Bierman, 2011) using the ASTER Global Digital Elevation Model (Lp Daac), determined
240 underlying basin rock types from the USGS Caribbean layer (French and Schenk, 2004), and utilized
precipitation data from the WorldClim dataset (Hijmans et al., 2005) to estimate basin-specific mean
annual precipitation (MAP).

We calculated erosion rates using version 3 of the online erosion rate calculator originally
described by Balco et al. (2008) and subsequently updated (wrapper: 3.0, erates: 3.0, muons: 3.1,
245 validate: validate_v2_input.m - 3.0 consts: 2020-08-26) using the effective elevation (Portenga and
Bierman, 2011) calculated for the basin upstream of the sample collection point, a sample thickness of 0
cm, a rock density of 2.6 g cm^{-3} , and assuming no topographic shielding across this low-relief
landscape. We report erosion rates using the Stone/Lal production scaling scheme.

For samples with the highest ^{10}Be concentrations ($n=4$), we also measured the concentration of
250 cosmogenic ^{21}Ne in quartz to further characterize exposure history (Supplement T10). Neon isotope
measurements were made at the Berkeley Geochronology Center on aliquots of the same purified quartz
samples used for $^{26}\text{Al}/^{10}\text{Be}$ analysis. They were done by vacuum degassing and noble gas mass
spectrometry using the method described in Balter-Kennedy et al. (2020) and Balco and Shuster (2009).

We used measurements of dissolved loads in stream water (Bierman et al., 2020) and modelled
255 annual flows from GLOH₂O (Beck et al., 2015; Beck et al., 2017) to calculate rock dissolution rates for
the 25 basins where we were able to collect water samples. To account for the wide range of lithologies
in our upstream watersheds, including some with evaporites, we modified the approach used by
Erlanger et al. (2021) (Supporting Fig. 2). We removed ions deposited as atmospheric inputs based on
published data on dissolved loads in Cuban rainfall (Préndez et al., 2014). We then determined
260 evaporite weathering rates by balancing Na with Cl and Ca with SO₄. The remaining Na was used to
determine the silicate contribution of Mg and Ca by using an assumed ratio of Na/Mg of 0.25 and

Na/Ca of 0.35 (Erlanger et al., 2021). Silicate weathering rates were calculated as the total of SiO₂ and HPO₄, assumed to result from silicate weathering. Finally, we balanced the remaining Mg and Ca with bicarbonate to determine carbonate weathering rates.

265 Considering a variety of landscape-scale metrics, we explored the relationship between ¹⁰Be-derived erosion rates and calculated rock dissolution (total and silicate, carbonate, and evaporite) rates using linear correlations and their associated p-values. All reported means of sample populations are arithmetic.

5 Results

270 Quartz sand, isolated from central Cuban river sediment, has high concentrations of cosmogenic nuclides (0.41 to 12.6 x 10⁵ atoms g⁻¹ ¹⁰Be and 0.27 to 5.9 x 10⁶ atoms g⁻¹ ²⁶Al). ²⁶Al/¹⁰Be ratios (Fig. 4, Table 1) vary considerably, ranging from 3.65-8.36 (mean = 5.72±1.14, median = 5.83). Sixteen of 24 samples plot below the window defined by continuous exposure and steady erosion on the two-isotope diagram (Fig. 5). Because these ²⁶Al/¹⁰Be data indicate significant burial of quartz during and/or after
275 exposure, many central Cuban drainage basins do not meet the assumption of insignificant nuclide decay inherent in calculations of erosion rates from cosmogenic nuclide concentrations in detrital sediment (Bierman and Steig, 1996). To minimize the impact of violating this assumption, we compare erosion rates based only on the longer-lived nuclide, ¹⁰Be, with landscape scale metrics and dissolved loads. The ¹⁰Be rates, because they cannot properly account for loss of nuclides during burial for
280 samples with depressed ²⁶Al/¹⁰Be ratios, are overestimates of the true rate of erosion.

Erosion rates (Supplement T8), calculated from measured concentrations of ¹⁰Be (Supplement T7), differed considerably between sites. ¹⁰Be-derived erosion rates (Fig. 3) range from 3.4-189 Mg km⁻² y⁻¹ (mean = 59±52, median = 45). Considered as bedrock lowering rates by assuming a bedrock density of 2.6 g cm⁻³, these are 1.3-73 m/My (mean = 23±20, median = 17). ¹⁰Be-derived erosion rates
285 in central Cuba are weakly and positively correlated with mean annual precipitation and slope (Fig. 6). Quartz yields for the samples we analyzed varied widely (0.5%-60%, mean = 20%, median = 17%) but

were not significantly correlated ($p \leq 0.05$) with any basin-scale variables or analytic results (Supplement T1).

290 Rock dissolution rates (Fig. 3) range from 10-176 $\text{Mg km}^{-2} \text{y}^{-1}$ (mean = 92 ± 39 , median = 97) and are higher than ^{10}Be -derived erosion rates in 18 of the 23 basins in which we were able to make both measurements. The median rock dissolution rate is 2.15 times higher than the median ^{10}Be -derived erosion rate. Rock dissolution rates and ^{10}Be -derived erosion rates are not correlated (Fig. 6). However, when total rock dissolution rates are partitioned into silicate, evaporite, and carbonate rates then the silicate dissolution rate is positively correlated with ^{10}Be -determined rates of erosion ($r^2 = 0.18$, $p <$
295 0.05 , Table 2). Rock dissolution rates are not separable by dominant basin lithology. ^{10}Be -inferred erosion rates are metamorphic $>$ sedimentary, igneous $>$ sedimentary (Fig. 4).

There is lithological dependence of ^{10}Be -derived erosion rates and the ratio of rock dissolution to ^{10}Be -derived erosion rates at the basin scale (Fig. 5). ^{10}Be -derived erosion rates of sedimentary rocks were lower than other rock types ($p = 0.01$). Samples with the lowest $^{26}\text{Al}/^{10}\text{Be}$ ratios and the highest
300 ^{10}Be concentrations (Fig. 5) were collected in the north-western part of the field area in basins predominately underlain by sedimentary rocks (Fig. 2). For the most part, samples from basins dominantly underlain by igneous and metamorphic rocks plot to the left on the two-isotope diagram and have higher $^{26}\text{Al}/^{10}\text{Be}$ ratios than quartz from basins underlain by sedimentary rocks. Basins draining primarily sedimentary lithologies have the highest ratio of rock dissolution to ^{10}Be -derived erosion
305 rates. Seven basins (CU-106, 119, 120, 121, 122, 131, and 132) stand out from the rest (Fig. 5) and are clustered in the northwestern part of our field area. These basins have much lower than average ^{10}Be -derived erosion rates (3.4-14.5 $\text{Mg km}^{-2} \text{y}^{-1}$), low $^{26}\text{Al}/^{10}\text{Be}$ ratios (3.80-5.87), and rock dissolution rates 1.7- 29 times higher than the ^{10}Be -derived rates of erosion. All but CU-131 are underlain primarily by sedimentary rocks.

310 Neon isotope measurements (Supplement T10) revealed high total neon concentrations with isotope composition indistinguishable from atmosphere, so excess ^{21}Ne was indistinguishable from zero and could not be quantified. Expected cosmogenic ^{21}Ne concentrations in the samples we analyzed, calculated from observed ^{10}Be concentrations and the assumption of steady erosion (3-6 M atoms g^{-1} cosmogenic ^{21}Ne), would comprise less than 2% of the total amount of ^{21}Ne we observed and would not

315 be detectable at typical measurement uncertainties. The neon isotope measurements are not inconsistent
with the ^{26}Al and ^{10}Be data, but provide no additional information.

6 Discussion

In central Cuba, erosion rates inferred from the concentration of ^{10}Be in river sand vary by more
than an order of magnitude. The lowest ^{10}Be -inferred erosion rate ($3.4 \text{ Mg km}^{-2} \text{ y}^{-1}$; 1.3 m/My) is less
320 than those measured in tectonically stable arid landscapes including Namibia and Australia (Bierman
and Caffee, 2001; Codilean et al., 2021). The highest ^{10}Be -inferred rate ($189 \text{ Mg km}^{-2} \text{ y}^{-1}$; 73 m My^{-1})
exceeds those measured in temperate, humid, tectonically stable areas such as the southern Appalachian
Mountains (Portenga et al., 2019; Duxbury et al., 2015; Linari et al.) and is similar to or less than rates
measured on other Caribbean Islands including Puerto Rico and Dominica (Quock et al., 2021; Brocard
325 et al., 2015; Brown et al., 1995b).

Variability in ^{10}Be concentration, and thus inferred rates of erosion, between central Cuban
drainage basins, many within just a few tens of kilometers of each other with similar basin slope,
suggests significant landscape-scale controls on ^{10}Be concentration and thus mass loss. Indeed, we find
that ^{10}Be -determined erosion rates are positively correlated with slope ($R^2=0.20$, $p=0.03$), mean annual
330 precipitation ($R^2=0.22$, $p=0.02$), and rates of silicate dissolution ($R^2=0.17$, $p=0.05$) (Fig. 6, Table 2).
Erosion rates are lowest for basins underlain dominantly by sedimentary rocks and highest for basins
underlain by metamorphic rocks (Fig. 3). However, accurately quantifying rates of denudation (total
mass loss) from central Cuban landscapes is complicated by significant export of mass in solution and
near-surface quartz enrichment. In the sections that follow, we discuss the ^{10}Be data in the context of
335 dissolved load export in river water and the landscape-scale insight on active processes provided by
dual-isotope measurements (^{26}Al and ^{10}Be) made in riverine quartz.

6.1 Cosmogenic erosion rates underestimate landscape scale mass loss in Cuba

Our data show clearly that significant, landscape-scale mass loss is occurring by solution in
central Cuba. Rock dissolution rates exceed, some by more than an order of magnitude, corresponding

340 ^{10}Be -derived mass loss rates in central Cuba, demonstrating that the cosmogenic nuclide measurements are an incomplete assessment of total mass loss from the landscape. Rock dissolution rates are greater than cosmogenic erosion rates for 18 of the 23 basins we analyzed and the median rock dissolution rate in Cuba is 2.15 times higher than median cosmogenic nuclide-derived rate (Table 1, Fig. 7). Rock dissolution rates that significantly exceed corresponding ^{10}Be -inferred rates have also been reported
345 from Uganda (Hinderer et al., 2013) and Cameroon (Regard et al., 2016), where they were attributed to the influence of easily weathered volcanic tephra and deep weathering associated with thick regolith, respectively. Most other studies that compare rock dissolution rates and ^{10}Be -derived erosion rates in the tropics documented rock dissolution rates within the range of cosmogenic nuclide-derived rates (Von Blackenburg et al., 2004; Salgado et al., 2006; Cherem et al., 2012; Sosa Gonzalez et al., 2016b; Quock et al., 2021). Cuba is different.

The discordance between high rock dissolution rates and low ^{10}Be -derived erosion rates in central Cuba suggests that significant rock weathering is occurring below the depth of most cosmogenic nuclide production (Bierman and Steig, 1996; Fig. 1). The discordance, along with high rates of carbonate and evaporite dissolution in some basins, suggests that many lithologies in our field area are
355 highly susceptible to dissolution. Bierman et al. (2020) attribute high rock dissolution rates and the relationship between stream water chemistry and bedrock type in central Cuba to extensive rock-groundwater interaction along subsurface flow paths, controlled by ongoing bedrock uplift and associated rock fracturing. The prevalence of rock dissolution at depth in Cuba is consistent with findings from other humid, tropical landscapes, including Puerto Rico (White et al., 1998; Kurtz et al.,
360 2011; Chapela Lara et al., 2017; Moore et al., 2019), Guadeloupe, Martinique, Dominica (Rad et al., 2007), and Hawaii (Schopka and Derry, 2012).

We observed no correlation between ^{10}Be -derived erosion and rock dissolution rates in central Cuba (Fig. 7), in contrast to other studies in the tropics that have observed generally positive correlations (Salgado et al., 2006; Cherem et al., 2012; Sosa Gonzalez et al., 2016b) The lack of
365 correlation suggests that mass loss below several meters, the depth at which most cosmogenic nuclides are produced, is an important component of denudation in Cuba. Discordance between high rock dissolution rates and low ^{10}Be -derived erosion rates observed in Cuba occurs in basins with different

underlying lithologies (Fig. 6). Such widespread discordance suggests deep chemical weathering is occurring throughout central Cuba.

370 Carbonate weathering dominates river water geochemistry in central Cuba. Our analysis of Cuban water composition suggests that the rate of carbonate dissolution varies widely and in most basins we sampled, exceeds by several-fold the rate of silicate dissolution (Fig. 4). Silicate dissolution rates are low ($<25 \text{ Mg km}^{-2} \text{ y}^{-1}$) and similar between all lithologies. Export rates of elements calculated to reflect the presence of evaporite minerals are also generally low ($<35 \text{ Mg km}^{-2} \text{ y}^{-1}$), except in four
375 basins dominated by sedimentary rocks (CU-120, 121, 122, 132). Water geochemistry data from four of these basins suggest the presence of significant evaporite deposits due to high concentrations of Cl, SO_4 , Br, and Na (Bierman et al., 2020). Together these data imply that lithologies underlying the basins we sampled are not uniform and that silicate rocks do not account for most of the dissolved mass loss in at least some, and likely many, of the basins we sampled.

380 Together, underlying lithology and topography are important controlling factors in how and how rapidly the Cuban landscapes we studied are losing mass by both physical and chemical weathering. Lowland basins, primarily underlain by sedimentary rocks, on average have low rates of ^{10}Be -inferred mass loss and high rates of dissolution. Six basins underlain by sedimentary lithologies (CU-106, -119, -120, -121, -122, and -132) have the highest ^{10}Be concentrations and lowest erosion rates indicating
385 longer total exposure durations for the quartz we analyzed. All are low slope (0.5 to 1.6 degrees). These six basins also demonstrate the greatest disparity between high rock dissolution rates and low ^{10}Be -derived erosion rates (5.7-29X). One basin underlain by igneous rocks (CU-131) has similarly low slope (0.6 degrees) and high ^{10}Be concentration but a much lower ratio of dissolution to erosion rates (1.7) likely reflecting the paucity of readily soluble minerals. As a result, ^{10}Be -derived erosion rates are
390 weakly and positively correlated with average basin slope ($R^2 = 0.20$, $p=0.03$), but rock dissolution rates are not correlated ($R^2 = 0.04$, $p=0.36$) with slope.

6.2 Low $^{26}\text{Al}/^{10}\text{Be}$ ratio evidence for a deep mixed surface layer and possible quartz enrichment

The $^{26}\text{Al}/^{10}\text{Be}$ ratios suggest that most sediment we collected from central Cuban rivers does not
395 have a simple exposure history. $^{26}\text{Al}/^{10}\text{Be}$ data in 16 of 24 sampled basins are inconsistent with steady
surface erosion (Fig. 5). Many of the basins with the lowest $^{26}\text{Al}/^{10}\text{Be}$ ratios drain predominantly marine
sedimentary lithologies, and have low average basin slopes ($0.5\text{-}0.7^\circ$); the remaining basin drains
primarily igneous rocks and has an average basin slope of 0.6° . These are the same seven basins
discussed in the section above, all but one of which have high ratios of dissolved load to ^{10}Be -inferred
400 erosion rates. There is a significant ($R^2 = 0.34$, $p = 0.003$) relationship between average basin slope and
 $^{26}\text{Al}/^{10}\text{Be}$.

Observed $^{26}\text{Al}/^{10}\text{Be}$ ratios in most of the low-ratio samples are consistent with bioturbation and
prolonged near-surface exposure (Struck et al., 2018). We suspect that at least some of the
inconsistency between measured $^{26}\text{Al}/^{10}\text{Be}$ ratios and those predicted by a simple, steady-state surface
405 erosion model is due to (deep) soil mixing. Typically, the lower boundary of the simple exposure region
of a two-isotope diagram (Fig. 5) is constructed based on the assumption that all grains move
monotonically towards the surface at the rate that the surface is eroding (Granger, 2006). Vertical
mixing, due to bioturbation or other soil processes taking place in the upper layers of soil, violates this
assumption. Within a mixed soil layer, grains circulate at a higher velocity than the erosion rate, and
410 therefore experience an average production rate lower than the surface rate and spend time below the
surface where the rate of nuclide decay may exceed the rate of nuclide production. During burial,
 $^{26}\text{Al}/^{10}\text{Be}$ ratios decrease and diverge from those predicted by the steady-state surface erosion model
(Fig. 5).

Rapid chemical mass loss due to the presence of readily soluble evaporite and marine or igneous
415 lithologies in some basins likely enriches the remaining sediment in quartz. The combination of mass
loss by rapid rock dissolution and the retention of weathering residuum favored by subdued topography
in low-relief basins allows less-soluble material (e.g., quartz) to accumulate at and near the surface,
creating thick regolith. Extensive vertical mixing of near-surface soil, as is expected for flat, forested
landscapes where the rate of bioturbation is likely high in relation to slow erosion rates, leads to longer

420 residence times for these residual mineral grains, and therefore a lower $^{26}\text{Al}/^{10}\text{Be}$ ratio, in a mixed surface layer compared to a surface eroding at the same rate without vertical mixing.

This assertion is supported by the consistency between measured $^{26}\text{Al}/^{10}\text{Be}$ ratios and expected nuclide concentrations and ratios calculated assuming the presence of a mixed surface layer (per Lal and Chen (2005), equation 12). Expected $^{26}\text{Al}/^{10}\text{Be}$ ratios calculated assuming a mixed layer depth of 40-
425 160 cm agree well with measured low $^{26}\text{Al}/^{10}\text{Be}$ ratios from basins CU-119, CU-122 and 132 (Fig. 5). This mixed layer depth range is consistent with the soil depths of 90-150 cm reported for the location of these basins (Bennett and Allison, 1928). In deeply weathered tropical soils, bioturbation can extend to depths of several meters (Von Blackenburg et al., 2004) so it is plausible that mixing depths are even greater than the model suggests, providing an explanation for the $^{26}\text{Al}/^{10}\text{Be}$ ratios measured in CU-120,
430 121, 131, and 106. We were not able to measure regolith depths in the drainage basins we sampled.

The $^{26}\text{Al}/^{10}\text{Be}$ ratio in other samples (e.g., CU-106, 118, and 110) is too low to be attributed solely to the effects of a deep mixed surface layer, and requires that some fraction of the sample has experienced both surface exposure and a significant period of burial well below the surface where cosmogenic nuclide production is negligible. Factors that could lead to this low ratio include the
435 incorporation of previously deeply buried sediment through channel avulsion (Wittmann et al., 2011) or incision into terraces (Hu et al., 2011). We conclude terrace storage, along with a combination of quartz enrichment due to high chemical weathering rates of soluble marine rocks in combination with very low slope basins and a deep mixing layer, combine to generate detrital quartz with high concentrations of ^{10}Be and lower than expected $^{26}\text{Al}/^{10}\text{Be}$ ratios.

440

6.3 Constraining total rates of landscape denudation

The disagreement between high rock dissolution rates and low ^{10}Be -derived erosion rates raises questions about how to best characterize total landscape denudation rates. It is clear from our data set that neither cosmogenic nuclide measurements nor stream solute flux are capturing all or even, in some
445 cases, the majority of landscape denudation in central Cuba. Evidence for deep rock dissolution presented in section 6.1 suggests that sediments and solutes are being sourced at least partially from

different depths in the landscape. Because most mass loss in much of central Cuba occurs in solution (rock dissolution rates are higher than ^{10}Be -derived erosion rates in 18 of 23 basins), rock dissolution rates typically represent a minimum bound on total landscape denudation.

450 Treating the removal of mass in solution and through physical erosion as entirely discrete processes happening at different depths in the landscape sets an upper limit on total landscape denudation: the sum of inferred rock dissolution rates and ^{10}Be -derived erosion rates. Summing ^{10}Be -derived erosion rates and chemical denudation rates increases estimates of total landscape denudation across study basins by a factor of 1.4-30 (mean = 6.3, median = 2.7) above ^{10}Be -derived erosion rates. 455 Disregarding the six basins with evidence of evaporite deposits (CU-106, -119, -120, -121, -122, and -132) leads to an average increase of a factor of 2.6 (median = 2.5) above ^{10}Be -derived erosion rates. These mean and median values for the basins without evaporites are between the reported CEF of 1.79 for the Luquillo Critical Zone Observatory in humid, tropical Puerto Rico (Riebe and Granger, 2013) and the CEF of 3.2 for the thickly saprolite-mantled, tropical environment of south Cameroon (Regard 460 et al., 2016). These comparisons suggest that for landscapes with a significant proportion of total denudation occurring through deep rock dissolution, summing rock dissolution rates and cosmogenic nuclide-derived rates provides a reasonable estimate of total landscape denudation.

In landscapes like central Cuba, total denudation rates may be difficult to predict based on landscape metrics. Summed chemical denudation rates and cosmogenic nuclide-derived erosion rates 465 are not correlated with rock type, as rock type appears to have opposing influences on these rates (i.e., basins underlain by sedimentary rocks had the highest rock dissolution rates but lowest cosmogenic nuclide-derived rates, Fig. 3). Summed rock dissolution rates and cosmogenic nuclide-derived rates do increase with mean basin slope ($R^2=0.19$, $p=0.04$) and mean basin elevation ($R^2=0.17$, $p=0.05$) (Fig. 5), but those relationships are confounded because ^{10}Be -derived rates are highest in high elevation, steep 470 basins and rock dissolution rates are highest in low slope, low elevation basins—relationships that are primarily controlled by the influence of rock type on these two different mass loss processes.

In central Cuba, the lack of correlation between rock dissolution rates and ^{10}Be -derived erosion rates (Fig. 7) suggests a possible mechanism for limiting total reduction in landscape relief. While global data demonstrates significant, positive correlations between ^{10}Be -derived erosion rates and basin

475 slope and relief (Portenga and Bierman, 2011), accounting for the influence of rock dissolution may
alter this dynamic. The possibility of combined physical and chemical processes limiting reductions in
relief has significant implications for the study of deeply weathered, high-relief tropical landscapes. The
dual importance of rock dissolution in low-lying areas and physical erosion in steeper terrain could
explain the relationship behind sustained high relief topography and low ^{10}Be -derived erosion rates
480 common across some tropical landscapes, such as Brazil (Vasconcelos et al., 2019) or Sri Lanka (Von
Blackenburg et al., 2004). Landscapes with high rock dissolution rates and low physical erosion rates
appear to be relatively common (Larsen et al., 2014b). As lowlands weather primarily through rock
dissolution and high relief areas by physical erosion, total relief would change more slowly than ^{10}Be -
estimated rate differentials would suggest.

485 Regardless of rock type, however, cosmogenic nuclide-derived erosion rates are positively
correlated with MAP ($R^2 = 0.22$, $p = 0.02$). While MAP does not vary widely across our study basins in
central Cuba (956 to 1555 mm y^{-1}), this correlation suggests a climatic control on denudation rates
across this landscape. This finding is contrary to other studies in the humid tropics (Von Blackenburg et
al., 2004), and beyond (Riebe et al., 2001b; Portenga and Bierman, 2011), that have found no
490 correlation between climate variables and cosmogenic nuclide-derived long-term erosion rates but
similar to recent findings in humid, temperate Tasmania (Vanlandingham et al., 2022). Since in Cuba
 ^{10}Be -derived erosion rates are positively correlated with MAP but chemical denudation rates are
uncorrelated with MAP, this trend likely highlights the importance of rainfall in allowing for the
physical export of sediment from a drainage basin that is transport-limited rather than weathering-
495 limited.

Our data clearly demonstrate that cosmogenic nuclide measurements can underestimate total
denudation in landscapes with significant rock dissolution at depth, particularly in the tropics. This
suggests that similar underestimates of total denudation rates produced by relying on measurements of
cosmogenic nuclides may be a factor in other tropical landscapes. While rock dissolution rates in the
500 tropics have been documented as among the highest globally (White and Blum, 1995; Rad et al., 2013;
Larsen et al., 2014a), a global compilation of ^{10}Be -derived erosion rates demonstrated that such
isotopically determined rates of erosion are lower in the tropics than in all other climate zones, apart

from arid regions (Portenga and Bierman, 2011). The contrast between these two depictions of tropical denudation suggests that ^{10}Be -derived erosion rates for tropical areas may be incomplete representations of total mass loss from these landscapes because dissolved loads are incompletely accounted for by measurements of ^{10}Be in river sand. This discrepancy highlights the need for more studies that compare rock dissolution rates and cosmogenic nuclide-derived rates to provide more accurate estimations of total landscape denudation (Vanlandingham et al., 2022).

7 Conclusions

The first cosmogenic nuclide measurements from the island of Cuba provide insight into how mass is lost from landscapes in humid, tropical settings. Solution plays a large role in total mass flux, and significant mineral dissolution occurs along weathering fronts meters below the landscape surface. Rock type exerts the primary control on the pace of denudation, and precipitation influences rates of landscape change. We find evidence for thick mixed surface layers in lowland basins and river water chemistry data suggest that deep rock dissolution dominates denudational processes in low slope basins where weathering products remain near the surface for long periods of time.

These findings highlight the necessity of accounting for mass loss by solution at depths below the penetration of most cosmic rays when interpreting cosmogenic nuclide-derived rates in landscapes with the potential for significant rock dissolution. The discrepancy between high rock dissolution rates and low ^{10}Be -derived erosion rates observed in central Cuba emphasizes how relying on cosmogenic nuclide measurements alone to determine total rates of mass loss from landscapes can lead to considerable underestimates of denudation. Summing mass loss rates in solution with mass loss rates inferred from cosmogenic nuclides provides an upper limit for total mass loss from landscapes when significant rock dissolution occurs below the penetration depth of cosmic ray neutrons. These findings suggest that estimating rock dissolution rates is important when applying cosmogenic nuclides in landscapes, especially those which are humid, tropical, have soluble rocks, and/or have deep weathered regolith.

8 Data availability

The data are under review at Pangaea (<https://doi.pangaea.de/10.1594/PANGAEA.940043>).

530 9 Author contributions


AHS, MKC, and PB contributed to project design. AGA, AGM, AHS, HCA, MKC, PBR, and RSH conducted fieldwork. MKC prepared samples for laboratory analysis. GB, LS, MC, AHS, PRB, and MKC contributed to cosmogenic nuclide analysis. MKC, PRB, and AHS led manuscript preparation; all authors assisted with data analysis and manuscript writing and review. GB, MKC, and AHS prepared
535 the figures. AHS completed the chemical weathering calculations.

10 Competing interests


Some authors are members of the editorial board of Geochronology. The peer-review process was guided by an independent editor, and the authors have no other competing interests to declare.

11 Acknowledgements


540 Support for fieldwork and analyses provided by NSF EAR-1719249 and NSF EAR-1719240 to Bierman and Schmidt, NSF EAR 1735676 to Bierman, and Oberlin College funding to Schmidt. Researchers from Centro de Estudios Ambientales de Cienfuegos were supported by the MICATIN and ISOAGRI projects. We thank Jay Racela and Marika Massey-Bierman for their assistance with laboratory work, and Marika Massey-Bierman, Monica Dix, Victor Manuel Fonseca Pérez, and
545 Santiago Gil Pérez for assistance with fieldwork. We thank Erica Erlanger and Claire Lukens for constructive reviews. Prepared in part by LLNL under Contract DE-AC52-07NA27344. LLNL-JRNL-824414.



550 Figure 1: Conceptual diagram showing cosmogenic nuclide production and concentration in a column of soil, saprolite, and rock. The dashed line shows decreasing production of cosmogenic nuclides with depth; solid line shows nuclide concentration with depth, and the white arrow represents mass loss by solution below the depth of significant cosmogenic nuclide production.




555 Figure 2: Maps showing underlying basin geology (panel A; French and Schenk, 2004), elevations (panel B; Lp Daac), and location of study area within the island of Cuba (panel C). Legend for panel A includes the category of the rock units in terms of sedimentary (S), igneous (I), and metamorphic (M). Note that the two marine units in A are separated because they have different chemical load signatures.




560 Figure 3: Maps showing rates of landscape change and isotopic data for each study watershed. A,B. Rock dissolution rates. C,D. ^{10}Be -derived erosion rates. In both maps, darker colors in the basins indicate faster rates of landscape change. E,F. Ratio between rock dissolution rates and ^{10}Be -derived erosion rates; darker colors indicate larger ratios. G,H. $^{26}\text{Al}/^{10}\text{Be}$ ratio. Darker colors are higher ratios. Box plots show the maximum and minimum values in the lines extending from the box; the upper side of the box represents the upper quartile, the line inside of the box represents the median value, and the bottom of the box represents the lower quartile. Y axis units are the same as shown in the corresponding map legend.

565



570 Figure 4: Dissolution rates partitioned by lithology. A,B. Silicate. C,D. Carbonate. E,F. Evaporite. Darker colors represent higher rates. Box plots show the maximum and minimum values in the lines extending from the box; the upper side of the box represents the upper quartile, the line inside of the box represents the median value, and the bottom of the box represents the lower quartile.



575

Figure 5: $^{26}\text{Al}/^{10}\text{Be}$ two-isotope plots. To permit comparison of data from different locations and elevations on the same plot, nuclide concentrations have been normalized by dividing measured concentrations by calculated mean production rates in the respective drainage basins, using production rate calculations from version 3 of the online exposure age calculator described by Balco et al. (2008) and subsequently updated. In both plots, uncertainty ellipses denote 68% confidence regions for the normalized nuclide concentrations, and the black lines are the boundaries of the simple exposure region (Lal, 1991) calculated using the conventional assumption of steady block erosion without vertical mixing. Panel A shows that $^{26}\text{Al}/^{10}\text{Be}$ ratios from basins with high nuclide concentrations, implying low erosion rates, are systematically lower than predicted by steady erosion without vertical mixing. Panel B shows that this inconsistency can, at least in part, be explained by the presence of a mixed layer of at least 160 cm. Circles show expected steady-state nuclide concentrations in a fully mixed surface layer calculated according to Lal and Chen (2005) for a range of erosion rates and mixed-layer thicknesses, which highlights that sediment derived from a deep mixed layer has lower nuclide concentrations and lower $^{26}\text{Al}/^{10}\text{Be}$ ratios than would be expected if the mixed layer were absent. Sample ID as in panel A.

590





Figure 6: Relationship between measured ^{10}Be -derived erosion rates, chemical denudation rates, and the sum of chemical denudation rates and ^{10}Be -derived erosion rates to basin characteristics. Different shaped/colored points represent the dominant underlying rock type in that basin. Plots with $p > 0.05$ are shown with a gray background. Small numbers in upper right are R^2 values; * indicates $p \leq 0.05$, ** $p < 0.01$.

595



600 Figure 7: Scatterplot of rock dissolution rates vs. ¹⁰Be-derived erosion rates. A. Data shown for calculations of carbonate, silicate, and evaporite dissolution rates. B. Data shown for dominant lithologies underlying each sampled basin. Horizontal lines extending from the points demonstrate the uncertainty associated with the calculation of ¹⁰Be-derived erosion rates. Histograms on axes show distribution of data. Dashed line is 1:1.

605

TABLE 2A. Correlation Coefficients for Linear Regressions

10Be-derived erosion	Ratio of rock dissolution to 10Be-derived erosion	²⁶ Al/ ¹⁰ Be ratio	Quartz yield	Carbonate dissolution rate	Silicate dissolution rate	Evaporite dissolution rate	Sum of rock dissolution and 10Be-derived erosion rate	Mean basin slope	Total basin area	Mean annual precipitation	% Agricultural land	Mean basin elevation	
0.12	0.36	0.25	0.32	0.59	0.32	0.65	0.65	0.19	-0.05	0.07	-0.27	0.22	Rock dissolution rate
	-0.55	0.64	-0.01	0.41	0.42	-0.32	0.83	0.44	0.18	0.47	-0.34	0.34	10Be-derived erosion
		-0.46	0.45	-0.43	-0.02	0.85	-0.22	-0.47	-0.20	-0.18	0.16	-0.45	Ratio of rock dissolution to 10Be-derived erosion
			0.17	0.60	0.14	-0.29	0.64	0.58	0.26	0.05	-0.49	0.56	²⁶ Al/ ¹⁰ Be ratio
				0.13	-0.11	0.32	0.17	0.39	-0.19	-0.12	-0.61	0.33	Quartz yield
					0.16	-0.22	0.64	0.68	-0.02	-0.10	-0.54	0.74	Carbonate dissolution rate
						0.04	0.46	-0.13	0.23	0.52	0.24	-0.12	Silicate dissolution rate
							0.11	-0.36	-0.09	0.07	0.10	-0.38	Evaporite dissolution rate
								0.44	0.09	0.33	-0.40	0.41	Sum of rock dissolution and 10Be-derived erosion rate
									-0.11	-0.14	-0.79	0.95	Mean basin slope
										0.23	0.02	-0.05	Total basin area
											0.09	-0.27	Mean annual precipitation
												-0.74	% Agricultural land

TABLE 2B. P-values for Linear Regressions													
10Be-derived erosion	Ratio of rock dissolution to 10Be-derived erosion	26Al/10Be ratio	Quartz yield	Carbonate dissolution rate	Silicate dissolution rate	Evaporite dissolution rate	Sum of rock dissolution and 10Be-derived erosion rate	Mean basin slope	Total basin area	Mean annual precipitation	% Agricultural land	Mean basin elevation	
0.583	0.090	0.260	0.147	0.002	0.124	0.000	0.001	0.363	0.816	0.753	0.184	0.296	Rock dissolution rate
	0.006	0.001	0.977	0.054	0.047	0.143	0.000	0.029	0.397	0.020	0.113	0.105	10Be-derived erosion
		0.028	0.034	0.039	0.941	0.000	0.315	0.024	0.369	0.421	0.467	0.031	Ratio of rock dissolution to 10Be-derived erosion
			0.443	0.002	0.530	0.184	0.001	0.003	0.222	0.818	0.017	0.004	26Al/10Be ratio
				0.563	0.629	0.141	0.442	0.064	0.393	0.581	0.002	0.118	Quartz yield
					0.456	0.291	0.001	0.000	0.941	0.626	0.006	0.000	Carbonate dissolution rate
						0.848	0.025	0.540	0.279	0.007	0.248	0.571	Silicate dissolution rate
							0.605	0.076	0.653	0.754	0.636	0.062	Evaporite dissolution rate
								0.037	0.683	0.120	0.056	0.051	Sum of rock dissolution and 10Be-derived erosion rate
									0.596	0.501	0.000	0.000	Mean basin slope
	p<0.01									0.265	0.936	0.817	Total basin area
											0.679	0.175	Mean annual precipitation
												0.000	% Agricultural land

References

Reference List

- 615 Balco, G. and Shuster, D. L.: Production rate of cosmogenic ^{21}Ne in quartz estimated from ^{10}Be , ^{26}Al , and ^{21}Ne concentrations in slowly eroding Antarctic bedrock surfaces, *Earth Planet Sc Lett*, 281, 48-58, 10.1016/j.epsl.2009.02.006, 2009.
- Balco, G., Stone, J. O., Lifton, N. A., and Dunai, T. J.: A complete and easily accessible means of calculating surface exposure ages or erosion rates from ^{10}Be and ^{26}Al measurements, *Quaternary Geochronology*, 3, 174-195, 10.1016/j.quageo.2007.12.001, 2008.
- 620 Balter-Kennedy, A., Bromley, G., Balco, G., Thomas, H., and Jackson, M. S.: A 14.5-million-year record of East Antarctic Ice Sheet fluctuations from the central Transantarctic Mountains, constrained with cosmogenic ^3He , ^{10}Be , ^{21}Ne , and ^{26}Al , *The Cryosphere*, 14, 2647-2672, 10.5194/tc-14-2647-2020, 2020.
- Barreto, H. N., Varajão, C. A. C., Braucher, R., Bourlès, D. L., Salgado, A. A. R., and Varajão, A. F. D. C.: Denudation rates of the Southern Espinhaço Range, Minas Gerais, Brazil, determined by in situ-produced cosmogenic beryllium-10, *Geomorphology*, 191, 1-13, 10.1016/j.geomorph.2013.01.021, 2013.
- 625 Beck, H. E., de Roo, A., and van Dijk, A. I. J. M.: Global Maps of Streamflow Characteristics Based on Observations from Several Thousand Catchments*, *J Hydrometeorol*, 16, 1478-1501, 10.1175/jhm-d-14-0155.1, 2015.
- Beck, H. E., Vergopolan, N., Pan, M., Levizzani, V., van Dijk, A. I. J. M., Weedon, G. P., Brocca, L., Pappenberger, F., Huffman, G. J., and Wood, E. F.: Global-scale evaluation of 22 precipitation datasets using gauge observations and hydrological modeling, *Hydrol Earth Syst Sc*, 21, 6201-6217, 10.5194/hess-21-6201-2017, 2017.
- 630 Bennett, H. H. and Allison, R. V.: *The soils of Cuba*, Tropical Plant Research Foundation, Washington, D.C.1928.
- Betancourt, C., Suárez, R., and Jorge, F.: Influencia de los procesos naturales y antrópicos sobre la calidad del agua en cuatro embalses cubanos, *Limnetica*, 31, 193-204, 2012.
- Bierman, P. and Steig, E.: Estimating rates of denudation using cosmogenic isotope abundances in sediment, *Earth Surf Proc Land*, 21, 103-203, 1996.
- 635 Bierman, P., Herdandez, R. S., Schmidt, A., Aguila, H. C., Alvarez, Y., Arruebarrena, A., Campbell, M. K., Dethier, D., Dix, M., Massey-Bierman, M., Moya, A. G., Perdrial, J., Racela, J., and Alonso-Hernandez, C.: ¡Cuba! River Water Chemistry Reveals Rapid Chemical Weathering, the Echo of Uplift, and the Promise of More Sustainable Agriculture, *GSA Today*, 30, 4-10, 2020.
- Bierman, P. R. and Caffee, M.: Slow Rates of Rock Surface Erosion and Sediment Production across the Namib Desert and Escarpment, Southern Africa, *Am J Sci*, 301, 326-358, 10.2475/ajs.301.4-5.326, 2001.
- 640 Bierman, P. R., Marsella, K. A., Patterson, C., Davis, P. T., & Caffee, M.: Mid-Pleistocene cosmogenic minimum-age limits for pre-Wisconsinan glacial surfaces in southwestern Minnesota and southern Baffin Island: a multiple nuclide approach, *Geomorphology*, 27, 27-39, 1999.
- Brocard, G. Y., Willenbring, J. K., Scatena, F. N., and Johnson, A. H.: Effects of a tectonically-triggered wave of incision on riverine exports and soil mineralogy in the Luquillo Mountains of Puerto Rico, *Appl Geochem*, 63, 586-598, 2015.
- 645 Brown, E. T., Stallard, R. F., Larsen, M. C., Raisbeck, G. M., and Yiou, F.: Denudation rates determined from the accumulation of in situ-produced ^{10}Be in the Luquillo Experimental Forest, Puerto Rico, *Earth Planet Sc Lett*, 129, 193-202, 1995a.
- Brown, E. T., Stallard, R. F., Larsen, M. C., Raisbeck, G. M., and Yiou, F.: Denudation rates determined from the accumulation of in situ-produced ^{10}Be in the luquillo experimental forest, Puerto Rico, *Earth Planet Sc Lett*, 129, 193-202, [https://doi.org/10.1016/0012-821X\(94\)00249-X](https://doi.org/10.1016/0012-821X(94)00249-X), 1995b.
- 650 Chapela Lara, M., Buss, H. L., Pogge von Strandmann, P. A. E., Schuessler, J. A., and Moore, O. W.: The influence of critical zone processes on the Mg isotope budget in a tropical, highly weathered andesitic catchment, *Geochim Cosmochim Ac*, 202, 77-100, 10.1016/j.gca.2016.12.032, 2017.
- Cherem, L. F. S., Varajão, C. A. C., Braucher, R., Bourlès, D., Salgado, A. A. R., and Varajão, A. C.: Long-term evolution of denudational escarpments in southeastern Brazil, *Geomorphology*, 173-174, 118-127, 10.1016/j.geomorph.2012.06.002, 2012.
- 655 Codilean, A. T., Munack, H., Cohen, T. J., Saktura, W. M., Gray, A., and Mudd, S. M.: OCTOPUS: An Open Cosmogenic Isotope and Luminescence Database, *Earth System Science Data*, 10.5194/essd-2018-32, 2018.
- Codilean, A. T., Fülöp, R. H., Munack, H., Wilcken, K. M., Cohen, T. J., Rood, D. H., Fink, D., Bartley, R., Croke, J., and Fifield, L. K.: Controls on denudation along the East Australian continental margin, *Earth-Science Reviews*, 214, 103543, <https://doi.org/10.1016/j.earscirev.2021.103543>, 2021.
- 660 Corbett, L. B., Bierman, P. R., and Rood, D. H.: An approach for optimizing in situ cosmogenic ^{10}Be sample preparation, *Quaternary Geochronology*, 33, 24-34, 10.1016/j.quageo.2016.02.001, 2016.

- 665 Corbett, L. B., Bierman, P. R., Woodruff, T. E., and Caffee, M. W.: A homogeneous liquid reference material for monitoring the quality and reproducibility of in situ cosmogenic ^{10}Be and ^{26}Al analyses, *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms*, 456, 180-185, 10.1016/j.nimb.2019.05.051, 2019.
- Derrieux, F., Siame, L. L., Bourlès, D. L., Chen, R.-F., Braucher, R., Léanni, L., Lee, J.-C., Chu, H.-T., and Byrne, T. B.: How fast is the denudation of the Taiwan mountain belt? Perspectives from in situ cosmogenic ^{10}Be , *J Asian Earth Sci*, 88, 230-245, 10.1016/j.jseaes.2014.03.012, 2014.
- 670 Dixon, J. L., Heimsath, A. M., and Amundson, R.: The critical role of climate and saprolite weathering in landscape evolution, *Earth Surface Processes and Landforms*, 34, 1507-1521, 10.1002/esp.1836, 2009a.
- Dixon, J. L., Heimsath, A. M., Kaste, J., and Amundson, R.: Climate-driven processes of hillslope weathering, *Geology*, 37, 975-978, 10.1130/g30045a.1, 2009b.
- Duxbury, J., Bierman, P. R., Portenga, E. W., Pavich, M. J., Southworth, S., and Freeman, S. P.: Erosion rates in and around Shenandoah National Park, Virginia, determined using analysis of cosmogenic ^{10}Be , *Am J Sci*, 315, 46-76, 2015.
- 675 Erlanger, E., Rugenstein, J., Bufer, A., Picotti, V., and Willett, S.: Controls on Physical and Chemical Denudation in a Mixed Carbonate-Siliciclastic Orogen, *Journal of Geophysical Research: Earth Surface*, 2021.
- Ferrier, K. L. and Kirchner, J. W.: Effects of physical erosion on chemical denudation rates: A numerical modeling study of soil-mantled hillslopes, *Earth Planet Sc Lett*, 272, 591-599, 10.1016/j.epsl.2008.05.024, 2008.
- 680 French, C. D. and Schenk, C. J.: Map showing geology, oil and gas fields, and geologic provinces of the Caribbean Region: U.S. Geological Survey Open-File Report 97-470-K, 2004.
- Galford, G. L., Fernandez, M., Roman, J., Monasterolo, I., Ahamed, S., Fiske, G., González, P., and Kaufman, L.: Cuban land use and conservation, from rainforests to coral reefs, *B Mar Sci*, 10.5343/bms.2017.1026, 2018.
- Granger, D. E.: A review of burial dating methods using ^{26}Al and ^{10}Be , in: *In-Situ Produced Cosmogenic Nuclides and Quantification of Geological Processes*, edited by: Siame, L. L., Bourlès, D. L., and Brown, E. T., The Geological Society of America, 1-16, 2006.
- 685 Granger, D. E. and Muzikar, P.: Dating sediment burial with in situ-produced cosmogenic nuclides: theory, techniques, and limitations, *Earth Planet Sc Lett*, 188, 269-281, 2001.
- Granger, D. E., Kirchner, J. W., and Finkel, R.: Spatially Averaged Long-Term Erosion Rates Measured from In Situ-Produced Cosmogenic Nuclides in Alluvial Sediment, *The Journal of Geology*, 104, 249-257, 1996.
- 690 Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G., and Jarvis, A.: Very high resolution interpolated climate surfaces for global land areas, *International Journal of Climatology*, 25, 1965-1978, 10.1002/joc.1276, 2005.
- Hinderer, M., Pflanz, D., and Schneider, S.: Chemical Denudation Rates in the Humid Tropics of East Africa and Comparison with ^{10}Be -Derived Erosion Rates, *Procedia Earth and Planetary Science*, 7, 360-364, 10.1016/j.proeps.2013.03.047, 2013.
- Hu, X., Kirby, E., Pan, B., Granger, D. E., and Su, H.: Cosmogenic burial ages reveal sediment reservoir dynamics along the Yellow River, China, *Geology*, 39, 839-842, 10.1130/g32030.1, 2011.
- 695 Iturralde-Vinent, M. A., García-Casco, A., Rojas-Agramonte, Y., Proenza, J. A., Murphy, J. B., and Stern, R. J.: The geology of Cuba: A brief overview and synthesis, *GSA Today*, 4-10, 10.1130/gsatg296a.1, 2016.
- Jonell, T. N., Clift, P. D., Hoang, L. V., Hoang, T., Carter, A., Wittmann, H., Böning, P., Pahnke, K., and Rittenour, T.: Controls on erosion patterns and sediment transport in a monsoonal, tectonically quiescent drainage, Song Gianh, central Vietnam, *Basin Res*, 29, 659-683, 10.1111/bre.12199, 2017.
- 700 Klein, J., Giegengack, R., Middleton, R., Sharma, P., Underwood, J., and Weeks, R.: Revealing histories of exposure using in situ produced ^{26}Al and ^{10}Be in Libyan Desert glass, *Radiocarbon*, 28, 547-555, 1986.
- Kohl, C. P. and Nishiizumi, K.: Chemical isolation of quartz for measurement of in-situ -produced cosmogenic nuclides, *Geochim Cosmochim Acta*, 56, 3583-3587, 1992.
- 705 Kurtz, A. C., Lugolobi, F., and Salvucci, G.: Germanium-silicon as a flow path tracer: Application to the Rio Icacos watershed, *Water Resour Res*, 47, 10.1029/2010wr009853, 2011.
- Lal, D.: Cosmic ray labeling of erosion surfaces: in situ nuclide production rates and erosion models, *Earth Planet Sc Lett*, 104, 424-439, 1991.
- Lal, D. and Chen, J.: Cosmic ray labeling of erosion surfaces II: Special cases of exposure histories of boulders, soils and beach terraces, *Earth Planet Sc Lett*, 236, 797-813, 10.1016/j.epsl.2005.05.025, 2005.
- 710 Larsen, I. J., Montgomery, D. R., and Greenberg, H. M.: The contribution of mountains to global denudation, *Geology*, 42, 527-530, 10.1130/g35136.1, 2014a.
- Larsen, I. J., Almond, P. C., Eger, A., Stone, J. O., Montgomery, D. R., and Malcolm, B.: Rapid soil production and weathering in the Southern Alps, New Zealand, *Science*, 343, 637-640, 2014b.
- 715 Linari, C., Bierman, P., Portenga, E., Pavich, M., and Finkel, R.: C. and Freeman, SPHT (2017) Rates of erosion and landscape change along the Blue Ridge escarpment, southern Appalachian Mountains, estimated from in situ cosmogenic ^{10}Be . *Earth Surface Processes and Landforms*, 42 (6), pp. 928-940.(doi: 10.1002/esp.4051),
- Llacer, I. D.: Comportamiento de la precipitación en estaciones meteorológicas seleccionadas de Cuba, Facultad de Geografía, La Universidad de la Habana, La Habana, Cuba, 2012.

- LP DAAC: ASTER GDEM (Volume 2019) [dataset],
- 720 Lukens, C. E., Riebe, C. S., Sklar, L. S., and Shuster, D. L.: Grain size bias in cosmogenic nuclide studies of stream sediment in steep terrain, *Journal of Geophysical Research: Earth Surface*, 121, 978-999, 10.1002/2016jgf003859, 2016.
- Makhubela, T. V., Kramers, J. D., Scherler, D., Wittmann, H., Dirks, P. H. G. M., and Winkler, S. R.: Effects of long soil surface residence times on apparent cosmogenic nuclide denudation rates and burial ages in the Cradle of Humankind, South Africa, *Earth Surf Proc Land*, 44, 2968-2981, 10.1002/esp.4723, 2019.
- 725 Mandal, S. K., Lupker, M., Burg, J.-P., Valla, P. G., Haghypour, N., and Christl, M.: Spatial variability of ^{10}Be -derived erosion rates across the southern Peninsular Indian escarpment: A key to landscape evolution across passive margins, *Earth Planet Sc Lett*, 425, 154-167, 10.1016/j.epsl.2015.05.050, 2015.
- Marshall, J. A., Roering, J. J., Gavin, D. G., and Granger, D. E.: Late Quaternary climatic controls on erosion rates and geomorphic processes in western Oregon, USA, *Geol Soc Am Bull*, 129, 715-731, 10.1130/b31509.1, 2017.
- 730 Modenesi-Gauttieri, M. C., de Toledo, M. C. M., Hiruma, S. T., Taioli, F., and Shimada, H.: Deep weathering and landscape evolution in a tropical plateau, *Catena*, 85, 221-230, 10.1016/j.catena.2011.01.006, 2011.
- Moore, O. W., Buss, H. L., and Dosseto, A.: Incipient chemical weathering at bedrock fracture interfaces in a tropical critical zone system, Puerto Rico, *Geochim Cosmochim Acta*, 252, 61-87, 10.1016/j.gca.2019.02.028, 2019.
- Nearing, M. A., Xie, Y., Liu, B., and Ye, Y.: Natural and anthropogenic rates of soil erosion, *International Soil and Water Conservation Research*, 5, 77-84, 10.1016/j.iswcr.2017.04.001, 2017.
- 735 Nishiizumi, K.: Preparation of ^{26}Al AMS standards, *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms*, 223-224, 388-392, <https://doi.org/10.1016/j.nimb.2004.04.075>, 2004.
- Nishiizumi, K., Imamura, M., Caffee, M. W., Southon, J. R., Finkel, R. C., and McAninch, J.: Absolute calibration of ^{10}Be AMS standards, *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms*, 258, 403-413, 10.1016/j.nimb.2007.01.297, 2007.
- 740 Nishiizumi, K., Winterer, E. L., Kohl, C. P., Klein, J., Middleton, R., Lal, D., and Arnold, J. R.: Cosmic ray production rates of ^{10}Be and ^{26}Al in quartz from glacially polished rocks, *Journal of Geophysical Research*, 94, 10.1029/JB094iB12p17907, 1989.
- Ollier, C. D.: Deep Weathering, Groundwater and Climate, *Geografiska Annaler*, 70, 285- 290, 1988.
- Pardo, G.: *Geology of Cuba*, AAPG Studies in Geology, The American Association of Petroleum Geologists, Tulsa, Oklahoma 2009.
- 745 Pérez Zorrilla, W. and Ya Karasik, G.: El escurrimiento solido y la erosion hidrica actual de Cuba, *Ciencias de la tierra y del espacio*, 15-16, 67-76, 1989.
- Pope, G. A.: Weathering in the tropics, and related extratropical processes, in: *Treatise on Geomorphology*, edited by: Shroder, J., Weathering and Soils Geomorphology, Academic Press, 2013.
- Portenga, E. W. and Bierman, P. R.: Understanding Earth's eroding surface with ^{10}Be , *GSA Today*, 21, 4-10, 10.1130/g111a.1, 2011.
- 750 Portenga, E. W., Bierman, P. R., Trodick Jr, C. D., Greene, S. E., DeJong, B. D., Rood, D. H., and Pavich, M. J.: Erosion rates and sediment flux within the Potomac River basin quantified over millennial timescales using beryllium isotopes, *Bulletin*, 131, 1295-1311, 2019.
- Préndez, M., López, R., and Carrillo, E.: Physical and Chemical Components of Cuba's Rain: Effects on Air Quality, *International Journal of Atmospheric Sciences*, 2014, 1-8, 10.1155/2014/680735, 2014.
- Pulina, M. and Fagundo, J. R.: Tropical karst and chemical denudation of western Cuba, *Geographia Polonica*, 60, 195-216, 1992.
- 755 Quock, M., Schmidt, A. H., Corbett, L. B., Bierman, P. R., Hidy, A. J., and Caffee, M.: Hurricanes alter ^{10}Be concentrations in tropical river sediment but do not change regional erosion rate estimates, *Earth Surf Proc Land*, 2021.
- Rad, S., Allegre, C., and Louvat, P.: Hidden erosion on volcanic islands, *Earth Planet Sc Lett*, 262, 109-124, 10.1016/j.epsl.2007.07.019, 2007.
- Rad, S., Rivé, K., Vittecoq, B., Cerdan, O., and Allègre, C. J.: Chemical weathering and erosion rates in the Lesser Antilles: An overview in Guadeloupe, Martinique and Dominica, *Journal of South American Earth Sciences*, 45, 331-344, 2013.
- 760 Regard, V., Carretier, S., Boeglin, J.-L., Ndam Ngoupayou, J.-R., Dzana, J.-G., Bedimo Bedimo, J.-P., Riotte, J., and Braun, J.-J.: Denudation rates on cratonic landscapes: comparison between suspended and dissolved fluxes, and ^{10}Be analysis in the Nyong and Sanaga River basins, south Cameroon, *Earth Surf Proc Land*, 41, 1671-1683, 10.1002/esp.3939, 2016.
- Reinhardt, L. J., Bishop, P., Hoey, T. B., Dempster, T. J., and Sanderson, D. C. W.: Quantification of the transient response to base-level fall in a small mountain catchment: Sierra Nevada, southern Spain, *Journal of Geophysical Research*, 112, 10.1029/2006jgf000524, 2007.
- 765 Reusser, L., Bierman, P., and Rood, D.: Quantifying human impacts on rates of erosion and sediment transport at a landscape scale, *Geology*, 43, 171-174, 2015.
- Riebe, C. S. and Granger, D. E.: Quantifying effects of deep and near-surface chemical erosion on cosmogenic nuclides in soils, saprolite, and sediment, *Earth Surf Proc Land*, 38, 523-533, 10.1002/esp.3339, 2013.
- 770 Riebe, C. S., Kirchner, J. W., and Finkel, R. C.: Long-term rates of chemical weathering and physical erosion from cosmogenic nuclides and geochemical mass balance, *Geochimica et Cosmochimica Acta*, 67, 4411-4427, 10.1016/s0016-7037(03)00382-x, 2003.
- Riebe, C. S., Kirchner, J. W., and Granger, D. E.: Quantifying quartz enrichment and its consequences for cosmogenic measurements of erosion rates from alluvial sediment and regolith, *Geomorphology*, 15-19, 2001a.

- 775 Riebe, C. S., Kirchner, J. W., Granger, D. E., and Finkel, R. C.: Minimal climatic control on erosion rates in the Sierra Nevada, California, *The Journal of Geology*, 29, 447-450, 2001b.
- Salgado, A. A. R., Braucher, R., Colin, F., Nalini, H. A., Varajão, A. F. D. C., and Varajão, C. A. C.: Denudation rates of the Quadrilátero Ferrífero (Minas Gerais, Brazil): Preliminary results from measurements of solute fluxes in rivers and in situ-produced cosmogenic ^{10}Be , *J Geochem Explor*, 88, 313-317, 10.1016/j.gexplo.2005.08.064, 2006.
- 780 Scherler, D., Bookhagen, B., and Strecker, M. R.: Tectonic control on ^{10}Be -derived erosion rates in the Garhwal Himalaya, India, *Journal of Geophysical Research: Earth Surface*, 119, 83-105, 10.1002/2013jfr002955, 2014.
- Schopka, H. H. and Derry, L. A.: Chemical weathering fluxes from volcanic islands and the importance of groundwater: The Hawaiian example, *Earth Planet Sc Lett*, 339-340, 67-78, 10.1016/j.epsl.2012.05.028, 2012.
- Small, E. E., Anderson, R. S., and Hancock, G. S.: Estimates of the rate of regolith production using ^{10}Be and ^{26}Al from an alpine hillslope, *Geomorphology*, 131-150, 1999.
- 785 Sosa Gonzalez, V., Bierman, P. R., Fernandes, N. F., and Rood, D. H.: Long-term background denudation rates of southern and southeastern Brazilian watersheds estimated with cosmogenic ^{10}Be , *Geomorphology*, 268, 54-63, 10.1016/j.geomorph.2016.05.024, 2016a.
- Sosa Gonzalez, V., Bierman, P. R., Nichols, K. K., and Rood, D. H.: Long-term erosion rates of Panamanian drainage basins determined using in situ ^{10}Be , *Geomorphology*, 275, 1-15, 10.1016/j.geomorph.2016.04.025, 2016b.
- 790 Struck, M., Jansen, J. D., Fujioka, T., Codilean, A. T., Fink, D., Egholm, D. L., Fülöp, R.-H., Wilcken, K. M., and Kotevski, S.: Soil production and transport on postorogenic desert hillslopes quantified with ^{10}Be and ^{26}Al , *GSA Bulletin*, 130, 1017-1040, 10.1130/b31767.1, 2018.
- VanLandingham, L. A., Portenga, E. W., Lefroy, E. C., Schmidt, A. H., Bierman, P. R., and Hidy, A. J.: Comparison of basin-scale in situ and meteoric ^{10}Be erosion and denudation rates in felsic lithologies across an elevation gradient at the George River, northeast Tasmania, Australia, *Geochronology*, 4, 153-176, 2022.
- 795 Vasconcelos, P. M., Farley, K. A., Stone, J., Piacentini, T., and Fifield, L. K.: Stranded landscapes in the humid tropics: Earth's oldest land surfaces, *Earth Planet Sc Lett*, 519, 152-164, 10.1016/j.epsl.2019.04.014, 2019.
- von Blanckenburg, F., Hewawasam, T., and Kubik, P. W.: Cosmogenic nuclide evidence for low weathering and denudation in the wet, tropical highlands of Sri Lanka, *Journal of Geophysical Research*, 109, 1-22, 10.1029/2003JF000049, 2004.
- Whitbeck, R. H.: Geographical Relations in the Development of Cuban Agriculture, *American Geographical Society*, 12, 223-240, 1922.
- 800 White, A., Blum, A., Schulz, M., Vivit, D., Stonestrom, D., Larsen, M., Murphy, S., and Eberl, D.: Chemical weathering in a tropical watershed, Luquillo Mountains, Puerto Rico: I. Long-term versus short-term weathering fluxes, *Geochemica et Cosmochimica Acta*, 62, 209-226, 1998.
- White, A. F. and Blum, A. E.: Effects of climate on chemical weathering in watersheds, *Geochemica et Cosmochimica Acta*, 59, 1729-1747, 1995.
- 805 Wittmann, H., von Blanckenburg, F., Maurice, L., Guyot, J. L., and Kubik, P. W.: Recycling of Amazon floodplain sediment quantified by cosmogenic ^{26}Al and ^{10}Be , *Geology*, 39, 467-470, 10.1130/g31829.1, 2011.

810