



## Improving age-depth correlations by using the LANDO model ensemble

5 Gregor Pfalz<sup>1,2,3,4</sup>, Bernhard Diekmann<sup>1,2</sup>, Johann-Christoph Freytag<sup>3,4</sup>, Liudmila Syrykh<sup>5</sup>,  
Dmitry A. Subetto<sup>5,6</sup>, Boris K. Biskaborn<sup>1,2</sup>

<sup>1</sup>Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Research Unit Potsdam, Telegrafenberg A45, 14473 Potsdam, Germany

<sup>2</sup>University of Potsdam, Institute of Geosciences, Karl-Liebknecht-Str. 24-25, 14476 Potsdam-Golm, Germany

<sup>3</sup>Einstein Center Digital Future, Robert-Koch-Forum, Wilhelmstraße 67, 10117 Berlin, Germany

10 <sup>4</sup>Humboldt-Universität zu Berlin, Department of Computer Science, Unter den Linden 6, 10099 Berlin, Germany

<sup>5</sup>Herzen State Pedagogical University of Russia, Moyka Emb. 48, St. Petersburg 191186, Russia

<sup>6</sup>Institute for Water and Environmental Problems of the Siberian Branch of the Russian Academy of Sciences, Molodezhnayastr.1, Barnaul 656038, Russia

*Correspondence to:* Gregor Pfalz (Gregor.Pfalz@awi.de), Boris K. Biskaborn (Boris.Biskaborn@awi.de)

### 15 **Abstract.**

Age-depth correlations are the key elements in paleoenvironmental studies to place proxy measurements into a temporal context. However, potential influencing factors of the available radiocarbon data and the associated modeling process can cause serious divergences of age-depth correlations from true chronologies, which is particularly challenging for paleolimnological studies in Arctic regions. This paper provides geoscientists with a tool-assisted approach to compare outputs from age-depth modeling systems and to strengthen the robustness of age-depth correlations. We primarily focused in the development on age determination data from a data collection of high latitude lake systems (50° N to 90° N, 62 sediment cores, and a total of 661 dating points). Our approach used five age-depth modeling systems (*Bacon*, *Bchron*, *clam*, *hamstr*, *Undatable*) that we linked through a multi-language Jupyter Notebook called **LANDO** (“**L**inked **a**ge and **d**epth modeling”). Within LANDO we have implemented a pipeline from data integration to model comparison to allow users to investigate the outputs of the modeling systems. In this paper, we focused on highlighting three different case studies: comparing multiple modeling systems for one sediment core with a continuous, undisturbed succession of dating points (CS1 - “*Undisturbed sequence*”), for one sediment core with scattered dating points (CS2 - “*Inconsistent sequence*”), and for multiple sediment cores (CS3 - “*Multiple cores*”). For the first case study (CS1), we showed how we facilitate the output data from all modeling systems to create an ensemble age-depth model. In the special case of scattered dating points (CS2), we introduced an adapted method that uses independent proxy data to assess the performance of each modeling system in representing lithological changes. Based on this evaluation, we reproduced the characteristics of an existing age-depth model (Lake Ilirney, EN18208) without removing age determination data. For the multiple sediment core (CS3) we found that when considering the Pleistocene-Holocene transition, the main regime changes in sedimentation rates do not occur synchronously for all lakes. We linked this behavior to the uncertainty within the modeling process as well as the local variability of the sediment cores within the collection.

### 1 **Introduction**

40 Lakes sediments are important terrestrial archives for recording climate variability within the Northern Hemisphere (Biskaborn et al., 2016; Smol, 2016; Lehnher et al., 2018; Subetto et al., 2017; Syrykh et al., 2021).



The identification of age-depth relationships in those lake sediments helps us to put their measured sediment properties in a temporal context (Bradley, 2015; Lowe and Walker, 2014; Blaauw and Heegaard, 2012). We can determine these relationships by directly counting the annual laminated layers (varves) (Brauer, 2004; Zolitschka et al., 2015), or by using indirect age determination methods such as radiocarbon, optically stimulated luminescence (OSL), or lead-cesium (Lead-210/Cesium-137) dating (Lowe and Walker, 2014; Bradley, 2015; Appleby, 2008; Hajdas et al., 2021). Defining a reliable age-depth correlation for paleoenvironmental studies in cold regions is particularly challenging, as varves only exist in rare cases and the determination of ages mostly depends on radiocarbon dating (Strunk et al., 2020 and references therein). Because of primarily financial restrictions, however, only a few selected samples are taken from sediment core sections to determine the corresponding ages of certain depths (Blaauw et al., 2018; Ciarletta et al., 2019; Olsen et al., 2017). We therefore rely on model calculations to define the ages between the samples. In addition to the mathematical challenges that arise when establishing age-depth correlations, the selection of appropriate dating material has an impact on the modeling process.

In the special case of Arctic lake systems, the amount of material for radiocarbon dating, i.e. aquatic/terrestrial macrofossils and organic remains, is extremely low (Abbott and Stafford, 1996; Colman et al., 1996; Strunk et al., 2020). Radiocarbon dating is therefore often based on the organic carbon content in bulk sediment samples, which can be relatively small due to the lower bioproductivity in those lakes (Strunk et al., 2020 and references therein). However, the use of bulk sediments is problematic, as some portions of contributing carbon are not occurring at the same time as the deposition but may reveal inherited ages from reworked older materials (Rudaya et al., 2016; Biskaborn et al., 2013, 2019). Several methods are available for pre-treating bulk sediment samples to address sample-based dating uncertainties (Brock et al., 2010; Strunk et al., 2020; Rethemeyer et al., 2019; Bao et al., 2019; Dee et al., 2020). Each pre-treatment method may yield a different result for the same material due to the influence of humic acids, fulvic acids, and humins (Brock et al., 2010; Strunk et al., 2020; Abbott and Stafford, 1996). Similarly, older, inert material incorporated by living organism, known as “reservoir effect” or “hard-water effect”, distorts the actual radiocarbon age by up to  $\pm 10\,000$  years (Ascough et al., 2005; Austin et al., 1995; Lougheed et al., 2016). Such a distortion creates methodological and mathematical errors in the development of age-depth correlations, which possibly leads to a misinterpretation of these relationships.

There are numerous geochronological software systems (from now on simply called modeling systems) available to the geoscientific community, which try to solve the challenges stated above (Trachsel and Telford, 2017; Wright et al., 2017; Lacourse and Gajewski, 2020). Implemented methods for detecting outliers, accounting for varying sedimentation rates, or using bootstrapping processes support the construction of an age-depth model (Parnell et al., 2011; Lougheed and Obrochta, 2019; Bronk Ramsey, 2009, 2008).

However, the correct usage of those systems requires a high degree of understanding of the underlying mathematical methods and models. Trachsel and Telford (2017) noted that, despite the users’ impact on the outcome of the model by setting priors and parameters, most users do not have any prior objective insights into appropriately choosing the right parameters. Wright et al. (2017), Trachsel and Telford (2017), and Lacourse and Gajewski (2020) even showed that the results produced by modeling systems could diverge from the true chronology. An in-depth comparison of the results is therefore extremely error-prone. Due to time constraints, usually, users only select and apply one modeling system for paleoenvironmental interpretation.



The objective of this paper is to reduce the effort to apply different methods for determining age-depth correlations and to make their results comparable. We provide a tool to link five selected modeling systems in a single multi-language Jupyter Notebook. We introduce an ensemble age-depth model that uses uninformed models to create data-driven, semi-informed age-depth correlation. We demonstrate the power of our tool by highlighting three case studies in which we examine our application for individual sediment cores and a collection of multiple sediment cores. Throughout this paper, the term “LANDO” refers to our implementation, which stands for “**Linked age and depth modeling**”.

In this paper, we use both published and unpublished age determination data from 62 sediment cores from high latitude lake systems (50° N to 90° N). This unique collection of age determination data allows us to thoroughly test LANDO by examining changes of sedimentation rates over time for various modeling and lake systems. We provide an overview on the acquired metadata in the repository mentioned in the “Code and data availability” section. The harmonization of the acquired data follows the conceptual framework described in Pfalz et al. (2021).

## 2 Methods

A key element in our data-science based approach for developing comparable age-depth correlations was to facilitate the use of modeling systems independent from their original proprietary development environment. A multi-language data analysis environment, such as the SoS notebook (Peng et al., 2018) or GraalVM (Niephaus et al., 2019), provides an interface that enables the comparison of modeling systems without being limited to one programming language or environment. Our implementation used the SoS notebook as its backbone. The SoS notebook is a native Python- and JavaScript-based Jupyter Notebook (Kluyver et al., 2016), which extends to other languages through so-called “Jupyter kernels”. We developed our implementation with the focus on four languages and their respective kernels: Python, R, Octave, and MATLAB. This selection allowed us to use the most common modeling systems.

According to Lacourse and Gajewski (2020), the most commonly used modeling systems are *Bacon* (Blaauw and Christen, 2011), *Bchron* (Haslett and Parnell, 2008; Parnell et al., 2008), *OxCal* (Bronk Ramsey, 1995; Bronk Ramsey and Lee, 2013), and *clam* (Blaauw, 2010). We additionally considered the MATLAB/Octave software *Undatable* (Lougheed and Obrochta, 2019), as an alternative to the classical Bayesian approach, and the R package *hamstr* (Dolman, 2021). The *hamstr* system is an implementation of the *Bacon* algorithm by Blaauw and Christen (2011), which improves the algorithm by additionally adding hierarchical accumulation structures.

In our study, we were able to connect five of the above-mentioned modeling systems in the SoS notebook, namely: *Bacon*, *Bchron*, *clam*, *hamstr*, and *Undatable*. The workflow of LANDO consists of five major components: Input – Preparation – Execution – Result aggregation – Evaluation of model performance.

### 2.1 Input

To work with LANDO users need to provide age determination data, e.g., data from radiocarbon or OSL dating, and associated metadata as listed in [Table 1](#). We developed two import options for the users: through a single spreadsheet or a connection to a database. For this study, we used a connection to a PostgreSQL database, which we developed after the conceptual framework as described in Pfalz et al. (2021), via the Python package “SQLAlchemy” (Bayer, 2012). We divided age determination input data into two attribute categories: necessary and recommended. The category “necessary” focused on the prerequisites of the individual modeling systems as



120

well as project-related attributes, such as unique identifiers, i.e., “measurementid”, “labid”. However, a larger comprehensive set of descriptive metadata helps a better understanding of the data (Cadena-Vela et al., 2020; Thanos, 2017). We added four additional attributes from the category “recommended” to facilitate the interpretation of age-depth models regarding their age determination data.

**Table 1** – *Necessary and recommended attributes for age determination input data, when used with LANDO. Attributes apply for both input methods through either a database or a spreadsheet.*

Attribute	Description	Data type	Necessary/ Recommended
<i>measurementid</i>	Composite key composed of a unique CoreID, a blank space, and the depth below sediment surface (mid-point cm) with max. two decimal digits of corresponding analytical age measurement - <i>example: “CoreA1 100.5”, when users obtained sample of CoreA1 between 100 and 101 cm depth</i>	string	Necessary
<i>thickness</i>	Thickness of the layer from which the sample for age determination was taken from in [cm]	float	Necessary
<i>labid</i>	Unique sample identifier that was provided by the laboratory for age determination	string	Necessary
<i>lab_location</i>	Name of city, where laboratory that conducted the analysis resides	string	Recommended
<i>material_category</i>	One of the eight categories that describes the material best, based on the categories from age-depth modeling system <i>Undatable</i> (Lougheed and Obrochta, 2019) <i>14C marine fossil</i> <i>tiepoint</i> <i>14C terrestrial fossil</i> <i>paleomag</i> <i>14C sediment</i> <i>U/Th</i> <i>tephra</i> <i>other</i>	string	Necessary
<i>material_description</i>	Short description of the used material	string	Recommended
<i>material_weight</i>	Weight of analyzed carbon used in radiocarbon dating in [μgC]	float	Recommended
<i>age</i>	Uncalibrated radiocarbon age in [uncal yr BP], or non-radiocarbon ages as values in [yr BP] (BP = Before Present (before 1950 CE))	float	Necessary
<i>age_error</i>	Error of the uncalibrated radiocarbon age and non-radiocarbon age in [yr]	float	Necessary
<i>pretreatment_dating</i>	Concise description or abbreviation of sample pre-treatment - <i>example: “ABA”, when radiocarbon pre-treatment comprises of an acid-base-acid sequence</i>	string	Recommended
<i>reservoir_age</i>	Additional reservoir effect (also known as hard-water effect or age offset) identified by the user in [yr]; if unknown, then insert 0	float	Necessary
<i>reservoir_error</i>	Error of reservoir age known to the user in [yr]; if unknown, then insert 0	float	Necessary

125

If users decide to use a spreadsheet as input option, then the spreadsheet should follow the same attribution as the database. In addition, we implemented an input prompt for further information, such as the year of core drilling



and core length, to ensure comparability to our database implementation. We provide an example spreadsheet with all attributes in the expected format in the repository mentioned in the “Code and data availability” section of this paper.

## 2.2 Preparation

The preparation component consisted of two separate steps. First, we checked each age determination dataset, whether a reservoir effect was influencing the radiocarbon data. In the absence of a known reservoir age or recent surface sample, we used available radiocarbon data points and a fast-calculating modeling system to predict the age of the upper most layer within a sediment core. In our approach, we used the *hamstr* package with a default value of 6000 iterations. We then compared the predicted value for the upper most layer with the year of the core retrieval, i.e., our target age. We accounted for an uncertainty in the estimate by allowing an extra 10% error between predicted age and target age. If a gap between predicted and target age is observable, then we assumed a reservoir effect is present. We calculated the reservoir effect by subtracting the target age from the mean predicted age, whereas the associated error we based on the two-sigma uncertainty ranges of the prediction. LANDO allow users to add the calculated reservoir age and its uncertainty range to the corresponding attributes (“reservoir\_age” and “reservoir\_error”). Depending on the choice of the user, this addition affects either all radiocarbon samples or only bulk sediment samples, or users completely discard the output for the subsequent modeling process.

As second step in the preparation component, we built a module that automatically changes the format of the available data to the individually desired input of each of the five modeling systems implemented in LANDO. We primarily used the Python package “pandas” (Reback et al., 2020) for the transformation within the module. We transferred the newly transformed age determination data to the corresponding programming language for age-depth modeling using the built-in “%get” function of SoS notebook.

## 2.3 Execution

We developed LANDO with the specific ability of create multiple age-depth models for multiple dating series from spatially distributed lake systems. Hence, reducing overall computing time was one of our highest priorities. We achieved this reduction by applying existing parallelization back-ends for both R and Python, such as “doParallel” (Microsoft Corporation and Weston, 2020a) and “Dask” (Dask Development Team, 2016), respectively. For each modeling system in R, we wrote a separate script that takes advantage of the parallelization back-end “doParallel”. Besides the individual modeling system packages, we made use of different R libraries, such as “tidyverse” (Wickham et al., 2019), “parallel” (R Core Team, 2021), “foreach” (Microsoft Corporation and Weston, 2020c), “doRNG” (Gaujoux, 2020), and “doSNOW” (Microsoft Corporation and Weston, 2020b). We neglected the use of parallelization for the *Undatable* software in MATLAB, since even the sequential execution for several sediment cores in our test setup was on the order of a few minutes. However, we achieved comparable results with *Undatable* in Octave using the parallelization package “parallel” (Fujiwara et al., 2021).

As mentioned before, the selection of model priors and parameters has an impact on the modeling outcome, if no objective prior knowledge exist. To lower our impact and to avoid introducing biases in the modeling process, we used the default values from each modeling system as our own default values (Blaauw et al., 2021; Blaauw, 2021; Parnell et al., 2008; Dolman, 2021; Lougheed and Obrochta, 2019). In our adaptation of clam, the parameter “poly\_degree” controls the polynomial degree of models for type 2, while the parameter “smoothing” controls the



170 degree of smoothing for type 4 and 5. In the original version of clam, users adjust both parameters with the single option “smooth” (Blaauw, 2021). Furthermore, the default value for “ssize” within the original version of Bacon is 2000. We increased this value to 8000 to ensure good MCMC mixing for problematic cores (Blaauw et al., 2021). In case the user has in-depth knowledge about his sediment core and wants to change certain values, we opted for making crucial parameters accessible within the SoS notebook outside of the executing scripts. **Table 2** provides an overview of all values which users can access and change for the individual systems. However, we limited the access to some parameters for operational purposes, such as the number of iterations or the resolution of the output.

**Table 2** – Default values for each modeling system, which users can access and change within LANDO.

Modeling system	Parameter	Default value
<i>Bacon</i>		
	acc.shape	1.5
	acc.mean	20
	mem.strength	10
	mem.mean	0.5
	ssize	8000
<i>Bchron</i>		
	<i>not applicable</i>	-
<i>clam</i>		
	types	1 to 5
	poly_degree	1 to 4
	smoothing	0.1 to 1.0
<i>hamstr</i>		
	K	c(10,10)
<i>Undatable</i>		
	xfactor	0.1
	bootpc	30

175

## 2.4 Result aggregation

180 After every model run, we received 10000 age estimates (also known as “iterations” or “realizations”) per centimeter from each modeling system for every sediment core. We transferred these results back to Python using the built-in “%put” function of SoS notebook, where in the next module, we calculated per centimeter the median and mean age values as well as one-sigma and two-sigma age ranges. For the summarizing statistics, we used standard Python libraries such as “pandas” (Reback et al., 2020) and “numpy” (Harris et al., 2020). We appended the model name as attribute to the statistics to allocate each result to its modeling system. In addition, we implemented a module, which helped us to push the aggregated result to our initial database to reuse in follow-up research projects. In a similar approach to the input component, we established the connection to our designed  
 185 PostgreSQL database via the package “SQLAlchemy” (Bayer, 2012).

Similarly, we used the 10000 age estimates per centimeter for calculating the sedimentation rates. Our calculation used three different approaches to calculate sedimentation rates: “naïve”, “moving average over three depths”, and “moving average over five depths”. **Table 3** lists the appropriate equations for each approach. The user can decide which one of the three approaches best applies to the individual sediment record. We summarized the output into  
 190 the basic summarizing statistics (mean, median, one-sigma ranges, and two sigma ranges) accessible to the users, but added the model name and employed approach as additional attributes. If users use more than one sediment



core for sedimentation rate calculation, then LANDO will automatically execute the sedimentation rate calculation in parallel using the “Dask” back-end (Dask Development Team, 2016) and the “joblib” Python package (Joblib Development Team, 2020).

195 **Table 3** – *Approaches to calculate sedimentation rates within LANDO. The value represents the layer of interest within a sediment core for which the calculation is necessary. Both  $x_{i+1}$  and  $x_{i+2}$  are the following layers, while  $x_{i-1}$  and  $x_{i-2}$  are the previous layers. The unit for the resulting sedimentation rate is centimeter per year [cm/yr].*

Approach	Equation
<i>Naïve (default)</i>	$\text{sedimentation rate } (x_i) = \frac{\text{depth}(x_i) - \text{depth}(x_{i-1})}{\text{age}(x_i) - \text{age}(x_{i-1})}$
<i>Moving average over three depths</i>	$\text{sedimentation rate } (x_i) = \frac{\text{depth}(x_{i+1}) - \text{depth}(x_{i-1})}{\text{age}(x_{i+1}) - \text{age}(x_{i-1})}$
<i>Moving average over five depths</i>	$\text{sedimentation rate } (x_i) = \frac{\text{depth}(x_{i+2}) - \text{depth}(x_{i-2})}{\text{age}(x_{i+2}) - \text{age}(x_{i-2})}$

## 200 2.5 Evaluation of model performance

To evaluate the performance of each modeling system, we looked at three different case studies:

Case Study no. 1 - Comparison of multiple modeling systems for one sediment core with a continuous, undisturbed sequence of dating points (“*Undisturbed sequence*” – CS1)

205 Case Study no. 2 - Comparison of multiple modeling systems for one sediment core with a disturbed sequence (including inversions) of dating points (“*Inconsistent sequence*” – CS2)


Case Study no. 3 - Comparison of sedimentation rate changes for multiple sediment cores (“*Multiple cores*” – CS3)

210 We examined both sedimentation rate and age-depth modeling results in each of the three case studies. For the first case study, we selected the sediment core EN18218 (Vyse et al., 2021) to showcase the generated output of LANDO. The 6.53 m long sediment record obtained from Lake Rauchuvagytygn, Chukotka (67.78938° N, 168.73352° E, core location water depth: 29.5 m) during an expedition in 2018 consisted of 23 bulk sediment samples used for radiocarbon sampling. The authors determined an existing age offset of  $785 \pm 31$  yr BP (years Before Present, i.e., before 1950 CE), which we used in our modeling process as well.

215 As counterexample for the second case study, we have chosen the sediment core EN18208 (Vyse et al., 2020). During the same expedition to Russia’s Far East in 2018, scientists recovered this EN18208 core from Lake Ilirney, Chukotka (67.34030° N, 168.29567° E, core length: 10.76 m, core location water depth: 19.0 m). The authors based their age-depth model on four OSL dates and 17 radiocarbon dates from bulk sediment samples as well as an age offset of  $1721 \pm 28$  yr BP. However, in addition to the age offset, we included all of the seven available OSL and 25 radiocarbon dates for this core in our study. More details on each sediment cores are accessible in the



220 corresponding references. Both cores are also part of the “Multiple cores” case study with a total of 62 sediment  
 cores (Figure 1).



225 **Figure 1** – Map of geographical distribution of lake sediment cores used for our study (triangles,  $n = 62$ ). The  
 orange triangles ( $n = 39$ ) represent sediment cores for which data of age determination were available either  
 through a corresponding publication or a publicly accessible database, e.g., Pangaia (Diepenbroek et al.,  
 2002). Purple triangles ( $n = 23$ ) indicate unpublished or referenced datasets, such as those provided by the  
 PaleoLake Database (Syrykh et al., 2021). ArcGIS Basemap: GEBCO Grid 2014 modified by AWI. The outer  
 ring in the graphic corresponds to 45° N.

### 2.5.1 Numerical combination of model outputs

230 To introduce the ensemble model in LANDO, we combined the outputs from all five modeling systems into one  
 composite model. We considered the outermost limits (min. and max. values) of all confidence intervals (one-  
 sigma or two-sigma) as our boundary for the ensemble model. By taking these outermost limits into account, we  
 artificially increased the area of uncertainty covered by the ensemble model, but we made sure that we were  
 representing all possible outcomes and maximizing the likelihood of including the true chronology. We also  
 235 included a weighted averages ( $\bar{x}$ ) of the age estimates and sedimentation rates, which we calculated using the  
 following equations:

$$\bar{x} = \sum_{k=1}^m \frac{n_k}{n} * \bar{x}_k \quad (\text{Eq. 1})$$

$$n = \sum_{k=1}^m n_k \quad (\text{Eq. 2})$$



with  $m$  being the number of participating modeling systems,  $n$  as the total number of iterations as well as  $\bar{x}_k$  and  
240  $n_k$  representing the median value (either for age estimate or sedimentation rate) and the associated number of  
iterations from each modeling system, respectively. In some cases, the weights from each modeling system are  
equal, as they produce the same number of iterations. Then we can simplify Eq. 1 to represent the arithmetic mean:

$$\bar{x} = \frac{1}{m} \sum_{k=1}^m \bar{x}_k \quad (\text{Eq. 3})$$

For our “*Multiple cores*” case study (CS3), we additionally had to ensure comparability of sedimentation rates  
245 between sediment cores, since each model assigns a different age value to its sedimentation rate value per  
centimeter. Therefore, we binned sedimentation rate results into 1000-year bins for each age-depth model as well  
as the ensemble model and calculated the weighted averages and their confidence intervals within these bins.  
Inside LANDO, users can change the initial bin size of 1000 years to the desired resolution.

### 2.5.2 Detection and filtering of unreasonable models

250 For cases in which age-depth models do not agree with each other, e.g., “*Inconsistent sequence*” case study (CS2),  
we have built in the option of importing data from measured sediment properties, also known as proxies. Because  
of compositional and density variations of deposits, changes in sedimentation rates imply changes in the deposition  
of proxies (Baud et al., 2021; Biskaborn et al., 2021; Vyse et al., 2021). By including appropriate, independent  
proxy data on lithological changes within the sediment core, we can weight each model based on its performance  
255 to represent these variations in sedimentation rate. Users should provide the independent sediment proxy data as  
file with two columns, namely “*compositedepth*” which should be the measurement depths (as mid-point  
centimeter below sediment surface), and “*value*” representing the values of the proxy. This simplification makes  
it possible to import different available proxies or statistical representations of proxy data, i.e., results from  
ordination techniques (PCA, MDS, etc.), into the optimization process and to visualize the behavior of the age-  
260 depth models in comparison to these proxies.

In order to evaluate the performance, we adapted the fuzzy change point approach by Hollaway et al. (2021) to  
work with our input data and desired outcome on a depth-dependent scale instead of a time series. Similarly to  
Hollaway et al. (2021), our approach firstly detected change points within the proxy data and each modeling system  
output by fitting an ARIMA model to the data and then extract change points by using the “*change point*” R package  
265 (Killick and Eckley, 2014; Killick et al., 2016) on the residuals of the ARIMA model. If we found no change points  
in the proxy data via this approach, we applied the “*change point*” R package on the raw independent sediment  
proxy data instead. Through the additional bootstrapping process introduced by Hollaway et al. (2021), we were  
able to set up confidence intervals for the extracted change points. Subsequently, we searched for the intersection  
between the change points plus their confidence interval for each age-depth model with the independent proxy  
270 data. After converting the change points for both age-depth model and independent proxy data into triangular fuzzy  
numbers, we obtained similarity scores using the Jaccard similarity score of the fuzzy number pairs as described  
in Hollaway et al. (2021). The similarity score can reach numbers between zero (no match) and one (perfect match).  
However, the threshold of excluding an age-depth model from the generated combined model depends on the  
imported proxy data and number of detected change points. Therefore, the user can set the threshold accordingly



275 to their proxy within LANDO, but we have implemented the default value for this threshold to 0.1, which corresponds to an overlap of 10% of the change points between model and proxy data.

In addition to the criterion of preparing the proxy data in the format of “depth vs. value” in a separate file, we suggest using a proxy with a high resolution. As a high-resolution proxy, we define a proxy with more than 50 measurements per meter of core length. For our “*Inconsistent sequence*” case study (CS2), we used high-  
280 resolution elemental proxy data from XRF (X-ray fluorescence) measurement as our independent proxy data. As our evaluation element to optimize the age-depth models, we selected zircon (“Zr”), which itself is an indicator for minerogenic/detrital input (Vyse et al., 2020 and references therein). The zircon proxy data of EN18208 has a resolution of 200 measurements per meter of core length.

To achieve a realistic comparison between sediment cores in the “*Multiple cores*” case study (CS3), we looked at  
285 the individual age-depth model outputs for each sediment core to determine whether an optimization step was required. We have only selected published sediment cores ( $n = 39$ ) so that we can refer to lithological boundaries from the original publication. During the analysis, we saw that nine sediment cores needed to be optimized due to strong inconsistencies between models over the entire length of each core. In twelve cases, where models within the lower section of the cores did not match, we considered proxy-based optimization to improve the model  
290 outcome when high-resolution data was available.

### 2.5.3 Sedimentation rate development over time

To identify similar temporal shifts in sedimentation regimes in our case study “*Multiple cores*” (CS3), we examined our data collection of 62 sediment cores regarding a general tendency in sedimentation rate shifts. First, we considered the 11 700-yr BP (Before Present, i.e., before 1950 CE) boundary as our marker for the change  
295 between Holocene and Pleistocene to separate the datasets (Rasmussen et al., 2006; Lowe and Walker, 2014; Walker et al., 2008). We selected this marker because numerous studies suggest a general difference in sedimentation regimes between these periods (e.g., Brosius et al., 2021; Vyse et al., 2021; Baumer et al., 2021; Bjune et al., 2021; Kublitskiy et al., 2020; Müller et al., 2009; Wolfe, 1996). As some of the models were below the 11 700-yr BP marker, the calculation of the mean sedimentation rate for the Pleistocene featured only a subset  
300 of sediment cores (total number of sediment cores with measurement in Pleistocene: 20). Then, for each age model of the sediment cores in the subset, we used the two-sigma ranges around 11 700 yr BP to determine whether the maximum absolute change occurred exactly at 11 700 yr BP or around our set marker. For this investigation, we changed the bin size to 100-year bins to allow comparison between each modeling system and the combined models. Using maximum from the interquartile ranges of the two-sigma ranges for each model (see supplementary material [Figure S3](#)), we defined the observation period from 8700 to 14 700 yr BP (corresponds to a range of  $\pm$   
305 3000 years). We then checked the data within the time span to see where the maximum change in sedimentation rate occurred. If the calculated age for the new marker was at the edge of our time span, we iteratively increased the outer limit by 100 years (up to a maximum of 18 000 yr BP) to see if the calculated age still reflected the maximum absolute change. We then used the newly defined marker to calculate the mean sedimentation rate for  
310 before and after the marker.

### 2.5.4 Display of models



To display the results from age-depth modeling and sedimentation rate calculation, we decided to create our own plots, instead of reusing the plots from each individual modeling system. Our plot header contains the unique CoreID; additionally, the header indicates whether the user decided to apply a reservoir correction on the radiocarbon data or not. Our single core plots consist of two main panels: On the left-hand side, the panel shows the results from the age-depth modeling process with the calibrated ages (in calibrated years Before Present, i.e., before 1950 CE) on the x-axis and the composite depth of the sediment core (in centimeter) on the inverted y-axis. On the right-hand side, the panel displays the result from the sedimentation rate calculation (in cm/yr, centimeter per year) on the x-axis plotted against the same composite depth on the inverted y-axis. For better readability of the strong variability of sedimentation rate, we used the log scale for the x-axis of the right panel. Generally, LANDO draws the ensemble age-depth model and sedimentation rate in grey with the weighted average as dashed line.

For all models, LANDO will display the median values for age and sedimentation rate as solid lines. Both panels further display the corresponding one-sigma range and two-sigma range per centimeter for each model. Depending on the user's selection, users can plot both sigma ranges, only one of the two sigma ranges, or just the median ages. To include age determination data within the plots, LANDO internally calibrates the radiocarbon data with the "BehronCalibrate" function of the *Behron* package (Haslett and Parnell, 2008; Parnell et al., 2008) with either the IntCal20 (Reimer et al., 2020) or Marine20 (Heaton et al., 2020) calibration curve. Per default, the left panel contains each calibrated age determination as single point with its uncertainty as error bars.

If users decide to filter out unreasonable age-depth models, similar to "Inconsistent sequence" case study (CS2), we added the option to plot the independent proxy data and therefrom derived lithology as an additional panel on the left-hand side for a better interpretability. Further, LANDO highlights the boundaries of lithological change and its confidence interval in both sedimentation rate and age-depth model plots. The optimized plot includes a goodness-of-fit for each involved modeling system to represent the change points at the bottom of the plot.


When using LANDO for multiple sediment cores, the overall plot holds for each sediment core the results from the binned weighted average sedimentation rate calculation (as median sedimentation rate in cm/yr, centimeter per year) against the selected age bins (in calibrated years Before Present, i.e., before 1950 CE) for each modeling system. This visual illustration allows user to compare multiple sediment cores based on the time axis.

For people with color vision deficiency, we incorporated the extra option to plot the resulting age-depth plots with different line styles and textures to support the visual differentiation between each model. Figure S4 in the supplementary material shows the color-blind friendly output created by LANDO. With LANDO we want to support inclusivity in science, but we look forward to feedback from the community on how we can improve LANDO in this regard.

### 3 Results

#### 3.1. "Undisturbed sequence" – Case Study no. 1

All five age-depth models were able to produce an age-depth correlation for sediment core EN18218 ("Lake Rauchuvagytygyn") with only small diversions in between some of the calibrated ages. Figure 2 depicts the two visual outputs produced by LANDO. Panel (a) displays all models side by side, while panel (b) shows the combined output from all models.



350

**Figure 2** – Generated output from LANDO for sediment core EN18218 ( $^{14}\text{C}$  data from Vyse et al., 2021). Panel (a) consists of a comparison between age-depth models from all five implemented modeling systems (left plot) and their calculated sedimentation rate (right plot). Colored solid lines indicate both the median age and median sedimentation rate for all models, while shaded areas represent their respective one-sigma and two-sigma ranges in the same colors with decreasing opacities. Panel (b) shows the ensemble age-depth model (left plot) and its sedimentation rate (right plot). The dashed line in panel (b) represents the weighted average age estimates (left plot) and the weighted average sedimentation rates (right plot) for the ensemble model, while the grey area represents the two-sigma uncertainty, i.e., the outermost limits of two-sigma ranges from all models. Both plots on the left of (a) and (b) show the depth below sediment surface on the inverted y-axis as composite depth of the sediment core in centimeter (cm) and the calibrated ages on the x-axis in calibrated years Before Present (cal. yr BP, i.e., before 1950 CE). Black circles within (a) and (b) indicate the calibrated dating points

360




with their uncertainty as error bar, which were calibrated using the *IntCal20* calibration curve (Reimer et al., 2020). The plots on the right display the sedimentation rate in centimeter per year (cm/yr, x-axis as log-scale) against the depth below sediment surface as the composite depth of the sediment core in centimeter (cm, inverted y-axis).

All models revealed highest sedimentation rates for the interval between 108 and 133 cm. Mean values ranged from 0.242 cm/yr (*hamstr*) to 0.764 cm/yr (*clam*), whereas the median sedimentation rate varied between 0.107 cm/yr (*Bacon*) and 0.314 cm/yr (*clam*). In three instances, the majority of models noticeable dropped to lower sedimentation rate values. We found the first significant downshift between 57 cm and 65 cm, but in contrast to the following declines, *hamstr* immediately increased the median sedimentation rate by tenfold from 0.015 to 0.15 cm/yr from 64 cm to 66 cm. Subsequent decreases occurred between 222 cm and 249 cm as well as between 339 and 366 cm with median sedimentation rates from 0.013 cm/yr (*hamstr*) to 0.025 cm/yr (*Bacon*) and from 0.012 cm/yr (*hamstr*) to 0.027 cm/yr (*Bacon*), respectively. In contrast to the upper segment, the models showed a stronger disagreement among each other in the lower segment of the sediment core, starting at 504 cm with larger varying mean and median values for sedimentation rate.

In our ensemble model, we found the highest value for weighted average sedimentation rate at 128 cm with 0.4483 cm/yr (two-sigma range: 0.032 - 2.338 cm/yr), which corresponded to weighted average age estimate of 4846 cal yr BP (two-sigma range: 4301 - 5384 cal yr BP). When considering the cumulative two-sigma uncertainty range of the ensemble model, our generated output covered sedimentation rates between 0.002 cm/yr and 2.486 cm/yr over the entire core.

### 3.2. “Inconsistent sequence” – Case Study no. 2

For the second case study, four out of five modeling systems produced an output for sediment core EN18208 (“Lake Ilirney”). The modeling system *clam* was unable to produce an age-depth model for this core. Figure 3 shows the visual outputs with all models in panel (a) and the combined model in panel (b). Figure 4 consists of three panels showing the results from the proxy-based optimization process using zircon (Zr). Panel (a) shows the visual output from the optimization process, while panel (b) and (c) illustrate the optimized age-depth model with the highest matching score and the resulting ensemble model, respectively.






390 **Figure 3** – Generated output from LANDO for sediment core EN18218 (OSL and  $^{14}\text{C}$  data from Vyse et al.,  
 2020). Panel (a) consists of a comparison between age-depth models from four out of five implemented modeling  
 systems (left plot) and their calculated sedimentation rate (right plot). The modeling system clam was unable to  
 produce an age-depth model for this core. Colored solid lines indicate both the median age and median  
 sedimentation rate for all four models, while shaded areas represent their respective one-sigma and two-sigma  
 395 ranges in the same colors with decreasing opacities. Panel (b) shows the ensemble age-depth model (left plot)  
 and its sedimentation rate (right plot). The dashed line in panel (b) represents the weighted average age  
 estimates (left plot) and the weighted average sedimentation rates (right plot) for the ensemble model, while the  
 grey area represents the two-sigma uncertainty, i.e., the outermost limit of two-sigma ranges from all four  
 models. Both plots on the left of (a) and (b) show the depth below sediment surface on the inverted y-axis as  
 400 composite depth of the sediment core in centimeter (cm) and the calibrated ages on the x-axis in calibrated years



405 *Before Present (cal. yr BP, i.e., before 1950 CE). Black circles within (a) and (b) indicate the calibrated dating points with their uncertainty as error bar, which were calibrated using the IntCal20 calibration curve (Reimer et al., 2020). The plots on the right display the sedimentation rate in centimeter per year (cm/yr, x-axis as log-scale) against the depth below sediment surface as the composite depth of the sediment core in centimeter (cm, inverted y-axis).*

410 Starting from the sediment surface, the first divergence between the age-depth models occurred with the age determination point at 184 cm. *Undatable* started to increase sedimentation rate between 183 cm and 185 cm with median sedimentation rate change from 0.123 cm/yr to 0.273 cm/yr. *Undatable* then followed the younger OSL dates (375.5 cm / 475.5 cm), while the other three models continued with the radiocarbon date at 210 cm. Shortly  
415 after both *hamstr* and *Bchron* increased sedimentation rates and directly at 325.5 cm went along the radiocarbon date. *Bchron* changed the median sedimentation rate between 209 cm and 211 cm from 0.007 cm/yr to 0.033 cm/yr, whereas *hamstr* increased from 0.004 cm/yr to 0.06 cm/yr between 214 cm and 216 cm. In contrast, *Bacon* assumed no abrupt changes in sedimentation rate while following the older age determination dates until 494.5 cm, before keeping a rather steady accumulation without any dating points (mean sedimentation rate: 0.04 – 0.06 cm/yr). At  
415 the depth of 795 cm, the three modeling systems *Undatable*, *hamstr*, and *Bchron* again overlapped their paths, before splitting at the existing dating point at 966 cm.





**Figure 4** – Optimized visual output for EN18208 (OSL and  $^{14}\text{C}$  data from Vyse et al., 2020). We used high-resolution X-ray fluorescence (XRF) measurements of zircon (Zr) as independent proxy to evaluate model performance to represent lithological changes. Panel (a) extends the existing panel (a) of Figure 3 by adding a plot on the left to show the proxy-derived lithology used to filter unreasonable models. This added plot consists of the proxy measurements of Zr (in counts per second) along the depth below sediment surface as the composite depth of the sediment core in centimeter (cm) and the derived lithological boundaries (solid horizontal lines) plus their uncertainty range (dashed horizontal lines). Both age-depth model and sedimentation rate plot contain the same lithological boundaries as visual aid. The text box in the bottom middle lists the models with their matching score related to the proxy-derived lithology. Panel (b) shows the model (Bchron) with the highest matching score (0.0796). Panel (c) depicts our ensemble model based on this model. The age-depth models displayed in panel (b) and (c) show strong similarities with the age-depth model developed by Vyse et al. (2020).

420  
 425  
 430  
 435

During the optimization process, our adapted algorithm located four lithological boundaries with its uncertainty range from the independent proxy data: 189.5 cm (182 – 192.5 cm), 646 cm (638 – 657 cm), 890.5 cm (874 – 912 cm), and 1051.5 cm (1043 – 1061.5 cm). We found the highest matching score from the optimization for Bchron (Score: 0.0796). Table 4 shows the average sedimentation rate for each proxy-derived lithological unit (PLU) of the ensemble model of EN1808.

**Table 4** – Average sedimentation rate of EN18208 divided into proxy-derived lithological units. The calibrated mean model range indicates the mean age estimates of the ensemble model for the corresponding depths of the proxy-derived lithological unit (PLU).

Proxy-derived lithological unit	Corresponding depths below sediment surface [cm]	Calibrated mean model range [cal yr BP]	Average sedimentation rate [cm/yr]
PLU1	0 – 190	-64 – 18295	0.0144
PLU2	190 – 646	18295 – 29363	0.1335
PLU3	646 – 891	29363 – 35308	0.1259
PLU4	891 - 1052	35308 – 40300	0.0634


### 3.3. “Multiple cores” – Case Study no. 3

440 In contrast to the previous case studies, this case study focused on studying development of sedimentation rates over time, with the emphasis on the transition from the Holocene to the Pleistocene. We used age determination data from 39 published sediment cores to show the standard output of LANDO for multiple sediment cores, while using both published and unpublished age data for the analyses. Figure 5 shows the ensemble models with weighted average sedimentation rates binned into 1000-year bins from our multi-core investigation with 39 published sediment cores (see Figure S1 for the individual models in the supplementary material). We set the boundaries from 0 to 21 000 cal yr BP within these figures to cover the time span from the present to the Last Glacial Maximum (LGM) (Clark et al., 2009). Below the number for each core in Figure 5 are the proxies used for their optimization. In 19 out of 62 cases within our entire collection, the ensemble model was based on four out of five models, as neither clam or Undatable was able to find a suitable age-depth model (for more details, please see Table S1 in the supplementary material). The maximum time span covered by the sediment cores varied between 2000 yr BP (CoreID: PG1972) and 320 000 yr BP (CoreID: PG1351). The average non-optimized

445  
 450




sedimentation rate ranged between 0.006 cm/yr (CoreID: 16-KP-03-L10) and 0.198 cm/yr (CoreID: PG1228). In total, we optimized seven sediment cores, as in most cases neither high-resolution data was available nor the provided proxy data represented a lithological proxy when crosschecked with the original publication. From these  
 455 seven sediment cores, we reconstructed the proxy-based lithology twice with TOC as a low-resolution proxy (CoreID: PG1228 & PG1437).




**Figure 5** – Optimized combined models for 39 published sediment cores displayed as weighted average  
 460 sedimentation rate (in centimeter per year, cm/yr – y-axis) binned into 1000-year bins (in calibrated years Before Present, cal. yr BP, i.e. before 1950 CE – x-axis) for the last 21 000 years. Dashed line represents the weighted average sedimentation rate, whereas the grey areas are the respective two-sigma ranges. Each grid cell contains the unique core identifier of each involved sediment core. In seven cases, the letters below each  
 number give the name of the independent proxy used for optimization process.


To visualize the difference in sedimentation rates between two neighboring and fundamentally different  
 465 environmental settings, i.e. Pleistocene glacial and Holocene interglacial, we used the datasets that were split at the Holocene-Pleistocene boundary at 11 700 yr BP. Figure 6 shows the mean sedimentation rate for Holocene and Pleistocene for each model with its one-sigma uncertainty. Figure S3 in the supplementary material gives an overview over the overall uncertainty for all models. Among all models, clam models have the lowest range on average for both Holocene (0.0125 cm/yr) and Pleistocene (0.0066 cm/yr), while the Undatable models show the greatest uncertainty on average in the Holocene (0.1061 cm/yr) as well as the combined models for the Pleistocene (0.0405 cm/yr). The sediment core PG1228 (latitude: 74.473° N) showed the highest individual sedimentation rate for the Holocene (median sedimentation rate: 2.1929 cm/yr). We observed a significant reduction of 89 % for the  
 470 optimized model of the same core (0.0684 cm/yr), compared to its combined model (0.6244 cm/yr).



- 475 **Figure 6** – Average sedimentation rate in centimeter per year (cm/yr) for each sediment core in our data collection of 62 sediment cores divided into Holocene dataset (from present up to 11 700 yr BP, orange lines) and Pleistocene dataset (from 11 700 yr BP up to a maximum of 115 000 yr BP, blue lines). Each plot displays the one-sigma range of sedimentation rate within each dataset for each model and sediment core. In addition, filled circles represent the mean value for the optimized models.
- 480 For our data compilation, we found the largest absolute change in sedimentation rates within the modeling systems on average between 10 500 and 13 800 yr BP (Figure 7). For our combined and optimized models, however, the largest change averaged between 10 800 yr BP and 10 900 yr BP. Still, all sediment cores covered the entire range of our initial time span from 8700 to 14 700 yr BP within the models. Using the results of the largest change in sedimentation rate for each sediment core and model as new markers, we again split the datasets into two separate
- 485 datasets. One dataset contained mostly Holocene sedimentation rate values (Holocene dataset), while the other contained mostly Pleistocene values (Pleistocene dataset). Therefore, the initial display (Figure 6) changed slightly to Figure 8. Most notable was the increase in total number of sediment cores in Pleistocene dataset with an individual separation (n = 39) compared to the Pleistocene dataset with the separation at 11 700 yr BP (n = 20).



490 **Figure 7** – Boxplot representing the years with the biggest absolute change in sedimentation rate for our data collection of 62 sediment cores. Sedimentation rate results from each model binned into 100-year bins to allow comparisons between the modeling systems. The initial observation time span covers 8700 to 14 700 yr BP. The orange line corresponds to the median value for each model.



495 **Figure 8** – Average sedimentation rate in centimeter per year (cm/yr) for each sediment core in our data collection of 62 sediment cores divided into Holocene dataset (orange lines) and Pleistocene dataset (blue lines). The exact value for the split of the datasets for each individual core and each model depends on the results of the maximum change in sedimentation rate within the observation period 8700 to 14 700 yr BP. Each plot displays



500 *the one-sigma range of sedimentation rate within each dataset for each model and sediment core. In addition,  
filled circles represent the mean value for the optimized models.*

## Discussion

### 4.1. Assessment of different case studies

By comparing the cases for the two single sediment cores, it becomes clear how age-depth correlations may diverge depending on the individual modeling system and its treatment of available dating points (cf. Wright et al., 2017; Trachsel and Telford, 2017; Lacourse and Gajewski, 2020). In the case of EN18218 (“Undisturbed sequence” – CS1), all five implemented modeling systems yield an agreeing and continuous chronology. However, the two radiocarbon dates at 81.25 cm and 114.75 cm have significant impact on the model’s interpretation for these depths. Vyse et al. (2021) argued that these two dates are outliers resulting from reworking and mixing effects within the sediment column. According to the authors, no additional proxy data from EN18218 would support the immediate increase in sedimentation rate for these depths and hence, they excluded both dates from the modeling process. Because we are not considering any additional proxy data to evaluate age-depth models in their geoscientific context, but rather include all provided age determination data into the modeling process, the consideration of these two radiocarbon dates on the basis of all available models leads to higher sedimentation rate. Nonetheless, the comprehensive application of the different modeling systems may help to identify doubtful dating points.

We saw a strong disagreement between the modeling systems in the case of sediment record EN18208 (“Inconsistent sequence” – CS2), which we expected prior to the execution of our application, due to the scattered dating points in the original data. Vyse et al. (2020) linked this scatter of age data points observed in the interval between 282 and 755 cm of EN18208 to the redeposition of older carbon. They implied that to produce reliable age-depth model they had to exclude both OSL and radiocarbon dating points for these depths. However, our optimized combined model agrees with their established age-depth model and can reproduce the characteristics of the existing model by Vyse et al. (2020), without removing dating points. In addition, in three out of four cases, our proxy-derived lithology with its uncertainty matches the lithological boundaries set by the authors of the EN18208 study, according to criteria based on acoustic sub-bottom profiling. Only the first original boundary (196 cm) is outside our confidence interval from 182 cm to 192 cm. We still showed that our approach could set logical boundaries for sediment cores by solely relying on high-resolution proxy data.

Even though LANDO can produce age-depth models for multiple sediment cores (“Multiple cores” – CS3), we must assume limitations in the geoscientific validity for some of the results. In a few cases, an optimization of age-depth models with independent proxy data would have been necessary, but such independent data were inaccessible or did not exist. As for these cases age-depth correlations between implemented modeling systems seem to disagree (see [Figure S1](#) in the supplementary material), the results from our combined model might over- or underestimate the true sedimentation rate. On the other hand, optimization using proxy data can reduce these biases. For instance, during the examination of the Holocene and the Pleistocene sedimentation rates ([Figure 6](#)), we noticed that one sediment core (PG1228) had an extremely high mean sedimentation rate for the Holocene dataset in *Undatable*. Similar to the second case study (“Inconsistent sequence” – CS2), we found scattered age data points for this sediment core, which influenced the modeling process of *Undatable*. Further, the result then affected our combined model by increasing the overall sedimentation rate for the Holocene in this core. However,



LANDO identified the *Undatable* model as an outlier based on the lithology established through independent TOC proxy data. The optimized model then agreed well with the original publication by Andreev et al., (2003), which  
540 further increased the validity of our approach. Our findings suggest that high-resolution proxy data should accompany geochronological studies to enable a more concise and realistic assessment of the development of sedimentation rates over time in high latitude lake systems.

The detection of sedimentation rate change as indicator for the Holocene-Pleistocene boundary yielded contrasting results. While the results from *hamstr* were closest to the 11 700-year boundary, all other modeling systems place  
545 the largest change in sedimentation rate either before or after 11 700 yr BP. We hypothesize that three factors may have influenced all model results. (1) The age uncertainty (one-sigma range) within each individual model varied on average between 1000 and 3000 years for the period of 11 600 to 11 800 yr BP (Figure S3 in the supplementary material). This wide range of uncertainty does not provide confidence in pinpointing the boundary to an exact time slice. We expect that a higher amount of dating points close to the Holocene-Pleistocene boundary could constrain  
550 the models (Blaauw et al., 2018; Lacourse and Gajewski, 2020; Trachsel and Telford, 2017), which would lead to a better estimate of the boundary. (2) The age output for each model is not evenly distributed, which means that in the period from 11 600 to 11 800 yr BP there are different numbers of observations for each core and each modeling system. We took this behavior into account by using binning (Alasadi and Bhaya, 2017). Otherwise, an interpolation between both age and sedimentation rate values could lead to potential biases in the interpretation.  
555 (3) While we assumed in our first setup that the main sedimentation rate change would occur at 11 700 yr BP consistently for all sediment cores (Figure 6), we cannot rule out the possibility that the sedimentation rate has changed significantly at different times for different lake systems. As our data collection covers a large area both in latitude and longitude (Figure 1), the variability between the models indicate the local variability between the climate and lithological preferences of the lake catchment for the involved sediment cores (e.g., Lozhkin et al.,  
560 2018; Finkenbinder et al., 2015; Anderson and Lozhkin, 2015; Kokorowski et al., 2008; Biskaborn et al., 2016; Courtin et al., 2021).

#### 4.2. Design of LANDO

From the beginning of the development of LANDO, we decided to integrate most of the default settings for each modeling system as default values (Table 2). Regional studies, such as the one performed by Goring et al. (2012),  
565 have shown that specific prior information for the Bayesian modeling systems are needed to best fit the models to lakes within a geographical area. Without this regional information, changing settings within the modeling system to an arbitrary higher or lower value without considering the regional diversity could lead to under- or overfitting, if the constraints are too loose or too strict (Trachsel and Telford, 2017). For the special case that users have in-depth knowledge for one lake or multiple lake system, users can easily adapt these parameters within LANDO, as  
570 we have made these settings accessible in the Jupyter Notebook itself.

Part of the reason we made this decision was that we acquired external age determination datasets where we may not necessarily have all the essential information to specify each model. But we also wanted to simplify the process for users who do not have in-depth modeling knowledge. By using the default values, we can compare models based on their ability to work with the available data. On the other hand, we are sure that the developers have set  
575 their default values based on systematic testing. Since we did not tune the age-depth models to the existing core, i.e. changing the parameters within each modeling system, we generated “uninformed” models that solely work



with the available age determination data. By combining these “uninformed” models into one model, we have created an ensemble model that we consider to be data-driven and “semi-informed”.

580 The advantage of this data-driven, semi-informed model approach is that we are reducing the risk of overfitting by considering the uncertainty of all modeling systems. This allows us to reevaluate existing geoscientific interpretations with larger uncertainty by taking advantage of the ensemble outcome. Additionally, we found that the more information is accessible to generate age-depth models, the more accurate and less uncertain these models become. A higher density of age determination along the depth of the sediment core is desirable for future drilling campaigns (cf. Blaauw et al., 2018).

585 The disadvantage arises in our second case study (“*Inconsistent sequence*” – CS2) and the multi-core investigation (“*Multiple cores*” – CS3). For both cases we needed the optimization step to narrow down the most suitable age-depth models for each sediment core, since the unoptimized uncertainty band was otherwise too wide for a clear interpretation. The optimization requires additional and independent proxy data, which are not available for some of our cores, especially for sediment cores obtained some decades ago. Our optimizing step is therefore mainly  
590 suitable for recently retrieved and analyzed sediment cores.

In addition to the assessment of age-modeling quality, we also checked the time and effort to conduct dating routines. We saw that *Bacon* had the highest runtime overall in all three case studies of our study design, which we link to our adjustment of the “ssize” parameter from 2000 (per default) to 8000 within the application. We increased this value to ensure good MCMC mixing for problematic cores, as suggested by Blaauw et al., (2021),  
595 as well as to guarantee we had enough iterations for our summarizing statistics to compare with other modeling systems. If users decide to reduce the value of “ssize”, we implemented an iterative process, which checks whether *Bacon* produced enough iterations. If this is not the case, then LANDO will iteratively rerun the same sediment core with a higher “ssize” to produce 10 000 iterations.

600 One unique feature of our application is the predominant use of parallelization within the age-depth modeling of multiple sediment cores. For instance, we used the “Dask” back-end for our sedimentation rate calculation. The advantage over popular Scala-based “Apache Spark” and its Python interface “PySpark” (Zaharia et al., 2016) is that the “Dask” back-end is Python-based and well integrated into the Python ecosystem (cf. Dask Development Team, 2016). Therefore, “Dask” natively works with Python packages already implemented in LANDO. The key difference is that “Dask” does neither provide a query optimizer nor rely on Map-Shuffle-Reduce, a data processing  
605 technique for distributed computing, but instead uses a generic task scheduling (cf. Dask Development Team, 2016). Still, parallelization libraries and back-ends provide LANDO with additional speed-up that can promote future multi-core studies.

610 Within the ensemble model, we faced the challenge that the combination of all age distributions from the underlying age-depth models per centimeter represents a multi-modal distribution, especially in cases such as the “*Inconsistent sequence*” case study (CS2). It also means that the output of the ensemble model in these cases is susceptible to inclusion/exclusion of any model. However, we consider using the weighted average median age to be a suitable solution for the multi-model distribution problem, as it is a good indicator on the most probable age within each centimeter based on all modeling systems. But we advise users to use the age confidence intervals per centimeter in subsequent analyses, instead of relying solely on the weighted average median age (cf. Telford et al.,



615 2004). By optimizing the ensemble model with the ability to include independent proxy data, users can increase  
the likelihood of a more probable mean age for their sediment core.

#### 4.3. Technical specifications of LANDO

In the further course of development, we decided to limit the resolution of the age-depth correlations. Using a  
resolution of one-centimeter increments allows us to match most proxy measurements from each sediment core  
620 with our age-depth models, apart from high-resolution measurement, such as XRF measurements. To allow a  
matching with high-resolution proxy data, we tested for a higher resolution of 0.25 cm for our application. In the  
single sediment core cases (CS1 and CS2), this change did not affect the workflow of LANDO. In turn, the  
“Multiple cores” case (CS3) ran into memory issues. Since the SoS notebook and our parallel back-ends store the  
result data frames in memory, expanding the resulting data frames to a 0.25 cm resolution causes a fourfold  
625 increase in memory use, which limits our capability to run our application on a single laptop. As an intermediate  
solution, we stored the results from each parallelization worker on disk to free the memory and performed  
combining operations later. Based on this experience, we recommend working with data centers or increasing the  
available main memory (RAM) of the operating computer for multi-core studies with expected high-resolution  
output.

630 Another advantage of parallelization is that most modeling systems only run on one CPU/thread. Nowadays,  
however, both personal computers and data centers are made up of multiple CPUs/threads. Especially for larger  
multi-site studies, our application has the advantage of cutting the overall computing time by running each  
modeling system on multiple CPUs/threads simultaneously, even for personal computers. In comparison to serial  
execution of multiple models on one CPU/thread, which would take several hours, our parallel execution reduced  
635 the computing time per modeling system by a factor up to four. When considering that our setup consisted of six  
CPUs (12 threads) and 16 GB RAM, user can even further increase this factor by using larger computing facilities.

Sediment core length is the most limiting factor that determines the overall computing time in our application.  
However, we want to ensure that users can model each sediment core over its entire length to match proxy data  
with the correct age-depth correlations. Within our LANDO system, we faced this problem by using extrapolation  
640 to calculate ages beyond available dating points. The exception here is the modeling system *Undatable*, which  
models only between the first and last dating point, as these two dating points act as anchors for the bootstrapping  
process (Lougheed and Obrochta, 2019). As a result, we saw the sedimentation rate dropping twice to zero at the  
end of the sedimentation rate calculations. We link this behavior to the end of the individual modeling processes  
of *Undatable* as well as the other implemented systems.

645 Extrapolating the age-depth models beyond age determination points always bares the risk that the extrapolated  
dates do not reflect the actual age. The implemented modeling systems account for this circumstance by increasing  
the uncertainty for these undated regions (Blaauw, 2010). While we are aware of this potential issue, we wanted  
to allow users to take advantage of the full age-depth coverage for their sediment core. Blaauw et al. (2018) pointed  
out in their findings that “most existing late-Quaternary studies contain fewer than one date per millennium” and  
650 recommended to increase the number of dating points to “a minimum of 2 dates per millennium”. This  
recommendation would further decrease the need of extrapolation and reduce the overall uncertainty of age-depth  
models. We agree that more age control can improve the age-depth modeling results, but until the associated costs



to analyze organic material for radiocarbon dating do not decrease more significantly (Hajdas et al., 2021; Zander et al., 2020), we recommend LANDO as tool to improve age-depth modeling.

#### 655 4.4. Current and future model implementation in LANDO

During the development of our approach, we realized that some programs were not executable or parallelizable under the current circumstances. For instance, we tested *OxCal* 4.4 as stand-alone version on Windows with NodeJS (version 12.13.1.0) and the R package “oxcAAR” (Martin et al., 2021) within our application. In the case of EN18208, execution duration was above 3 hours until the notebook lost connection to the *OxCal* interface.

660 Furthermore, some cores never fully reached convergence within *OxCal*. We tried adapting our set-ups including changing the internal constraints, i.e. placement and number of boundaries, or using different depositions models, i.e. alternating between sequential model (“Sequence()”) and Poisson-process deposition model (“P\_Sequence()”). According to Bronk Ramsey and Lee (2013), the long-term plan of *OxCal* is to make the entire source code openly accessible, which we fully support. An open source code would allow us to identify the current bottleneck so that  
665 we could implement *OxCal* in a future release.

To determine the most fitting age-depth model through the *clam* modeling software, we added the “best fit” option to LANDO by default. The “best fit” option utilizes the negative log fit results from all *clam* outputs and identifies the fit with the lowest result as best fit. We included two further exclusion criteria for *clam* models within LANDO:  
670 if a) there are too many age reversals within the models, or b) the fit reaches infinity. Under specific circumstances, some sediment cores will not have a fitting model, as is the case, for instance, in the “*Inconsistent sequence*” case study (CS2). Including models that do not fit the data would lead to erroneous estimations of the age-depth correlation. This comes with the cost of losing an established model in the combined model, if no fitting *clam* model is available. However, we think that the benefit of having a more fitting model outweighs this cost.

Although *Undatable* is open source and the fastest modeling system within LANDO, its original development  
675 environment (MATLAB) is not free of charge. That is why we implemented *Undatable* in the open source MATLAB-equivalent Octave. Since the Octave version of *Undatable* was slower than the original MATLAB version, we used the parallelization package “parallel” (Fujiwara et al., 2021) to provide comparable results in terms of computing time. To use *Undatable* with MATLAB within our application, users must acquire a license of MATLAB and link the MATLAB kernel to their license. Unfortunately, we do not have the capacity to provide  
680 individual licenses with LANDO. For users with an active MATLAB license, we provide in the repository mentioned in the “Code and data availability” section the appropriate code to run the MATLAB version of *Undatable* in LANDO.

A potential expansion option of LANDO within the multi-language environment is to extent the application and allow future data analysis to use powerful tools, such as Python’s machine learning libraries, e.g., keras (Chollet  
685 and and others, 2015) and tensorflow (Abadi et al., 2016). We anticipate that other developers can use LANDO as their starting point in building larger limnological data analysis application.

#### 4 Conclusion

This paper introduced our application LANDO – a linked age-depth modeling notebook approach. We presented an improved age-depth modeling procedure for sediment cores from high-latitude lake systems by linking five  
690 established systems: *Bacon*, *Bchron*, *clam*, *hamstr*, and *Undatable*. The added value of our application is the



reduced effort to use established modeling systems in a single Jupyter Notebook for both single and multiple dating series and at the same time make the results comparable. In addition, we introduced an ensemble model that uses the output from all models to create a more robust age-depth correlation. In the case of scattered age determination data, we further implemented an adapted version of the fuzzy change point approach that allow users to integrate independent proxy data as indicator of lithological changes. This option helps evaluate the performance of modeling systems across lithological boundaries while providing a more reliable ensemble age-depth model by filtering inappropriate model runs for problematic datasets. Our application also allows users to run large datasets with multiple sediment cores in parallel to reduce the overall computation time. In our data collection of 62 sediment cores from northern lake systems at high latitudes, we found that the main regime changes in sedimentation rates do not occur synchronously for all lakes at the Pleistocene-Holocene boundary. However, we linked this behavior to the uncertainty within the modeling process as well as the local variability of the sediment cores within the collection.

#### Code and data availability

The LANDO code is accessible at GitHub (<https://github.com/GPawi/LANDO>) (Pfalz, 2021). We provide files containing accessible links to the used datasets, contact details for unpublished data, and five example spreadsheets. Contact details comprise names of research group and personal communication addresses of working group leaders. A stand-alone version of the LANDO application will be available for download upon publication.

#### Competing interests

The authors declare that they have no conflict of interest.

#### Author contribution

GP wrote the manuscript with inputs from all co-authors. GP developed the application, designed and implemented the LANDO system, and conducted testing. BKB, BD, and JCF advised and supervised the work of GP. BKB, BD, LS, and DAS provided published and unpublished age determination data for this publication.

#### Acknowledgements

The authors acknowledge the support of the Helmholtz Einstein Berlin International Berlin Research School in Data Science (HEIBRiDS), the Alfred Wegener Institute - Helmholtz Centre for Polar and Marine Research, the Einstein Center Digital Future, the Humboldt University of Berlin, and the Ministry of Education of the Russian Federation as part of a state task (project no. FSZN-2020-0016). The authors highly appreciate all the work that went into developing the stand-alone versions of each modeling system. Hence, we like to thank Maarten Blaauw, Andrew C. Parnell, Andrew Dolman, Bryan C. Lougheed, and Stephen P. Obrochta for their continuous work on *Bacon*, *Bchron*, *clam*, *hamstr*, and *Undatable*.

#### References

Abadi, M., Barham, P., Chen, J., Chen, Z., Davis, A., Dean, J., Devin, M., Ghemawat, S., Irving, G., Isard, M., Kudlur, M., Levenberg, J., Monga, R., Moore, S., Murray, D. G., Steiner, B., Tucker, P., Vasudevan, V., Warden, P., Wicke, M., Yu, Y., and Zheng, X.: TensorFlow: A system for large-scale machine learning, in: 12th USENIX Symposium on Operating Systems Design and Implementation (OSDI 16), 265–283,



[https://doi.org/10.1016/0076-6879\(83\)01039-3](https://doi.org/10.1016/0076-6879(83)01039-3), 2016.

Abbott, M. B. and Stafford, T. W.: Radiocarbon geochemistry of modern and Ancient Arctic lake systems, Baffin Island, Canada, *Quat. Res.*, 45, 300–311, <https://doi.org/10.1006/qres.1996.0031>, 1996.

730 Alasadi, S. A. and Bhaya, W. S.: Review of data preprocessing techniques in data mining, *J. Eng. Appl. Sci.*, 12, 4102–4107, <https://doi.org/10.3923/jeasci.2017.4102.4107>, 2017.

Anderson, P. M. and Lozhkin, A. V.: Late Quaternary vegetation of Chukotka (Northeast Russia), implications for Glacial and Holocene environments of Beringia, *Quat. Sci. Rev.*, 107, 112–128, <https://doi.org/10.1016/j.quascirev.2014.10.016>, 2015.

735 Andreev, A. A., Tarasov, P. E., Siegert, C., Ebel, T., Klimanov, V. A., Melles, M., Bobrov, A. A., Dereviagin, A. Y., Lubinski, D. J., and Hubberten, H.-W.: Late Pleistocene and Holocene vegetation and climate on the northern Taymyr Peninsula, Arctic Russia, 32, 484–505, <https://doi.org/10.1080/03009480310003388>, 2003.

Appleby, P. G.: Three decades of dating recent sediments by fallout radionuclides: A review, 18, 83–93, <https://doi.org/10.1177/0959683607085598>, 2008.

740 Ascough, P., Cook, G., and Dugmore, A.: Methodological approaches to determining the marine radiocarbon reservoir effect, *Prog. Phys. Geogr.*, 29, 532–547, <https://doi.org/10.1191/0309133305pp461ra>, 2005.

Austin, W. E. N., Bard, E., Hunt, J. B., Kroon, D., and Peacock, J. D.: The  $^{14}\text{C}$  Age of the Icelandic Vedde Ash: Implications for Younger Dryas Marine Reservoir Age Corrections, *Radiocarbon*, 37, 53–62, <https://doi.org/10.1017/S0033822200014788>, 1995.

745 Bao, R., McNichol, A. P., Hemingway, J. D., Lardie Gaylord, M. C., and Eglinton, T. I.: Influence of different acid treatments on the radiocarbon content spectrum of sedimentary organic matter determined by RPO/accelerator mass spectrometry, *Radiocarbon*, 61, 395–413, <https://doi.org/10.1017/RDC.2018.125>, 2019.

Baud, A., Jenny, J. P., Francus, P., and Gregory-Eaves, I.: Global acceleration of lake sediment accumulation rates associated with recent human population growth and land-use changes, *J. Paleolimnol.*, 66, 453–467,

750 <https://doi.org/10.1007/s10933-021-00217-6>, 2021.

Baumer, M. M., Wagner, B., Meyer, H., Leicher, N., Lenz, M., Fedorov, G., Pestryakova, L. A., and Melles, M.: Climatic and environmental changes in the Yana Highlands of north-eastern Siberia over the last c. 57 000 years, derived from a sediment core from Lake Emanda, 50, 114–133, <https://doi.org/10.1111/bor.12476>, 2021.

Bayer, M.: SQLAlchemy, in: *The Architecture of Open Source Applications Volume II: Structure, Scale, and a Few More Fearless Hacks*, edited by: Brown, A. and Wilson, G., aosabook.org, 2012.

Biskaborn, B. K., Herzschuh, U., Bolshiyarov, D. Y., Schwamborn, G., and Diekmann, B.: Thermokarst processes and depositional events in a Tundra Lake, Northeastern Siberia, *Permafrost. Periglac. Process.*, 24, 160–174, <https://doi.org/10.1002/ppp.1769>, 2013.

760 Biskaborn, B. K., Subetto, D. A., Savelieva, L. A., Vakhrameeva, P. S., Hansche, A., Herzschuh, U., Klemm, J., Heinecke, L., Pestryakova, L. A., Meyer, H., Kuhn, G., and Diekmann, B.: Late Quaternary vegetation and lake system dynamics in north-eastern Siberia: Implications for seasonal climate variability, *Quat. Sci. Rev.*, 147,



406–421, <https://doi.org/10.1016/j.quascirev.2015.08.014>, 2016.

Biskaborn, B. K., Nazarova, L., Pestryakova, L. A., Syrykh, L., Funck, K., Meyer, H., Chaplgin, B., Vyse, S.,  
765 Gorodnichev, R., Zakharov, E., Wang, R., Schwamborn, G., Bailey, H. L., and Diekmann, B.: Spatial  
distribution of environmental indicators in surface sediments of Lake Bolshoe Toko, Yakutia, Russia, 16, 4023–  
4049, <https://doi.org/10.5194/bg-16-4023-2019>, 2019.

Biskaborn, B. K., Nazarova, L., Kröger, T., Pestryakova, L. A., Syrykh, L., Pfalz, G., Herzsuh, U., and  
Diekmann, B.: Late Quaternary Climate Reconstruction and Lead-Lag Relationships of Biotic and Sediment-  
770 Geochemical Indicators at Lake Bolshoe Toko, Siberia, *Front. Earth Sci.*, 9, 703,  
<https://doi.org/10.3389/feart.2021.737353>, 2021.

Bjune, A. E., Greve Alsos, I., Brendryen, J., Edwards, M. E., Hafliason, H., Johansen, M. S., Mangerud, J.,  
Paus, A., Regnéll, C., Svendsen, J., and Clarke, C. L.: Rapid climate changes during the Lateglacial and the early  
Holocene as seen from plant community dynamics in the Polar Urals, Russia, *J. Quat. Sci.*, 00, jqs.3352,  
<https://doi.org/10.1002/jqs.3352>, 2021.

775 Blaauw, M.: Methods and code for “classical” age-modelling of radiocarbon sequences, *Quat. Geochronol.*, 5,  
512–518, <https://doi.org/10.1016/j.quageo.2010.01.002>, 2010.

Blaauw, M.: clam: Classical Age-Depth Modelling of Cores from Deposits, [https://cran.r-  
project.org/package=clam](https://cran.r-project.org/package=clam), 2021.

Blaauw, M. and Christen, J. A.: Flexible paleoclimate age-depth models using an autoregressive gamma process,  
780 *Bayesian Anal.*, 6, 457–474, <https://doi.org/10.1214/11-BA618>, 2011.

Blaauw, M. and Heegaard, E.: Estimation of Age-Depth Relationships, 379–413, [https://doi.org/10.1007/978-94-  
007-2745-8\\_12](https://doi.org/10.1007/978-94-007-2745-8_12), 2012.

Blaauw, M., Christen, J. A., Bennett, K. D., and Reimer, P. J.: Double the dates and go for Bayes — Impacts of  
model choice, dating density and quality on chronologies, *Quat. Sci. Rev.*, 188, 58–66,  
785 <https://doi.org/10.1016/j.quascirev.2018.03.032>, 2018.

Blaauw, M., Christen, J. A., and Aquino Lopez, M. A.: rbacon: Age-Depth Modelling using Bayesian Statistics,  
<https://cran.r-project.org/package=rbacon>, 2021.

Bradley, R. S.: *Paleoclimatology: Reconstructing Climates of the Quaternary Second Edition*, 3rd ed., Elsevier,  
Oxford, 557 pp., <https://doi.org/10.1029/eo081i050p00613-01>, 2015.

790 Brauer, A.: Annually Laminated Lake Sediments and Their Palaeoclimatic Relevance, 109–127,  
[https://doi.org/10.1007/978-3-662-10313-5\\_7](https://doi.org/10.1007/978-3-662-10313-5_7), 2004.

Brock, F., Higham, T., Ditchfield, P., and Ramsey, C. B.: Current Pretreatment Methods for AMS Radiocarbon  
Dating at the Oxford Radiocarbon Accelerator Unit (Orau), *Radiocarbon*, 52, 103–112,  
<https://doi.org/10.1017/S0033822200045069>, 2010.

795 Bronk Ramsey, C.: Radiocarbon Calibration and Analysis of Stratigraphy: The OxCal Program, *Radiocarbon*,  
37, 425–430, <https://doi.org/10.1017/s0033822200030903>, 1995.



- Bronk Ramsey, C.: Deposition models for chronological records, *Quat. Sci. Rev.*, 27, 42–60, <https://doi.org/10.1016/j.quascirev.2007.01.019>, 2008.
- Bronk Ramsey, C.: Dealing with Outliers and Offsets in Radiocarbon Dating, *Radiocarbon*, 51, 1023–1045, 800 <https://doi.org/10.1017/s0033822200034093>, 2009.
- Bronk Ramsey, C. and Lee, S.: Recent and Planned Developments of the Program OxCal, *Radiocarbon*, 55, 720–730, <https://doi.org/10.1017/s0033822200057878>, 2013.
- Cadena-Vela, S., Mazón, J.-N., and Fuster-Guilló, A.: Defining a Master Data Management Approach for Increasing Open Data Understandability, in: *Lecture Notes in Computer Science (including subseries Lecture Notes in Artificial Intelligence and Lecture Notes in Bioinformatics)*, vol. 11878 LNCS, 169–178, 805 [https://doi.org/10.1007/978-3-030-40907-4\\_17](https://doi.org/10.1007/978-3-030-40907-4_17), 2020.
- Chollet, F. and others: Keras: The Python Deep Learning library, <https://keras.io>, 2015.
- Ciarletta, D. J., Shawler, J. L., Tenebruso, C., Hein, C. J., and Lorenzo-Trueba, J.: Reconstructing Coastal Sediment Budgets From Beach- and Foredune-Ridge Morphology: A Coupled Field and Modeling Approach, *J. Geophys. Res. Earth Surf.*, 124, 1398–1416, <https://doi.org/10.1029/2018JF004908>, 2019. 810
- Clark, P. U., Dyke, A. S., Shakun, J. D., Carlson, A. E., Clark, J., Wohlfarth, B., Mitrovica, J. X., Hostetler, S. W., and McCabe, A. M.: The Last Glacial Maximum, *Science (80-. )*, 325, 710–714, <https://doi.org/10.1126/science.1172873>, 2009.
- Colman, S. M., Jones, G. A., Rubin, M., King, J. W., Peck, J. A., and Orem, W. H.: AMS radiocarbon analyses 815 from Lake Baikal, Siberia: Challenges of dating sediments from a large, oligotrophic lake, *Quat. Sci. Rev.*, 15, 669–684, [https://doi.org/10.1016/0277-3791\(96\)00027-3](https://doi.org/10.1016/0277-3791(96)00027-3), 1996.
- Courtin, J., Andreev, A. A., Raschke, E., Bala, S., Biskaborn, B. K., Liu, S., Zimmermann, H., Diekmann, B., Stouf-Leichsenring, K. R., Pestryakova, L. A., and Herzschuh, U.: Vegetation Changes in Southeastern Siberia During the Late Pleistocene and the Holocene, *Front. Ecol. Evol.*, 9, 233, 820 <https://doi.org/10.3389/fevo.2021.625096>, 2021.
- Dask Development Team: Dask: Library for dynamic task scheduling, <https://dask.org>, 2016.
- Dee, M. W., Palstra, S. W. L., Aerts-Bijma, A. T., Bleeker, M. O., De Bruijn, S., Ghebru, F., Jansen, H. G., Kuitens, M., Paul, D., Richie, R. R., Spriensma, J. J., Scifo, A., Van Zonneveld, D., Verstappen-Dumoulin, B. M. A. A., Wietzes-Land, P., and Meijer, H. A. J.: Radiocarbon Dating at Groningen: New and Updated 825 Chemical Pretreatment Procedures, *Radiocarbon*, 62, 63–74, <https://doi.org/10.1017/RDC.2019.101>, 2020.
- Diepenbroek, M., Grobe, H., Reinke, M., Schindler, U., Schlitzer, R., Sieger, R., and Wefer, G.: PANGAEA - an information system for environmental sciences, *Comput. Geosci.*, 28, 1201–1210, [https://doi.org/10.1016/S0098-3004\(02\)00039-0](https://doi.org/10.1016/S0098-3004(02)00039-0), 2002.
- Dolman, A. M.: hamstr: Hierarchical Accumulation Modelling with Stan and R, 830 <https://github.com/EarthSystemDiagnostics/hamstr>, 2021.
- Finkenbinder, M. S., Abbott, M. B., Finney, B. P., Stoner, J. S., and Dorfman, J. M.: A multi-proxy



- reconstruction of environmental change spanning the last 37,000 years from Burial Lake, Arctic Alaska, *Quat. Sci. Rev.*, 126, 227–241, <https://doi.org/10.1016/j.quascirev.2015.08.031>, 2015.
- 835 Fujiwara, H., Hajek, J., and Till, O.: Octave Forge - The “parallel” package, <https://octave.sourceforge.io/parallel/index.html>, 2021.
- Gaujoux, R.: doRNG: Generic Reproducible Parallel Backend for “foreach” Loops, <https://cran.r-project.org/package=doRNG>, 2020.
- 840 Goring, S., Williams, J. W., Blois, J. L., Jackson, S. T., Paciorek, C. J., Booth, R. K., Marlon, J. R., Blaauw, M., and Christen, J. A.: Deposition times in the northeastern United States during the Holocene: Establishing valid priors for Bayesian age models, *Quat. Sci. Rev.*, 48, 54–60, <https://doi.org/10.1016/j.quascirev.2012.05.019>, 2012.
- Hajdas, I., Ascough, P., Garnett, M. H., Fallon, S. J., Pearson, C. L., Quarta, G., Spalding, K. L., Yamaguchi, H., and Yoneda, M.: Radiocarbon dating, *Nat. Rev. Methods Prim.*, 1, 62, <https://doi.org/10.1038/s43586-021-00058-7>, 2021.
- 845 Harris, C. R., Millman, K. J., van der Walt, S. J., Gommers, R., Virtanen, P., Cournapeau, D., Wieser, E., Taylor, J., Berg, S., Smith, N. J., Kern, R., Picus, M., Hoyer, S., van Kerkwijk, M. H., Brett, M., Haldane, A., del Río, J. F., Wiebe, M., Peterson, P., Gérard-Marchant, P., Sheppard, K., Reddy, T., Weckesser, W., Abbasi, H., Gohlke, C., and Oliphant, T. E.: Array programming with NumPy, *Nature*, 585, 357–362, <https://doi.org/10.1038/s41586-020-2649-2>, 2020.
- 850 Haslett, J. and Parnell, A.: A simple monotone process with application to radiocarbon-dated depth chronologies, *J. R. Stat. Soc. Ser. C Appl. Stat.*, 57, 399–418, <https://doi.org/10.1111/j.1467-9876.2008.00623.x>, 2008.
- 855 Heaton, T. J., Köhler, P., Butzin, M., Bard, E., Reimer, R. W., Austin, W. E. N., Bronk Ramsey, C., Grootes, P. M., Hughen, K. A., Kromer, B., Reimer, P. J., Adkins, J., Burke, A., Cook, M. S., Olsen, J., and Skinner, L. C.: Marine20 - The Marine Radiocarbon Age Calibration Curve (0-55,000 cal BP), *Radiocarbon*, 62, 779–820, <https://doi.org/10.1017/RDC.2020.68>, 2020.
- Hollaway, M. J., Henrys, P. A., Killick, R., Leeson, A., and Watkins, J.: Evaluating the ability of numerical models to capture important shifts in environmental time series: A fuzzy change point approach, *Environ. Model. Softw.*, 139, 104993, <https://doi.org/10.1016/j.envsoft.2021.104993>, 2021.
- 860 Joblib Development Team: Joblib: running Python functions as pipeline jobs, <https://joblib.readthedocs.io/>, 2020.
- Killick, R. and Eckley, I. A.: changepoint: An R Package for Changepoint Analysis, *J. Stat. Softw.*, 58, 1–19, 2014.
- Killick, R., Haynes, K., and Eckley, I. A.: changepoint: An R package for changepoint analysis, <https://cran.r-project.org/package=changepoint>, 2016.
- 865 Kluyver, T., Ragan-Kelley, B., Pérez, F., Granger, B., Bussonnier, M., Frederic, J., Kelley, K., Hamrick, J., Grout, J., Corlay, S., Ivanov, P., Avila, D., Abdalla, S., and Willing, C.: Jupyter Notebooks -- a publishing



- format for reproducible computational workflows, in: *Positioning and Power in Academic Publishing: Players, Agents and Agendas*, 87–90, 2016.
- 870 Kokorowski, H. D., Anderson, P. M., Mock, C. J., and Lozhkin, A. V.: A re-evaluation and spatial analysis of evidence for a Younger Dryas climatic reversal in Beringia, *Quat. Sci. Rev.*, 27, 1710–1722, <https://doi.org/10.1016/j.quascirev.2008.06.010>, 2008.
- Kublitskiy, Y., Kulkova, M., Druzhinina, O., Subetto, D., Stančikaitė, M., Gedminienė, L., and Arslanov, K.: Geochemical Approach to the Reconstruction of Sedimentation Processes in Kamyshovoye Lake (SE Baltic, Russia) during the Late Glacial and Holocene, 10, 764, <https://doi.org/10.3390/min10090764>, 2020.
- 875 Lacourse, T. and Gajewski, K.: Current practices in building and reporting age-depth models, *Quat. Res.*, 96, 28–38, <https://doi.org/10.1017/qua.2020.47>, 2020.
- Lehnerr, I., St Louis, V. L., Sharp, M., Gardner, A. S., Smol, J. P., Schiff, S. L., Muir, D. C. G., Mortimer, C. A., Michelutti, N., Tarnocai, C., St Pierre, K. A., Emmerton, C. A., Wiklund, J. A., Köck, G., Lamoureux, S. F., and Talbot, C. H.: The world’s largest High Arctic lake responds rapidly to climate warming, *Nat. Commun.*, 9, 1–9, <https://doi.org/10.1038/s41467-018-03685-z>, 2018.
- 880 Lougheed, B. C. and Obrochta, S. P.: A Rapid, Deterministic Age-Depth Modeling Routine for Geological Sequences With Inherent Depth Uncertainty, *Paleoceanogr. Paleoclimatology*, 34, 122–133, <https://doi.org/10.1029/2018PA003457>, 2019.
- Lougheed, B. C., Van Der Lubbe, H. J. L., and Davies, G. R.:  $^{87}\text{Sr}/^{86}\text{Sr}$  as a quantitative geochemical proxy for  $^{14}\text{C}$  reservoir age in dynamic, brackish waters: Assessing applicability and quantifying uncertainties, *Geophys. Res. Lett.*, 43, 735–742, <https://doi.org/10.1002/2015GL066983>, 2016.
- 885 Lowe, J. J. and Walker, M.: *Reconstructing Quaternary Environments*, Routledge, <https://doi.org/10.4324/9781315797496>, 2014.
- Lozhkin, A., Anderson, P., Minyuk, P., Korzun, J., Brown, T., Pakhomov, A., Tsygankova, V., Burnatny, S., and Naumov, A.: Implications for conifer glacial refugia and postglacial climatic variation in western Beringia from lake sediments of the Upper Indigirka basin, 47, 938–953, <https://doi.org/10.1111/bor.12316>, 2018.
- 890 Martin, H., Schmid, C., Knitter, D., and Tietze, C.: oxcAAR: Interface to “OxCal” Radiocarbon Calibration, <https://cran.r-project.org/package=oxcAAR>, 2021.
- Microsoft Corporation and Weston, S.: doParallel: Foreach Parallel Adaptor for the “parallel” Package, <https://cran.r-project.org/package=doParallel>, 2020a.
- 895 Microsoft Corporation and Weston, S.: doSNOW: Foreach Parallel Adaptor for the “snow” Package, <https://cran.r-project.org/package=doSNOW>, 2020b.
- Microsoft Corporation and Weston, S.: foreach: Provides Foreach Looping Construct, <https://cran.r-project.org/package=foreach>, 2020c.
- 900 Müller, S., Tarasov, P. E., Andreev, A. A., and Diekmann, B.: Late Glacial to Holocene environments in the present-day coldest region of the Northern Hemisphere inferred from a pollen record of Lake Billyakh,



- Verkhoyansk Mts, NE Siberia, *Clim. Past*, 5, 73–84, <https://doi.org/10.5194/cp-5-73-2009>, 2009.
- Niephaus, F., Felgentreff, T., and Hirschfeld, R.: Towards polyglot adapters for the GraalVM, *ACM Int. Conf. Proceeding Ser.*, <https://doi.org/10.1145/3328433.3328458>, 2019.
- 905 Olsen, J., Ascough, P., Lougheed, B. C., and Rasmussen, P.: Radiocarbon Dating in Estuarine Environments, 141–170, [https://doi.org/10.1007/978-94-024-0990-1\\_7](https://doi.org/10.1007/978-94-024-0990-1_7), 2017.
- Parnell, A. C., Haslett, J., Allen, J. R. M., Buck, C. E., and Huntley, B.: A flexible approach to assessing synchronicity of past events using Bayesian reconstructions of sedimentation history, *Quat. Sci. Rev.*, 27, 1872–1885, <https://doi.org/10.1016/j.quascirev.2008.07.009>, 2008.
- 910 Parnell, A. C., Buck, C. E., and Doan, T. K.: A review of statistical chronology models for high-resolution, proxy-based Holocene palaeoenvironmental reconstruction, *Quat. Sci. Rev.*, 30, 2948–2960, <https://doi.org/10.1016/j.quascirev.2011.07.024>, 2011.
- Peng, B., Wang, G., Ma, J., Leong, M. C., Wakefield, C., Melott, J., Chiu, Y., Du, D., and Weinstein, J. N.: SoS notebook: An interactive multi-language data analysis environment, 34, 3768–3770, <https://doi.org/10.1093/bioinformatics/bty405>, 2018.
- 915 Pfalz, G.: GPawi/LANDO: LANDO public release v1.0, <https://doi.org/10.5281/ZENODO.5734334>, 2021.
- Pfalz, G., Diekmann, B., Freytag, J.-C., and Biskaborn, B. K.: Computers and Geosciences Harmonizing heterogeneous multi-proxy data from lake systems, *Comput. Geosci.*, 153, 11, <https://doi.org/10.1016/j.cageo.2021.104791>, 2021.
- 920 R Core Team: R: A Language and Environment for Statistical Computing, <https://www.r-project.org/>, 2021.
- Rasmussen, S. O., Andersen, K. K., Svensson, A. M., Steffensen, J. P., Vinther, B. M., Clausen, H. B., Siggaard-Andersen, M. L., Johnsen, S. J., Larsen, L. B., Dahl-Jensen, D., Bigler, M., Röthlisberger, R., Fischer, H., Goto-Azuma, K., Hansson, M. E., and Ruth, U.: A new Greenland ice core chronology for the last glacial termination, *J. Geophys. Res. Atmos.*, 111, 1–16, <https://doi.org/10.1029/2005JD006079>, 2006.
- 925 Reback, J., McKinney, W., jbrockmendel, Bossche, J. Van den, Augspurger, T., Cloud, P., gfyong, Sinhrks, Hawkins, S., Roeschke, M., Klein, A., Petersen, T., Tratner, J., She, C., Ayd, W., Naveh, S., Garcia, M., Schendel, J., Hayden, A., Saxton, D., Jancauskas, V., McMaster, A., Battiston, P., Seabold, S., patrick, Dong, K., chris-b1, h-vetinari, Hoyer, S., and Gorelli, M.: pandas-dev/pandas: Pandas 1.1.5, <https://doi.org/10.5281/ZENODO.4309786>, 2020.
- 930 Reimer, P. J., Austin, W. E. N., Bard, E., Bayliss, A., Blackwell, P. G., Bronk Ramsey, C., Butzin, M., Cheng, H., Edwards, R. L., Friedrich, M., Grootes, P. M., Guilderson, T. P., Hajdas, I., Heaton, T. J., Hogg, A. G., Hughen, K. A., Kromer, B., Manning, S. W., Muscheler, R., Palmer, J. G., Pearson, C., van der Plicht, J., Reimer, R. W., Richards, D. A., Scott, E. M., Southon, J. R., Turney, C. S. M., Wacker, L., Adolphi, F., Büntgen, U., Capano, M., Fahrni, S. M., Fogtmann-Schulz, A., Friedrich, R., Köhler, P., Kudsk, S., Miyake, F.,
- 935 Olsen, J., Reinig, F., Sakamoto, M., Sookdeo, A., and Talamo, S.: THE INTCAL20 NORTHERN HEMISPHERE RADIOCARBON AGE CALIBRATION CURVE (0–55 CAL kBP), *Radiocarbon*, 1–33, <https://doi.org/10.1017/RDC.2020.41>, 2020.



- Rethemeyer, J., Gierga, M., Heinze, S., Stolz, A., Wotte, A., Wischhöfer, P., Berg, S., Melchert, J., and Dewald, A.: Current Sample Preparation and Analytical Capabilities of the Radiocarbon Laboratory at CologneAMS, *Radiocarbon*, 61, 1449–1460, <https://doi.org/10.1017/rdc.2019.16>, 2019.
- 940
- Rudaya, N., Nazarova, L., Novenko, E., Andreev, A., Kalugin, I., Daryin, A., Babich, V., Li, H. C., and Shilov, P.: Quantitative reconstructions of mid- to late holocene climate and vegetation in the north-eastern alтай mountains recorded in lake teletskoye, *Glob. Planet. Change*, 141, 12–24, <https://doi.org/10.1016/j.gloplacha.2016.04.002>, 2016.
- 945
- Smol, J. P.: Arctic and Sub-Arctic shallow lakes in a multiple-stressor world: a paleoecological perspective, *Hydrobiologia*, 778, 253–272, <https://doi.org/10.1007/s10750-015-2543-3>, 2016.
- Strunk, A., Olsen, J., Sanei, H., Rudra, A., and Larsen, N. K.: Improving the reliability of bulk sediment radiocarbon dating, *Quat. Sci. Rev.*, 242, 106442, <https://doi.org/10.1016/j.quascirev.2020.106442>, 2020.
- Subetto, D. A., Nazarova, L. B., Pestryakova, L. A., Strykh, L. S., Andronikov, A. V., Biskaborn, B., Diekmann, B., Kuznetsov, D. D., Sapelko, T. V., and Grekov, I. M.: Paleolimnological studies in Russian northern Eurasia: A review, *Contemp. Probl. Ecol.*, 10, 327–335, <https://doi.org/10.1134/S1995425517040102>, 2017.
- 950
- Strykh, L., Subetto, D., and Nazarova, L.: Paleolimnological studies on the East European Plain and nearby regions: the PaleoLake Database, *J. Paleolimnol.*, 65, 369–375, <https://doi.org/10.1007/s10933-020-00172-8>, 2021.
- 955
- Telford, R. J., Heegaard, E., and Birks, H. J. B.: The intercept is a poor estimate of a calibrated radiocarbon age, 14, 296–298, <https://doi.org/10.1191/0959683604hl707fa>, 2004.
- Thanos, C.: Research Data Reusability: Conceptual Foundations, Barriers and Enabling Technologies, 5, 2, <https://doi.org/10.3390/publications5010002>, 2017.
- 960
- Trachsel, M. and Telford, R. J.: All age–depth models are wrong, but are getting better, 27, 860–869, <https://doi.org/10.1177/0959683616675939>, 2017.
- Vyse, S. A., Herzsuh, U., Andreev, A. A., Pestryakova, L. A., Diekmann, B., Armitage, S. J., and Biskaborn, B. K.: Geochemical and sedimentological responses of arctic glacial Lake Ilirney, chukotka (far east Russia) to palaeoenvironmental change since ~51.8 ka BP, *Quat. Sci. Rev.*, 247, 106607, <https://doi.org/10.1016/j.quascirev.2020.106607>, 2020.
- 965
- Vyse, S. A., Herzsuh, U., Pfalz, G., Pestryakova, L. A., Diekmann, B., Nowaczyk, N., and Biskaborn, B. K.: Sediment and carbon accumulation in a glacial lake in Chukotka (Arctic Siberia) during the late Pleistocene and Holocene: Combining hydroacoustic profiling and down-core analyses, *Biogeosciences Discuss.*, 2021, 1–40, <https://doi.org/10.5194/bg-2021-39>, 2021.
- 970
- Walker, M., Johnsen, S., Rasmussen, S. O., Steffensen, J.-P., Popp, T., Gibbard, P., Hoek, W., Lowe, J., Andrews, J., Björck, S., Cwynar, L., Hughen, K., Kershaw, P., Kromer, B., Litt, T., Lowe, D. J., Nakagawa, T., Newnham, R., and Schwander, J.: The Global Stratotype Section and Point (GSSP) for the base of the Holocene Series/Epoch (Quaternary System/Period) in the NGRIP ice core, 31, 264–267,



<https://doi.org/10.18814/epiiugs/2008/v3i12/016>, 2008.

- 975 Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T., Miller, E., Bache, S., Müller, K., Ooms, J., Robinson, D., Seidel, D., Spinu, V., Takahashi, K., Vaughan, D., Wilke, C., Woo, K., and Yutani, H.: Welcome to the Tidyverse, *J. Open Source Softw.*, 4, 1686, <https://doi.org/10.21105/joss.01686>, 2019.

- Wolfe, A. P.: A high-resolution late-glacial and early Holocene diatom record from Baffin Island, eastern Canadian Arctic, *Can. J. Earth Sci.*, 33, 928–937, <https://doi.org/10.1139/e96-070>, 1996.

Wright, A. J., Edwards, R. J., van de Plassche, O., Blaauw, M., Parnell, A. C., van der Borg, K., de Jong, A. F. M., Roe, H. M., Selby, K., and Black, S.: Reconstructing the accumulation history of a saltmarsh sediment core: Which age-depth model is best?, *Quat. Geochronol.*, 39, 35–67, <https://doi.org/10.1016/j.quageo.2017.02.004>, 2017.

- 985 Zaharia, M., Xin, R. S., Wendell, P., Das, T., Armbrust, M., Dave, A., Meng, X., Rosen, J., Venkataraman, S., Franklin, M. J., Ghodsi, A., Gonzalez, J., Shenker, S., and Stoica, I.: Apache spark: A unified engine for big data processing, *Commun. ACM*, 59, 56–65, <https://doi.org/10.1145/2934664>, 2016.

- Zander, P. D., Szidat, S., Kaufman, D. S., Żarczyński, M., Poraj-Górska, A. I., Boltshauser-Kaltenrieder, P., and Grosjean, M.: Miniature radiocarbon measurements (< 150 µg C) from sediments of Lake Żabińskie, Poland: effect of precision and dating density on age–depth models, 2, 63–79, <https://doi.org/10.5194/gchron-2-63-2020>, 2020.

Zolitschka, B., Francus, P., Ojala, A. E. K., and Schimmelmann, A.: Varves in lake sediments - a review, *Quat. Sci. Rev.*, 117, 1–41, <https://doi.org/10.1016/j.quascirev.2015.03.019>, 2015.