



$^{230}\text{Th}/\text{U}$ Isochron Dating of Cryogenic Cave Carbonates

Paul Töchterle¹, Simon D. Steidle¹, R. Lawrence Edwards², Yuri Dublyansky¹, Christoph Spötl¹, Xianglei Li², John Gunn³, Gina E. Moseley¹

¹Institute of Geology, University of Innsbruck, Innsbruck, 6020, Austria

5 ²Dept. Earth and Environmental Sciences, University of Minnesota, Minneapolis, 55455, MN/USA

³School of Geography, Earth and Environmental Sciences, University of Birmingham, Birmingham, B15 2TT, UK

Correspondence to: Paul Töchterle (paul.toechterle@uibk.ac.at)

Abstract. Cryogenic Cave Carbonates (CCCs) are a type of speleothem, typically dated with $^{230}\text{Th}/\text{U}$ disequilibrium methods, that provide evidence of palaeo-permafrost conditions. In the field, CCCs occur as distinct patches of millimetre- to centimetre-
10 sized loose crystals and crystal aggregates on the floor of cave chambers, lacking a framework to validate ages by stratigraphic order. Correction factors for initial ^{230}Th ($^{230}\text{Th}_0$) are often based on the bulk-earth derived ratio of initial $^{230}\text{Th}/^{232}\text{Th}$ activity ($(^{230}\text{Th}/^{232}\text{Th})_0$), which is a well-established approach when $^{230}\text{Th}_0$ is moderately low. For samples with elevated levels of $^{230}\text{Th}_0$, however, accuracy can be improved by constraining $(^{230}\text{Th}/^{232}\text{Th})_0$ independently. Here, we combine detailed morphological observations from three CCC patches found in Water Icicle Close Cavern in the Peak District (UK) with $^{230}\text{Th}/\text{U}$
15 analyses. We find that individual CCC crystals show a range of morphologies that arise from non-crystallographic branching in response to the chemical evolution of the freezing solution. Results of $^{230}\text{Th}/\text{U}$ dating indicate that samples with a large surface area relative to the sample volume are systematically more affected by contamination with $^{230}\text{Th}_0$. By fitting isochrons to these results, we test whether the CCCs in an individual patch formed during the same freezing event, and demonstrate that $(^{230}\text{Th}/^{232}\text{Th})_0$ can deviate substantially from the bulk-earth derived value and also vary between the different CCC patches.
20 Where CCCs display elevated $^{230}\text{Th}_0$, isochrons are a useful tool to constrain $(^{230}\text{Th}/^{232}\text{Th})_0$ and obtain ages with improved accuracy. Detritus absorbed to the crystal surface is shown to be the most likely source of $^{230}\text{Th}_0$. Our results suggest that some previously published CCC ages may merit re-assessment and we provide suggestions on how to approach future dating efforts.

1 Introduction

25 Cryogenic Cave Carbonates (CCCs) are speleothems that typically occur as an accumulation of loose crystals on the floor of cave chambers. Despite being described in earlier literature (Skřivánek, 1954), it was not until the early 2000s that their value as palaeoclimate archives became apparent (Žák et al., 2004). The widely accepted model of CCC formation involves precipitation in slowly freezing pools of water on top of cave ice bodies. This particular setting requires pre-existing, sizeable bodies of cave ice with cave air temperatures between approximately -1 and 0 °C (i.e. the ‘CCC window’, Spötl et al., 2021; Koltai et al., 2021), indicating that the cave was surrounded by permafrost at the time of CCC formation. A study of Winter



30 Wonderland Cave, USA (Munroe et al., 2021), validated this genetic model by observing very recent formation of CCCs in the exact setting as proposed by Žák et al. (2004).

As with common speleothems (e.g. stalagmites and flowstones), the formation age of CCCs is determined by $^{230}\text{Th}/\text{U}$ disequilibrium dating. The principle of this method relies on the large solubility difference between uranium and thorium in water (Gascoyne, 1992). Age precisions as low as 2‰ are routinely achieved using the latest analytical protocols (Cheng et al., 2013). However, the accuracy of dating results can be compromised if substantial amounts of thorium incorporated at the time of formation ($^{230}\text{Th}_0$) are not adequately corrected for (Dorale et al., 2004). In practice, ^{230}Th is identified and corrected by measuring the chemically equivalent ^{232}Th (expressed herein as measured ratio $(^{230}\text{Th}/^{232}\text{Th})_t$; parenthesis signify activity ratio hereafter), a long-lived isotope that is not part of the ^{238}U decay chain. Subsequently, $^{230}\text{Th}_0$ is subtracted by applying an initial $^{230}\text{Th}/^{232}\text{Th}$ activity ratio ($(^{230}\text{Th}/^{232}\text{Th})_0$). However, in nature, values for $(^{230}\text{Th}/^{232}\text{Th})$ vary by orders of magnitude (from <1 to >50, e.g., R. Scott, 1968; Moore and Sackett, 1964; Moore, 1981; Hubert et al., 2006; Hirose et al., 2012). Consequently, $(^{230}\text{Th}/^{232}\text{Th})_0$ needs to be accurately constrained if a sample's $^{230}\text{Th}_0$ is high enough to make the age equation sensitive to the correction term. There are four widely used approaches to derive $(^{230}\text{Th}/^{232}\text{Th})_0$: (1) by using the bulk-earth composition (e.g. Wedepohl, 1995); (2) by deriving it from stratigraphic constraints (e.g. Cheng et al., 2000; Hellstrom, 2006); (3) by using modern values measured at the field site, for example in groundwater; (4) by calculating values derived from isochrons (e.g., Lin et al., 1998; Henderson and Slowey, 2000; Cobb et al., 2003).

Since the early 2000's, more than 20 studies published $^{230}\text{Th}/\text{U}$ ages of CCCs, commonly from multiple samples, and interpreted them with regard to regional permafrost conditions (e.g., Richter and Riechelmann, 2008; Žák et al., 2009; Richter et al., 2010; Žák et al., 2012; Orvošová et al., 2014; Spötl and Cheng, 2014; Chaykovskiy et al., 2014; Richter et al., 2018; Koltai et al., 2021; Spötl et al., 2021). However, reported ages from a given patch of CCCs were rarely synchronous and sometimes spanned thousands of years and different climatic events (Luetscher et al., 2013; Dublyansky et al., 2018). This spread of ages led investigators to conclude that in those cases, CCCs must have formed during multiple freezing events. Since CCCs typically occur as accumulations of crystals with no internal stratigraphy, it is generally not possible to constrain $(^{230}\text{Th}/^{232}\text{Th})_0$ by stratigraphic order, validate individual ages or detect potential outliers and/or erroneous measurements (i.e., 'age reversals'), hence, in almost all cases, $^{230}\text{Th}_0$ was corrected by assuming a bulk-earth composition for $(^{230}\text{Th}/^{232}\text{Th})_0$.

55 In this study, we test the hypothesis that CCCs from a single patch formed coevally within achievable dating precision. We present a comprehensive dataset of three CCC patches from Water Icicle Close Cavern comprising 26 $^{230}\text{Th}/\text{U}$ ages, carbonate stable isotope data, and detailed morphological observations. For the first time, we apply an isochron approach to constrain $(^{230}\text{Th}/^{232}\text{Th})_0$ in CCCs in order to test whether an observed spread in ages is the result of multiple generations of CCC being present in a patch or arises from variable amounts of $^{230}\text{Th}_0$ across different samples.



60 **2 Study Site**

The samples used in this study were collected from Water Icicle Close Cavern (WICC), a limestone dissolution cave located in the Peak District (United Kingdom). The cave entrance (53.1781 °N, 1.7605 °W) is situated on a plateau at 338 m above sea level and is surrounded by a low-gradient hilly topography that has been incised by valleys, most of which are dry. The bedrock geology comprises a Lower Carboniferous limestone sequence, several hundred metres thick. In total, the cave
65 comprises over 1000 m of explored passages that formed along a horizontal level 30-40 m below the surface. The climate of the region is temperate with a mean annual air temperature of 8.9 °C and 1160 mm of annual precipitation (UK METoffice, 2020).

Previous research reported evidence of extensive former ice presence in the cave including speleothem damage, ice attachments and solifluction deposits (Gunn et al., 2020). Dating of several broken and re-sealed speleothems points towards at least one
70 period of cave glaciation between 87,000 and 83,000 years ago that can be related to the presence of permafrost. The area around Water Icicle Close Cavern was not glaciated during the Last Glacial Maximum but the British-Irish Ice Sheet terminated only around 30 km northwest of the cave at its maximum extent (Clark et al., 2012). Overall, 11 patches of CCC were documented in the cave, three of which were sampled in the course of this study.

3 Methods

75 **3.1 Sampling Strategy**

Three patches of CCCs were sampled according to different strategies depending on the respective grain size of the occurrence. If the mean crystal size was larger than approximately 5 mm, several individual crystals were selected while trying to obtain a representative subsample of all morphological varieties of the patch. For smaller grain sizes, a subsample was simply scooped up with a knife. With all sampling, damage to the cave was minimised such that the overall appearance of the CCC patches
80 was kept intact.

3.2 Carbonate Stable Isotope Analysis

While CCCs can be identified by their characteristic crystal morphology and field occurrence, the diagnostic property separating them from common speleothems is the stable isotope composition (Žák et al., 2018). Several morphological varieties of each sampled CCC patch were selected for stable isotope analysis. If grain size permitted, aliquots were taken from
85 the same specimen that was also used for petrographic analysis.

Samples were cleaned in an ultrasonic bath of de-ionised water before drilling between 0.10 and 0.80 mg of carbonate powder from random points on the surface with a carbide burr-tipped dental drill. The stable isotope composition of these carbonate powders was then determined using a Thermo Scientific Delta V Plus isotope ratio mass spectrometer coupled to a GasBench



II (Spötl and Vennemann, 2003). An analytical uncertainty
90 (1σ) of $\pm 0.06\%$ for $\delta^{13}\text{C}$ and $\pm 0.08\%$ for $\delta^{18}\text{O}$ applies to all
stable isotope data (Spötl, 2011) and values are reported
relative to the Vienna PeeDee Belemnite (VPDB) standard.

3.3 $^{230}\text{Th}/\text{U}$ analysis

As with stable isotope analyses, smaller crystals were rinsed
95 with deionised water for several minutes in an ultrasonic
cleaner. Visible surficial detritus was removed with a
toothbrush. Samples were then processed as a whole. For two
large samples of patch 3 (PT25 and PT26, see table 1) it was
possible to cut them in half using a diamond-coated wire saw,
100 polish and drill an aliquot from the inner part of the sample
of approximately 30 mg with a burr-tipped dental drill in a
laminar-flow hood.

Chemical preparation and mass spectrometry were performed at the Trace Metal Isotope Geochemistry Laboratory at the
University of Minnesota. Elemental extraction of U and Th was performed according to the procedures described in Edwards
105 et al. (1987). The resulting solutions were measured on a Thermo Fisher Neptune Plus multi-collector inductively coupled
plasma mass spectrometer in peak-jumping mode on a secondary electron multiplier (Shen et al., 2012).

Ages were calculated according to Edwards et al. (1987) with decay constants for uranium $\lambda_{238} = 1.55125 \cdot 10^{-6}$ (Jaffey et al.,
1971), $\lambda_{234} = 2.82206 \cdot 10^{-6}$ and thorium $\lambda_{230} = 9.1705 \cdot 10^{-6}$ (Cheng et al., 2013). Measurement uncertainties are given at the
 2σ level and ages are reported as years before 1950 CE (BCE).

110 Each uncorrected age was subsequently corrected for $^{230}\text{Th}_0$ based on a range of $(^{230}\text{Th}/^{232}\text{Th})_0$ values, starting from a bulk-
earth value of 0.82 ± 0.41 , representing secular equilibrium with a $(^{232}\text{Th}/^{238}\text{U})$ value of 3.8 (Wedepohl, 1995), to 15 times this
bulk-earth ratio (i.e.: $(^{230}\text{Th}/^{232}\text{Th})_0 = [0.82 \pm 0.41, \dots, 12.23 \pm 0.41]$, precision arbitrarily assumed as 50% of bulk-earth). The
resulting array of corrected ages and their respective uncertainties was converted to probability density functions (PDFs)
assuming a Gaussian distribution and summed per patch in order to compare with isochron dating results.

115 Isochrons were constructed by applying least squares regressions in a three-dimensional space of $(^{234}\text{U}/^{238}\text{U})$, $(^{232}\text{Th}/^{238}\text{U})$ and
 $(^{230}\text{Th}/^{238}\text{U})$ (Ludwig and Titterton, 1994). Isochron ages are calculated from the $(^{234}\text{U}/^{238}\text{U})$ and $(^{230}\text{Th}/^{238}\text{U})$ intercept \pm
the 95% confidence interval of the respective regression.




Figure 1: Selected CCC morphologies of patch 1. a) Rhombohedral calcite crystal with stepped surface, b) complex intergrown crystals composed of radial-fibrous calcite, c) SEM image of the surface of the sample depicted in a), d) SEM image of b).

3.4 Morphological Analysis

The cleaned samples were examined using a Keyence
120 VHX-6000 digital microscope. To investigate the
surface texture of CCCs, selected samples were gold-
coated and analysed using a JEOL JSM-6010LV
scanning electron microscope. Secondary electron
images were acquired using 15 kV accelerating
125 voltage.

4 Results and Discussion

4.1 Sample Characterisation

The sampled CCC patches in Water Icicle Close
Cavern are composed of calcite. Grain size varies
130 within each patch, but most specimens are
approximately 1-10 mm in diameter. One out of the
three sampled patches also contains larger crystals up
to 3 cm in diameter. Each occurrence comprises a variety of crystal morphologies, as is typical for CCCs (Žák et al., 2018).
Prevalent shapes include rhombohedra (fig. 1a) and aggregates of crystals (fig. 1b). Some specimens comprise elongated sheets
135 or needles branching radially from a common centre point and show an almost spherulitic shape. Secondary electron
microscopy (SEM) images reveal that rhombohedra-like morphologies have comparably smooth, stepped surfaces that mimic
the overall rhombohedral geometry (fig. 1c). More complex grains on the other hand have a highly irregular surface but
elongated rhombohedral shapes can still be recognised (fig. 1d).

The stable isotope composition of CCCs from patch 1 varies systematically with crystal morphology (fig. 2). Rhombohedral
140 crystals have $\delta^{18}\text{O}$ values from -12.4 to -14.1 ‰ and $\delta^{13}\text{C}$ values from -6.4 to -5.4 ‰. In comparison, complex grains show
lower $\delta^{18}\text{O}$ (-14.2 to -15.4 ‰) and higher $\delta^{13}\text{C}$ values (-5.1 to -4.5 ‰).

These observations can be interpreted in the context of the genetic model of CCCs (Žák et al., 2004). As water freezes, the
 $\delta^{18}\text{O}$ of residual water lowers progressively because ^{18}O is preferentially incorporated into the growing ice. Conversely, $\delta^{13}\text{C}$
increases due to preferential loss of ^{12}C as CO_2 degasses. This trend is well established for CCCs (Žák et al., 2008; Žák et al.,
145 2018) and is also observed from core to rim within single CCC grains (Töchterle, 2018).

The concept of a morphological evolution of CCCs has been formulated in previous studies (Žák et al., 2012), but analytical
evidence supporting it has been lacking. The observed correlation between crystal morphology and stable isotope composition




Figure 2: Stable isotope composition of selected crystals from patch 1. The data follow a trend with a negative slope, that is typical for CCC. The stable isotope composition changes systematically with the progression from rhombohedral to radially branching morphologies. All depicted samples are between 2 and 5 mm in diameter, but images were rescaled for better comparability.



(fig. 2) indicates that these CCCs formed during a single freezing event. Furthermore, complex aggregates apparently formed later in the freezing process than rhombohedral grains. During freezing, it is not only the stable isotope composition that changes, but also the overall concentration of solutes. A large body of literature confirms the effects of thermodynamic variables such as supersaturation, ionic strength and the concentration of foreign ions or organic compounds on calcite morphologies and fabrics (e.g. Frisia et al., 2000; Sunagawa, 2005; Sand et al., 2012; Frisia, 2015; Hong et al., 2016; Mercedes-Martín et al., 2021). The morphological variety of CCCs observed in this study can be viewed as an expression of changing crystallisation driving forces in response to progressive freezing. Two fundamental pathways can alter the shape of a grain. Firstly, multiple individual crystals can intergrow to form crystal aggregates. Secondly, the crystal fabric can be influenced by various degrees of non-crystallographic branching in response to fast growth and/or high supersaturation of the parent solution, resulting in radial growth with the endmember being a spherulite (Sunagawa, 2005; Shtukenberg et al., 2012; Yoreo et al., 2015). Crucial for this study, by building a more complex geometry or creating a rougher surface, both of these mechanisms potentially increase the relative surface area of a grain (i.e., the ratio between surface area and volume).

4.2 $^{230}\text{Th}/\text{U}$ Analyses

Overall, 26 $^{230}\text{Th}/\text{U}$ analyses were carried out on CCCs from three patches (table 1). From each patch, three samples of three morphological types were selected. All analyses yielded high concentrations of ^{238}U (1035 ± 1 to $3843 \pm 5 \text{ ng g}^{-1}$) and mostly low concentrations of detrital thorium (expressed as higher values of $(^{230}\text{Th}/^{232}\text{Th})_i$). In the following sections the results from each patch will be discussed individually. Samples are grouped and sorted into morphological types (A, B and C) according to the relative surface area and roughness on a visual basis.

4.2.1 Patch 1

The samples selected from patch 1 comprise rhombohedral single crystals (morphology A), branched aggregates (morphology B), and also a set of transitional forms that show a progression from lower to higher degrees of non-crystallographic branching




Figure 3: Uncorrected ages plotted against the respective $(^{230}\text{Th}/^{232}\text{Th})_t$ ratio. Colour represents morphology type. Also shown for each patch are corresponding isochron ages ($\pm 95\%$ confidence interval) and the summed normalised probability density functions (PDFs), calculated from an array of $(^{230}\text{Th}/^{232}\text{Th})_0$ values. Samples of type A in patch 2, which are much older, are not shown for better scaling.



180 (morphology C, suppl. fig. 1). In this patch, $(^{230}\text{Th}/^{232}\text{Th})_t$ varies by more than one order of magnitude (from 7.7 ± 0.2 to 150 ± 3) and correlates with the morphology type whereby the complex morphologies B and C show lower $(^{230}\text{Th}/^{232}\text{Th})_t$ values (table 1) indicative of higher $^{230}\text{Th}_0$. More complex morphologies therefore yielded older uncorrected ages. The
185 difference between these uncorrected ages is as large $\sim 6,200$ years (fig 3). Given the substantial spread in nominal ages and lack of constraints on $(^{230}\text{Th}/^{232}\text{Th})_0$, it is tempting to infer distinct formation events for the different morphological types. However, CCCs in this patch conform to a morphological evolution as described in section 4.1, implying that they
190 likely formed from the same parent solution during a single freezing event. It seems unlikely, however, that such an event lasted several thousand years, and that the cave temperature remained stable throughout this entire period. The question arises if it is possible to find a $(^{230}\text{Th}/^{232}\text{Th})_0$ ratio that would result in overlapping corrected ages, supporting the hypothesis that all
195 samples formed at the same time and thus reconcile dating results with morphological observations.

We calculated an array of corrected ages based on multiple possible $(^{230}\text{Th}/^{232}\text{Th})_0$ ratios (30 ratios between 1x and 15x bulk-earth) and derived a summed PDF (shaded area in fig 3), representing a cumulative likelihood
200 distribution of all corrected ages in this patch. Naturally, samples with low $(^{230}\text{Th}/^{232}\text{Th})_t$ are highly sensitive to $(^{230}\text{Th}/^{232}\text{Th})_0$ and yielded very wide distributions compared to samples with $(^{230}\text{Th}/^{232}\text{Th})_t > 100$.

Figure 4 shows the ^{238}U -normalised activities of ^{230}Th and ^{232}Th . These variables correlate with an R^2 of 0.98 (r1.1 in fig. 4, regression parameters
205 in suppl. table 1), meaning that 98 % of the observed variance in $(^{230}\text{Th}/^{238}\text{U})$ (which mainly determines the ages of these samples) can be explained by different levels of ^{232}Th . From the least-squares regression, it is possible to calculate an isochron age ($12,224 \pm 478$ BCE) and a $(^{230}\text{Th}/^{232}\text{Th})_0$ activity ratio of 2.7 ± 0.2 . This result partly overlaps the peak of the PDF. However, the linear regression model strongly depends on the oldest uncorrected age (PT33).
210 When disregarding PT33, the isochron age shifts to $11,865 \pm 172$ BCE (r1.2, $R^2 = 0.997$) with a $(^{230}\text{Th}/^{232}\text{Th})_0$ of 3.4 ± 0.1 , improving the agreement between the isochron and PDF ages. The isochron age of r1.2 also intercepts high- $(^{230}\text{Th}/^{232}\text{Th})_t$ samples notably better, which are less sensitive to differences in the $(^{230}\text{Th}/^{232}\text{Th})_0$ values. Excluding samples with very high




Figure 4: Isochron plots of ^{238}U -normalised thorium measurements of each CCC patch. Colour represents morphological type. The shaded area represents the 95% confidence interval of the regression slope and intercept. Regression lines are labelled with an identification number corresponding to supplementary table 1.



levels of $^{230}\text{Th}_0$ may be justified in cases where multiple sources of $^{230}\text{Th}_0$ were mixed with varying ratios, as will be discussed in section 4.3.

Table 1: Results of $^{230}\text{Th}/\text{U}$ analyses of CCC from Water Ice-Ice Close Cavern. Uncertainties are reported at the 2σ level.

Patch	Morph. Type	ID	^{238}U [ng g ⁻¹]	^{232}Th [pg g ⁻¹]	$(^{230}\text{Th}/^{232}\text{Th})_0$	$g^{234}\text{U}^*$	$(^{230}\text{Th}/^{238}\text{U})$	^{230}Th age [BCE] uncorrected	^{230}Th age [BCE] isochron corrected	$g^{234}\text{U}_{\text{initial}}^*$	isochron $(^{230}\text{Th}/^{232}\text{Th})_0$
1	A	PT27	1466 ±1	4733 ±95	150 ±3	489.6 ±2.0	0.1583 ±0.0003	12110 ±29	11850 ±30	506.3 ±2.0	
		PT28	1487 ±1	5510 ±110	133 ±3	488.7 ±1.5	0.1614 ±0.0002	12369 ±24	12071 ±27	505.8 ±1.5	
		PT29	1680 ±3	6514 ±131	125 ±3	492.8 ±2.5	0.1589 ±0.0004	12128 ±39	11817 ±41	509.6 ±2.6	
	B	PT30	1035 ±1	25456 ±510	23 ±0	491.6 ±1.7	0.1823 ±0.0006	14051 ±50	12093 ±102	508.8 ±1.7	
		PT31	1092 ±1	30804 ±617	20 ±0	492.7 ±1.9	0.1853 ±0.0004	14281 ±39	12040 ±109	509.8 ±2.0	3.4 ±0.1
		PT32	1072 ±2	52299 ±1049	13 ±0	490.9 ±2.4	0.2117 ±0.0008	16493 ±75	12646 ±188	508.9 ±2.5	
2	A	PT12	3843 ±5	2368 ±48	2176 ±44	497.5 ±2.1	0.4386 ±0.0007	36971 ±94	36868 ±94	552.1 ±2.3	
		PT13	2324 ±3	1843 ±37	1692 ±34	498.5 ±1.8	0.4390 ±0.0008	36983 ±96	36850 ±97	553.3 ±2.0	
		PT15	1707 ±2	981 ±20	1111 ±22	1039.3 ±2.7	0.2090 ±0.0004	11620 ±30	11548 ±31	1073.9 ±2.8	
	B	PT16	1703 ±2	1074 ±22	1012 ±20	1041.5 ±2.3	0.2089 ±0.0003	11599 ±24	11520 ±26	1076.2 ±2.3	
		PT17	1758 ±2	1884 ±38	596 ±12	1042.3 ±1.6	0.2090 ±0.0003	11602 ±19	11467 ±26	1076.8 ±1.7	7.3 ±0.7
		PT09	3010 ±4	6218 ±125	316 ±6	1038.9 ±2.2	0.2137 ±0.0004	11899 ±29	11640 ±46	1073.9 ±2.3	
3	A	PT18	2230 ±2	1588 ±32	1626 ±33	151.4 ±1.4	0.3790 ±0.0006	43058 ±103	42967 ±108	170.9 ±1.6	
		PT19	2271 ±2	1289 ±26	2042 ±41	150.2 ±1.3	0.3792 ±0.0005	43145 ±94	43072 ±98	169.6 ±1.4	
		PT20	2537 ±2	2889 ±58	1020 ±20	150.7 ±1.4	0.3800 ±0.0005	43223 ±98	43077 ±111	170.2 ±1.6	4.2 ±1.1
	B	PT21	1772 ±1	1463 ±29	1404 ±28	150.0 ±1.3	0.3794 ±0.0005	43168 ±89	43062 ±97	169.5 ±1.4	
		PT22	1688 ±1	2431 ±49	807 ±16	150.3 ±1.3	0.3801 ±0.0005	43260 ±95	43076 ±117	169.8 ±1.5	
		PT23	1889 ±3	2025 ±41	1084 ±22	150.5 ±1.7	0.3801 ±0.0009	43254 ±151	43117 ±159	170.0 ±1.9	
C	PT24	1112 ±2	10208 ±205	130 ±3	150.5 ±1.6	0.3893 ±0.0010	44530 ±162	43029 ±178	170.0 ±1.9		
	PT25	1484 ±3	92 ±5	18343 ±974	150.1 ±1.6	0.3736 ±0.0009	42366 ±147	42356 ±147	169.2 ±1.8	5.4 ±0.2	
	PT26	1547 ±3	168 ±5	10505 ±313	150.1 ±1.7	0.3729 ±0.0009	42269 ±144	42251 ±144	169.1 ±1.9		

* $g^{234}\text{U} = ((^{234}\text{U}/^{238}\text{U}) - 1) * 1000$.



215

4.2.2 Patch 2

Patch 2 also contains morphologies of variable relative surface area (suppl. fig. 2). Samples of morphology A are translucent with a brownish hue and tend to form smooth, euhedral crystal faces. Morphology B is milky-white and shows a high degree of non-crystallographic branching close to spherulitic growth. Morphology C also has a milky-white colour but is characterised by complex aggregates of blocky crystals with a large relative surface area per grain.

As for patch 1, the uncorrected ages vary for each morphological type (table 1). Types B and C cluster around 11,600 and 12,100 BCE respectively, while type A yielded much older ages around 37,000 BCE. Compared to patch 1, samples generally have higher $(^{230}\text{Th}/^{232}\text{Th})_t$ values (117 ± 2 to $5,994 \pm 121$) indicating relatively low $^{230}\text{Th}_0$. Thus, correction for $^{230}\text{Th}_0$ yields a much narrower PDF sum (fig. 3) and sub-sample ages for type A and B are identical within the uncertainty of each respective morphology. Again, a relationship between morphology and $(^{230}\text{Th}/^{232}\text{Th})_t$ can be observed, where samples with a smaller relative surface area (morphologies A and B) are less affected by $^{230}\text{Th}_0$ contamination than the most complex morphology (type C).

The dating results show that there are two distinct generations of CCC present in this patch (table 1, fig. 4). Types B and C plot along a common regression line ($r=2.1$, $R^2 = 0.98$), whereas type A samples are offset. For types B and C, the resulting isochron yields an age of $11,524 \pm 85$ BCE, indistinguishable from the PDF peak (fig. 3), and a $(^{230}\text{Th}/^{232}\text{Th})_0$ activity ratio of 7.3 ± 0.7 .

Note that the isochron-derived value for $(^{230}\text{Th}/^{232}\text{Th})_0$ is approximately 9 times higher than bulk-earth, meaning that the correction for $^{230}\text{Th}_0$ is proportionally larger than for patch 1, especially for samples with even moderately elevated $(^{230}\text{Th}/^{232}\text{Th})_t$. Previous U-series dating studies have proposed threshold values of $(^{230}\text{Th}/^{232}\text{Th})_t > 100$ or even > 300 (Li et al., 1989; Fensterer et al., 2010), above which detrital thorium contamination is deemed insignificant to the accuracy of dating results. Results from patch 2 indicate that in cases where the true value of $(^{230}\text{Th}/^{232}\text{Th})_0$ is multiple times higher than the bulk-earth estimate, threshold values may need to be adjusted to even higher levels (potentially as high as $(^{230}\text{Th}/^{232}\text{Th})_t > 500$ for relatively young samples) in order to achieve the same level of accuracy. We argue that such $(^{230}\text{Th}/^{232}\text{Th})_t$ limits should be considered as guides, and that each dataset should be evaluated on a case-by-case basis as acceptable limits will be affected by age, the relationship between analytical uncertainty and uncertainty in $^{230}\text{Th}_0$, and the age accuracy needed to address the question at hand. As presented here, isochrons can be a useful and reliable approach if high-quality, reliably dateable samples with high $(^{230}\text{Th}/^{232}\text{Th})_t$ are not available.

4.2.3 Patch 3

In patch 3, morphologies of type A and B consist of translucent calcite with a brownish hue (suppl. fig. 3, similar to type A of patch 2). Type A resembles rhombohedral single crystals with subtle non-crystallographic branching while Type B branches more strongly giving the grains a dumbbell-like shape. Type C is significantly larger (up to 3 cm in diameter) and shows a



transition from the brownish rhombohedral type to milky white calcite with a radial-fibrous texture indicative of high degrees of non-crystallographic branching.

250 The U and Th concentrations of type A and B are similar within an order of magnitude (table 1) with high $(^{230}\text{Th}/^{232}\text{Th})_t$ values $(807 \pm 16$ to $2,042 \pm 41)$ indicating very clean samples with minimal $^{230}\text{Th}_0$ contamination. The resulting uncorrected ages of these morphological types are identical within analytical uncertainty (fig. 3).

In contrast, morphological type C shows an extreme range of $(^{230}\text{Th}/^{232}\text{Th})_t$ values. The two large specimens, where aliquots from the core of the samples were analysed (suppl. fig. 3), have extremely high $(^{230}\text{Th}/^{232}\text{Th})_t$ (10,505 and 18,343 respectively), yielding a very narrow PDF sum that is indistinguishable from uncorrected ages. The third aliquot however, which was a piece
255 of surface material chipped off from a larger type C grain, yielded the lowest $(^{230}\text{Th}/^{232}\text{Th})_t$ (130 ± 3) of this patch and hence an uncorrected age ~ 2000 years older.

For this patch, the PDF sum clearly shows a bimodal distribution of corrected ages (fig. 3). An isochron regression can be fitted to the data when exempting morphology C ($r3.1$, $R^2 = 0.89$). The resulting isochron age of $43,065 \pm 139$ BCE ($(^{230}\text{Th}/^{232}\text{Th})_0 = 4.2 \pm 1.1$, fig. 4c) agrees with the older PDF peak corresponding to types A and B. The isochron age of type
260 C ($r3.2$, $42,297 \pm 109$ BCE) also confirms the results of the drilled aliquots and the corresponding $(^{230}\text{Th}/^{232}\text{Th})_0$ (5.4 ± 0.2) is similar to the value obtained from types A and B. However, it is noted that the regression is largely controlled by a single, highly inaccurate sample (PT24). Disregarding this sample, a very steep regression with $(^{230}\text{Th}/^{232}\text{Th})_0 = 16.8 \pm 3.5$ (i.e., approx. 21x bulk-earth) would be necessary to connect the two clusters of data. Consequently, it is reasonable to propose two distinct generations of CCC at $43,065 \pm 139$ BCE (type A and B) and $42,297 \pm 109$ BCE (type C). The isochron approach is thus able
265 to also resolve differing formation ages within a single patch.

4.3 Implications

In theory, $(^{230}\text{Th}/^{232}\text{Th})_0$ values derived from CCC isochrons represent the Th activity of the residual water from which the CCCs precipitated. All of the $(^{230}\text{Th}/^{232}\text{Th})_0$ values we found $(2.7 \pm 0.2$ to 7.2 ± 0.7 , suppl. table 1) are comparable to literature data on common speleothems (e.g., Beck et al., 2001; Hubert et al., 2006; Hoffmann et al., 2010; Carolin et al., 2013; Moseley
270 et al., 2013; Arienzo et al., 2015; Carolin et al., 2016). However, there is significant variability in $(^{230}\text{Th}/^{232}\text{Th})_0$ across the three patches. Other studies have reported a much lower in-cave variability with respect to $(^{230}\text{Th}/^{232}\text{Th})$ in sediments, speleothems and drip water (Olley et al., 1997; Kaufman et al., 1998). A possible explanation for this high variability can be entertained by considering the dynamic nature of CCC-forming pools. Munroe et al. (2021) reported that the residual pools holding CCCs in Winter Wonderland Cave show different degrees of discolouration of the residual water, indicating that
275 progressive freezing altered the physio-chemical properties of the residual water. One possible explanation could be that as colloids build-up within the residual water, the ratio between the two main sources of $^{230}\text{Th}_0$ (i.e., (i) colloids and (ii) silicates and iron (oxyhydr)oxides; (Short et al., 1988; Henderson and Slowey, 2000; Dorale et al., 2004) changes, thereby altering the Th isotopic composition. A non-linear behaviour of $(^{230}\text{Th}/^{232}\text{Th})_0$ could also explain why the isochron approach in patch 1



agrees better with the PDF sum, when sample PT33 is excluded from the regression. At this point however, it remains unclear
280 whether progressive freezing alters $(^{230}\text{Th}/^{232}\text{Th})_0$.

We have shown that for distinct patches of CCCs, an apparent spread in corrected ages can be an artefact arising from
underestimating $(^{230}\text{Th}/^{232}\text{Th})_0$ in samples with substantial $^{230}\text{Th}_0$. The data indicate that the relative surface area of a CCC
grain correlates with ^{232}Th content, likely because more detrital material adsorbs onto a larger surface. Thus, CCCs with smaller
surface areas will tend to require smaller corrections for $^{230}\text{Th}_0$ and yield more accurate ages. Further evidence supporting this
285 is provided by the drilled aliquots of core material from patch 3, which show the highest $(^{230}\text{Th}/^{232}\text{Th})_t$ of all analyses and
consequently yielded ages that are not sensitive to correction.

The dating issues that we describe in this study are ultimately tied to sample preparation. Grain sizes of CCCs are highly
variable and also seem to vary geographically. Very large specimens, sometimes over 10 cm in size, seem to be more prevalent
in Russian caves (e.g., Chaykovskiy and Kadebskaya, 2015 and Y. Dublyansky, unpublished data) while studies from central
290 Europe and the European Alps report mostly mm-sized crystals (Luetscher et al., 2013; Spötl and Cheng, 2014; Richter et al.,
2014; Richter et al., 2015; Pavuza and Spötl, 2017; Richter et al., 2017; Colucci et al., 2017; Žák et al., 2018; Richter et al.,
2019; Koltai et al., 2021; Richter et al., 2021; Spötl et al., 2021; Kluge et al., 2014). Drilling aliquots to obtain core material
from small grains is unfeasible in practise, which consequently demands a critical assessment method. If high-quality samples
with high $(^{230}\text{Th}/^{232}\text{Th})_t$ are not available, dating multiple morphological types (if present) and constructing isochrons may be
295 a key strategy in obtaining accurate and precise CCC chronologies.

5 Conclusions

- The morphological variability of CCCs in Water Icicle Close Cavern arises from aggregation and varying degrees of
non-crystallographic branching, which in turn reflect changes in the chemical environment during the CCC-forming
freezing process. Samples range from rhombohedral single crystals that form early in the process to highly branched
300 crystal aggregates that form when the driving forces of crystallisation increased at a later stage.
- Patches of CCCs likely formed during a single freezing event if a) the CCC morphologies show progressive non-
crystallographic branching, and b) aliquots define a highly correlated isochron.
- An apparent spread in uncorrected ages can arise from varying degrees of contamination by $^{230}\text{Th}_0$. To guarantee high
accuracy of samples with substantial $^{230}\text{Th}_0$, we recommend that ages from a single patch are further analysed using
305 an isochron approach from which both the age of formation and $(^{230}\text{Th}/^{232}\text{Th})_0$ can be calculated. Furthermore, age
offsets can be minimised by avoiding surface material where grain size permits.
- To reliably determine the age (range) of a patch of CCCs it is necessary to analyse multiple CCC individuals and – if
present – multiple morphologies.

These conclusions have implications for the application of CCCs in palaeoclimatology. If confirmed by studies of CCCs from
310 other caves, the accuracy of some previously published $^{230}\text{Th}/\text{U}$ results (corrected with nominal values of $(^{230}\text{Th}/^{232}\text{Th})_0$) should



be viewed with caution. Of particular concern are the ages reported for those studies for which reported values of ($^{230}\text{Th}/^{232}\text{Th}$)_t are low. As a consequence, the climatic interpretation of some studies that used CCCs to reconstruct past permafrost conditions may need to be re-assessed. This applies in particular to studies that analysed whole grains as opposed to aliquots of core material. In addition to increasing the number of aliquots per CCC patch, we recommend applying isochron dating to increase
315 the robustness of CCC-based palaeoclimate interpretations.

6 Code/Data availability

All data necessary to reproduce the results presented herein are provided in the text and supplementary material.

7 Author Contribution

PT performed analyses and prepared the manuscript. SDS and GEM performed additional calculations and contributed to
320 manuscript preparation. XL contributed with additional $^{230}\text{Th}/\text{U}$ analyses. RLE, CS, YD and JG were instrumental in the study design, directing analyses and preparing the manuscript.

8 Competing Interests

The authors declare no competing interests

9 Acknowledgements

325 The authors extend their gratitude to Alan Brentnall for cave access and guiding, Natural England for permitting fieldwork and sampling, and Peter Schroedl and Dylan Parmenter for assistance with $^{230}\text{Th}/\text{U}$ analyses at the University of Minnesota. We also thank Andy Freem for discovering the CCCs in Water Icicle Close Cavern as well as Robbie Shone for supporting field work, photography and discovering additional CCC samples.

10 References

- 330 Arienzo, M. M., Swart, P. K., Pourmand, A., Broad, K., Clement, A. C., Murphy, L. N., Vonhof, H. B., and Kakuk, B.: Bahamian speleothem reveals temperature decrease associated with Heinrich stadials, *Earth and Planetary Science Letters*, 430, 377–386, <https://doi.org/10.1016/j.epsl.2015.08.035>, 2015.
- Beck, J. W., Richards, D. A., Lawrence, R., Edwards, R. L., Silverman, B. W., Smart, P. L., Donahue, D. J., Hererra-Osterheld, S., Burr, G. S., Calsoyas, L., Timothy, A. J., Jull, and Biddulph, D.: Extremely Large Variations of
335 Atmospheric ^{14}C Concentration During the Last Glacial Period, *Science*, 292, 2453–2458, <https://doi.org/10.1126/science.1056649>, 2001.



- Carolin, S. A., Cobb, K. M., Lynch-Stieglitz, J., Moerman, J. W., Partin, J. W., Lejau, S., Malang, J., Clark, B., Tuen, A. A., and Adkins, J. F.: Northern Borneo stalagmite records reveal West Pacific hydroclimate across MIS 5 and 6, *Earth and Planetary Science Letters*, 439, 182–193, <https://doi.org/10.1016/j.epsl.2016.01.028>, 2016.
- 340 Carolin, S. A., Cobb, K. M., Adkins, J. F., Clark, B., Conroy, J. L., Lejau, S., Malang, J., and Tuen, A. A.: Varied Response of Western Pacific Hydrology to Climate Forcings over the Last Glacial Period, *Science*, 340, 1564–1566, <https://doi.org/10.1126/science.1233797>, 2013.
- Chaykovskiy, I. I. and Kadebskaya, O.: Morphology of cryogenic calcite from Rossiyskaya cave (Central Ural): Problems of mineralogy, petrography and metallogeny., *Scientific readings in memory of P.N. Chirvinsky (in Russian)*, 18, 102–112, 2015.
- 345 Chaykovskiy, I. I., Kadebskaya, O., and Žák, K.: Morphology, composition, age and origin of carbonate spherulites from caves of Western Urals, *Geochem. Int.*, 52, 336–346, <https://doi.org/10.1134/S0016702914020049>, 2014.
- Cheng, H., Edwards, R. L., Shen, C.-C., Polyak, V. J., Asmerom, Y., Woodhead, J., Hellstrom, J., Wang, Y., Kong, X., Spötl, C., Wang, X., and Calvin Alexander, E.: Improvements in ²³⁰Th dating, ²³⁰Th and ²³⁴U half-life values, and U–²³⁸Th isotopic measurements by multi-collector inductively coupled plasma mass spectrometry, *Earth and Planetary Science Letters*, 371–372, 82–91, <https://doi.org/10.1016/j.epsl.2013.04.006>, 2013.
- 350 Cheng, H., Adkins, J. F., Edwards, R. L., and Boyle, E. A.: U–Th dating of deep-sea corals, *Geochimica et Cosmochimica Acta*, 64, 2401–2416, [https://doi.org/10.1016/S0016-7037\(99\)00422-6](https://doi.org/10.1016/S0016-7037(99)00422-6), 2000.
- Clark, C. D., Hughes, A. L.C., Greenwood, S. L., Jordan, C., and Sejrup, H. P.: Pattern and timing of retreat of the last British-Irish Ice Sheet, *Quaternary Science Reviews*, 44, 112–146, <https://doi.org/10.1016/j.quascirev.2010.07.019>, 2012.
- 355 Cobb, K. M., Charles, C. D., Cheng, H., Kastner, M., and Edwards, R.L.: U/Th-dating living and young fossil corals from the central tropical Pacific, *Earth and Planetary Science Letters*, 210, 91–103, [https://doi.org/10.1016/S0012-821X\(03\)00138-9](https://doi.org/10.1016/S0012-821X(03)00138-9), 2003.
- 360 Colucci, R. R., Luetscher, M., Forte, E., Guglielmin, M., Lenaz, D., Princivale, F., and Vita, F.: First alpine evidence of in situ voarse cryogenic cave carbonates (CCC_coarse), *Geogr. Fis. Dinam. Quat.*, 40, 53–59, 2017.
- Dorale, J. A., Edwards, R. L., Alexander, E. C., Shen, C.-C., Richards, D. A., and Cheng, H.: Uranium-Series Dating of Speleothems: Current Techniques, Limits, & Applications, in: *Studies of Cave Sediments: Physical and Chemical Records of Paleoclimate*, edited by: I.D. Sasowsky and Mylroie, J. E., Kluwer Academic/Plenum Publishers, 177–197, https://doi.org/10.1007/978-1-4419-9118-8_10, 2004.
- 365 Dublyansky, Y., Moseley, G. E., Lyakhnitsky, Y., Cheng, H., Edwards, R. L., Scholz, D., Koltai, G., and Spötl, C.: Late Palaeolithic cave art and permafrost in the Southern Ural, *Scientific Reports*, 8, 12080, <https://doi.org/10.1038/s41598-018-30049-w>, 2018.
- Edwards, R. L., Chen, J. H., and Wasserburg, G. J.: ²³⁸U/²³⁴U/²³⁰Th/²³²Th systematics and the precise measurement of time over the past 500,000 years, *Earth and Planetary Science Letters*, 81, 175–192, [https://doi.org/10.1016/0012-821X\(87\)90154-3](https://doi.org/10.1016/0012-821X(87)90154-3), 1987.
- 370 Fensterer, C., Scholz, D., Hoffmann, D., Mangini, A., and Pajón, J. M.: ²³⁰Th/U-dating of a late Holocene low uranium speleothem from Cuba, *Geochimica et Cosmochimica Acta*, 9, 12015, <https://doi.org/10.1088/1755-1315/9/1/012015>, 2010.
- 375 Frisia, S., Borsato, A., Fairchild, I. J., and McDermott, F.: Calcite Fabrics, Growth Mechanisms, and Environments of Formation in Speleothems from the Italian Alps and Southwestern Ireland, *Journal of Sedimentary Research*, 70, 1183–1196, <https://doi.org/10.1306/022900701183>, 2000.
- Frisia, S.: Microstratigraphic logging of calcite fabrics in speleothems as tool for palaeoclimate studies, *IJS*, 44, 1–16, 2015.
- 380 Gascoyne, M.: Palaeoclimate determination from cave calcite deposits, *Quaternary Science Reviews*, 11, 609–632, [https://doi.org/10.1016/0277-3791\(92\)90074-I](https://doi.org/10.1016/0277-3791(92)90074-I), 1992.



- Gunn, J., Fairchild, I. J., Moseley, G. E., Töchterle, P., Ashley, K. E., Hellstrom, J., and Edwards, R. L.: Palaeoenvironments in the central White Peak District (Derbyshire, UK): evidence from Water Icicle Close Cavern, *Cave and Karst Science*, 47, 153–168, 2020.
- Hellstrom, J.: U–Th dating of speleothems with high initial ^{230}Th using stratigraphical constraint, *Quaternary Geochronology*, 1, 289–295, <https://doi.org/10.1016/j.quageo.2007.01.004>, 2006.
- Henderson, G. M. and Slowey, N. C.: Evidence from U–Th dating against Northern Hemisphere forcing of the penultimate deglaciation, *Nature*, 404, 61–66, <https://doi.org/10.1038/35003541>, 2000.
- Hirose, K., Kikawada, Y., and Igarashi, Y.: Temporal variation and provenance of thorium deposition observed at Tsukuba, Japan, *Journal of Environmental Radioactivity*, 108, 24–28, <https://doi.org/10.1016/j.jenvrad.2011.10.004>, 2012.
- Hoffmann, D. L., Beck, J. W., Richards, D. A., Smart, P. L., Singarayer, J. S., Ketchum, T., and Hawkesworth, C. J.: Towards radiocarbon calibration beyond 28ka using speleothems from the Bahamas, *Earth and Planetary Science Letters*, 289, 1–10, <https://doi.org/10.1016/j.epsl.2009.10.004>, 2010.
- Hong, M., Xu, J., and Teng, H. H.: Evolution of calcite growth morphology in the presence of magnesium: Implications for the dolomite problem, *Geochimica et Cosmochimica Acta*, 172, 55–64, <https://doi.org/10.1016/j.gca.2015.09.022>, 2016.
- Hubert, A., Bourdon, B., Pili, E., and Meynadier, L.: Transport of radionuclides in an unconfined chalk aquifer inferred from U-series disequilibria, *Geochimica et Cosmochimica Acta*, 70, 5437–5454, <https://doi.org/10.1016/j.gca.2006.08.008>, 2006.
- Jaffey, A. H., Flynn, K. F., Glendenin, L. E., Bentley, W. C., and Essling, A. M.: Precision Measurement of Half-Lives and Specific Activities of U^{235} and U^{238} , *Phys. Rev. C*, 4, 1889–1906, <https://doi.org/10.1103/PhysRevC.4.1889>, 1971.
- Kaufman, A., Wasserburg, G. J., Porcelli, D., Bar-Matthews, M., Ayalon, A., and Halicz, L.: U–Th isotope systematics from the Soreq cave, Israel and climatic correlations, *Earth and Planetary Science Letters*, 156, 141–155, [https://doi.org/10.1016/S0012-821X\(98\)00002-8](https://doi.org/10.1016/S0012-821X(98)00002-8), 1998.
- Kluge, T., Affek, H. P., Zhang, Y. G., Dublyansky, Y., Spötl, C., Immenhauser, A., and Richter, D. K.: Clumped isotope thermometry of cryogenic cave carbonates, *Geochimica et Cosmochimica Acta*, 126, 541–554, <https://doi.org/10.1016/j.gca.2013.11.011>, 2014.
- Koltai, G., Spötl, C., Jarosch, A. H., and Cheng, H.: Cryogenic cave carbonates in the Dolomites (northern Italy): insights into Younger Dryas cooling and seasonal precipitation, *Clim. Past*, 17, 775–789, <https://doi.org/10.5194/cp-17-775-2021>, 2021.
- Li, W.-X., Lundberg, J., Dickin, A. P., Ford, D. C., Schwarcz, H. P., McNutt, R., and Williams, D.: High-precision mass-spectrometric uranium-series dating of cave deposits and implications for palaeoclimate studies, *Nature*, 339, 534–536, <https://doi.org/10.1038/339534a0>, 1989.
- Lin, J. C., Broecker, W. S., Hemming, S. R., Hajdas, I., Anderson, R. F., Smith, G. I., Kelley, M., and Bonani, G.: A Reassessment of U–Th and ^{14}C Ages for Late-Glacial High-Frequency Hydrological Events at Searles Lake, California, *Quat. res.*, 49, 11–23, <https://doi.org/10.1006/qres.1997.1949>, 1998.
- Ludwig, K. R. and Titterton, D. M.: Calculation of $^{230}\text{Th}/\text{U}$ isochrons, ages, and errors, *Geochimica et Cosmochimica Acta*, 58, 5031–5042, [https://doi.org/10.1016/0016-7037\(94\)90229-1](https://doi.org/10.1016/0016-7037(94)90229-1), 1994.
- Luetscher, M., Borreguero, M., Moseley, G. E., Spötl, C., and Edwards, R. L.: Alpine permafrost thawing during the Medieval Warm Period identified from cryogenic cave carbonates, *The Cryosphere*, 7, 1073–1081, <https://doi.org/10.5194/tc-7-1073-2013>, 2013.
- Mercedes-Martín, R., Rogerson, M., Prior, T. J., Brasier, A. T., Reijmer, J. J.G., Billing, I., Matthews, A., Love, T., Lepley, S., and Pedley, M.: Towards a morphology diagram for terrestrial carbonates: Evaluating the impact of carbonate supersaturation and alginic acid in calcite precipitate morphology, *Geochimica et Cosmochimica Acta*, 306, 340–361, <https://doi.org/10.1016/j.gca.2021.04.010>, 2021.



- Moore, W. S.: The thorium isotope content of ocean water, *Earth and Planetary Science Letters*, 53, 419–426,
425 [https://doi.org/10.1016/0012-821X\(81\)90046-7](https://doi.org/10.1016/0012-821X(81)90046-7), 1981.
- Moore, W. S. and Sackett, W. M.: Uranium and thorium series inequilibrium in sea water, *J. Geophys. Res.*, 69, 5401–5405,
<https://doi.org/10.1029/JZ069i024p05401>, 1964.
- Moseley, G. E., Smart, P. L., Richards, D. A., and Hoffmann, D. L.: Speleothem constraints on marine isotope stage (MIS) 5
relative sea levels, Yucatan Peninsula, Mexico, *J. Quaternary Sci.*, 28, 293–300, <https://doi.org/10.1002/jqs.2613>, 2013.
- 430 Munroe, J., Kimble, K., Spötl, C., Marks, G. S., McGee, D., and Herron, D.: Cryogenic cave carbonate and implications for
thawing permafrost at Winter Wonderland Cave, Utah, USA, *Scientific Reports*, 11, 6430,
<https://doi.org/10.1038/s41598-021-85658-9>, 2021.
- Olley, J. M., Roberts, R. G., and Murray, A. S.: Disequilibria in the uranium decay series in sedimentary deposits at Allen's
cave, nullarbor plain, Australia: Implications for dose rate determinations, *Radiation Measurements*, 27, 433–443,
435 [https://doi.org/10.1016/S1350-4487\(96\)00114-X](https://doi.org/10.1016/S1350-4487(96)00114-X), 1997.
- Orvošová, M., Deininger, M., and Milovský, R.: Permafrost occurrence during the Last Permafrost Maximum in the Western
Carpathian Mountains of Slovakia as inferred from cryogenic cave carbonate, *Boreas*, 43, 750–758,
<https://doi.org/10.1111/bor.12042>, 2014.
- Pavuzá, R. and Spötl, C.: Neue Dazen zu Vorkommen und Entstehung kryogener Calcite in ostalpinen Höhlen, *Die Höhle*,
440 68, 2017.
- R. Scott, M.: Thorium and uranium concentrations and isotope ratios in river sediments, *Earth and Planetary Science Letters*,
4, 245–252, [https://doi.org/10.1016/0012-821X\(68\)90042-3](https://doi.org/10.1016/0012-821X(68)90042-3), 1968.
- Richter, D. K. and Riechelmann, D. F.: Late Pleistocene cryogenic calcite spherulites from the Malachitdom Cave (NE
Rhenish Slate Mountains, Germany): origin, unusual internal structure and stable C-O isotope composition, *IJS*, 37,
445 119–129, 2008.
- Richter, D. K., Mueller, M., Platte, A., and Scholz, D.: Erste weichselzeitliche Kryocalcite im Attendorn-Elsper Riffkomplex
(Frettermühler Wasserhöhle, Südwestfalen), *Geologie und Paläontologie in Westfalen*, 93, 1–16, 2019.
- Richter, D. K., Goll, K., Gerbe, W., Niederbayr, A., Platte, A., and Scholz, D.: Weichselzeitliche Kryocalcite als Hinweis für
Eisseen in der Hüttenbläuserschachthöhle (Iserlohn/NRW), *E&G Quaternary Science Journal*, 64, 67–81, 2015.
- 450 Richter, D. K., Harder, M., Niedermayr, A., and Scholz, D.: Zopfsinter in der Zoolithenhöhle: Erstfund kryogener Calcite in
der Fränkischen Alb, *Mitt. Verb. dt. Höhlen- u. Karstforscher*, 60, 36–41, 2014.
- Richter, D. K., Meissner, P., Immenhauser, A., Schulte, U., and Dorsten, I.: Cryogenic and non-cryogenic pool calcites
indicating permafrost and non-permafrost periods: A case study from the Herbstlabyrinth-Advent Cave system
(Germany), *The Cryosphere*, 4, 501–509, <https://doi.org/10.5194/tc-4-501-2010>, 2010.
- 455 Richter, D. K., Schudelski, A., Neuser, R. D., and Scholz, D.: Weichselzeitliche Umbrellacalcite aus der Höhle
"Malachitdom" (NE-Sauerland): vom Kaltwasser- zum Ausfrierstadium in Pools auf Eis, *Geologie und Paläontologie in
Westfalen*, 2021.
- Richter, D. K., Scholz, D., Jöns, N., Neuser, R. D., and Breitenbach, S. F.M.: Coarse-grained cryogenic aragonite as end-
member of mineral formation in dolomite caves, *Sedimentary Geology*, 376, 136–146,
460 <https://doi.org/10.1016/j.sedgeo.2018.08.006>, 2018.
- Richter, D. K., Knolle, F., Meyer, S., and Scholz, D.: Erste weichselzeitliche Kryocalcit-Vorkommen in Höhlen des
Iberg/Winterberg Riffkomplexes (Harz), *Mitt. Verb. dt. Höhlen- u. Karstforscher*, 63(2), 52–57, 2017.
- Sand, K. K., Rodriguez-Blanco, J. D., Makovicky, E., Benning, L. G., and Stipp, S. L. S.: Crystallization of CaCO₃ in
Water–Alcohol Mixtures: Spherulitic Growth, Polymorph Stabilization, and Morphology Change, *Crystal Growth &
465 Design*, 12, 842–853, <https://doi.org/10.1021/cg2012342>, 2012.



- Shen, C.-C., Wu, C.-C., Cheng, H., Edwards, R. L., Hsieh, Y.-T., Gallet, S., Chang, C.-C., Li, T.-Y., Lam, D. D., Kano, A., Hori, M., and Spötl, C.: High-precision and high-resolution carbonate ^{230}Th dating by MC-ICP-MS with SEM protocols, *Geochimica et Cosmochimica Acta*, 99, 71–86, <https://doi.org/10.1016/j.gca.2012.09.018>, 2012.
- 470 Short, S. A., Lowson, R. T., and Ellis, J.: $^{234}\text{U}/^{238}\text{U}$ and $^{230}\text{Th}/^{234}\text{U}$ activity ratios in the colloidal phases of aquifers in lateritic weathered zones, *Geochimica et Cosmochimica Acta*, 52, 2555–2563, [https://doi.org/10.1016/0016-7037\(88\)90026-9](https://doi.org/10.1016/0016-7037(88)90026-9), 1988.
- Shtukenberg, A. G., Punin, Y. O., Gunn, E., and Kahr, B.: Spherulites, *Chemical Reviews*, 112, 1805–1838, <https://doi.org/10.1021/cr200297f>, 2012.
- 475 Skřivánek, F.: *Jeskyně na Chlumu v Českém krasu (Caves at Chlum in the Bohemian Karst)*, in: *Československý kras 7 (in Czech)*, edited by: Skřivánek, F., 24–34, 1954.
- Spötl, C.: Long-term performance of the Gasbench isotope ratio mass spectrometry system for the stable isotope analysis of carbonate microsamples, *Rapid Comm. Mass Spec.*, 25, 1683–1685, <https://doi.org/10.1002/rcm.5037>, 2011.
- Spötl, C. and Cheng, H.: Holocene climate change, permafrost and cryogenic carbonate formation: Insights from a recently deglaciated, high-elevation cave in the Austrian Alps, *Clim. Past*, 10, 1349–1362, <https://doi.org/10.5194/cp-10-1349-2014>, 2014.
- 480 Spötl, C. and Vennemann, T. W.: Continuous-flow isotope ratio mass spectrometric analysis of carbonate minerals, *Rapid Comm. Mass Spec.*, 17, 1004–1006, <https://doi.org/10.1002/rcm.1010>, 2003.
- Spötl, C., Koltai, G., Jarosch, A. H., and Cheng, H.: Increased autumn and winter precipitation during the Last Glacial Maximum in the European Alps, *Nat Commun*, 12, 1839, <https://doi.org/10.1038/s41467-021-22090-7>, 2021.
- 485 Sunagawa, I.: *Crystals: Growth, morphology, and perfection*, Cambridge University Press, Cambridge, 295 pp., 2005.
- Töchterle, P.: *Cryogenic cave carbonates from the Ural Mountains (Russia)*, MSc Thesis, Institute of Geology, University of Innsbruck, Innsbruck, 2018.
- UK METoffice: Climatic Research Unit (CRU) Time-Series (TS) version 4.04 of high-resolution gridded data of month-by-month variation in climate (Jan. 1901- Dec. 2019)., available at: <https://catalogue.ceda.ac.uk/uuid/89e1e34ec3554dc98594a5732622bce9>, last access: 6 October 2020, 2020.
- 490 Wedepohl, H.: The composition of the continental crust, *Geochimica et Cosmochimica Acta*, 59, 1217–1232, [https://doi.org/10.1016/0016-7037\(95\)00038-2](https://doi.org/10.1016/0016-7037(95)00038-2), 1995.
- Yoreo, J. J. de, Gilbert, P. U. P. A., Sommerdijk, N. A. J. M., Penn, R. L., Whitlam, S., Joester, D., Zhang, H., Rimer, J. D., Navrotsky, A., Banfield, J. F., Wallace, A. F., Michel, F. M., Meldrum, F. C., Cölfen, H., and Dove, P. M.: CRYSTAL GROWTH. Crystallization by particle attachment in synthetic, biogenic, and geologic environments, *Science*, 349, aaa6760, <https://doi.org/10.1126/science.aaa6760>, 2015.
- 495 Žák, K., Onac, B. P., Kadebskaya, O., Filippi, M., Dublyansky, Y., and Luetscher, M.: Cryogenic Mineral Formation in Caves, in: *Ice caves*, edited by: Perşoiu, A. and Lauritzen, S.-E., Elsevier, Amsterdam, Oxford, Cambridge, Mass., 2018.
- 500 Žák, K., Richter, D. K., Filippi, M., Živor, R., Deininger, M., Mangini, A., and Scholz, D.: Coarsely crystalline cryogenic cave carbonate; a new archive to estimate the Last Glacial minimum permafrost depth in Central Europe, *Clim. Past*, 8, 1821–1837, <https://doi.org/10.5194/cp-8-1821-2012>, 2012.
- Žák, K., Hercman, H., Orvošová, M., and Jačková, I.: Cryogenic cave carbonates from the Cold Wind Cave, Nízke Tatry Mountains, Slovakia: Extending the age range of cryogenic cave carbonate formation to the Saalian, *IJS*, 38, 139–152, <https://doi.org/10.5038/1827-806X.38.2.5>, 2009.
- 505 Žák, K., Onac, B. P., and Perşoiu, A.: Cryogenic carbonates in cave environments: A review, *Quaternary International*, 187, 84–96, <https://doi.org/10.1016/j.quaint.2007.02.022>, 2008.
- Žák, K., Urban, J., Cílek, V., and Hercman, H.: Cryogenic cave calcite from several Central European caves: Age, carbon and oxygen isotopes and a genetic model, *Chemical Geology*, 206, 119–136, <https://doi.org/10.1016/j.chemgeo.2004.01.012>, 2004.