

In situ U-Pb dating of 4 billion year old carbonates in the martian meteorite Allan Hills 84001

Romain Tartèse¹, Ian C. Lyon¹

¹Department of Earth and Environmental Sciences, The University of Manchester, Manchester, M13 9PL, UK

5 Correspondence to: Romain Tartèse (romain.tartese@manchester.ac.uk)

Abstract. *In situ* carbonate U-Pb dating studies have proliferated dramatically in recent years. Almost all these studies have targeted relatively young terrestrial calcite up to Carboniferous in age. To assess the robustness of the carbonate U-Pb chronometer in deep-time, we carried out *in situ* U-Pb analyses in magnesite-ankerite-calcite carbonates in the martian meteorite Allan Hills (ALH) 84001. Carbonates in ALH 84001 formed at *ca.* 3.94 Ga, and there is little evidence that much
10 happened to this rock since then, making it an ideal sample to test the robustness of the U-Pb system in old carbonates. We obtained a concordant date of $3941 \pm 49/110$ Ma ($n = 14$, MSWD = 2.0), which is identical to the step-leaching Rb/Sr date determined previously. These results thus confirm that old carbonates are amenable to U-Pb dating in samples that have had a relatively simple history post-carbonate formation.

1 Introduction

15 Analytical developments in laser ablation – inductively coupled plasma mass spectrometry (LA-ICP-MS) over the last decade have driven important progress in *in situ* dating of carbonates, and in particular of calcite and occasionally dolomite, using the radioactive decay of uranium (U) into lead (Pb) (see the recent review by Roberts *et al.*, 2020, for example). Indeed, biogenic, diagenetic, and vein carbonates can typically incorporate up to *ca.* 10-20 $\mu\text{g/g}$ U, and up to *ca.* 100 $\mu\text{g/g}$ U in speleothems (e.g., Roberts *et al.*, 2020). Carbonates typically also incorporate initial Pb, meaning that multiple analyses on carbonate
20 samples often yield linear arrays in a Tera-Wasserburg inverse concordia diagram, providing information on both the $^{207}\text{Pb}/^{206}\text{Pb}$ composition of the initial Pb and the age of formation of the carbonates. Recent applications of carbonate U-Pb dating using LA-ICP-MS include constraining the timing of sedimentation, lithification, and diagenesis (e.g., Drost *et al.*, 2018; Godeau *et al.*, 2018; Mueller *et al.*, 2020; Brigaud *et al.*, 2021), faulting (e.g., Ring and Gerdes, 2016; Roberts and Walker, 2016; Goodfellow *et al.*, 2017; Nuriel *et al.*, 2017; 2019; Hansman *et al.*, 2018; Beaudoin *et al.*, 2018; Holdsworth *et al.*, 2019; Smeraglia *et al.*, 2019), aragonite to calcite conversion in ammonites (Li *et al.*, 2014), alteration of oceanic crust (Coogan *et al.*, 2016), veining, hydrothermalism, and mineralisation (Burisch *et al.*, 2017; 2018; Parrish *et al.*, 2018; Walter *et al.*, 2018; Bertok *et al.*, 2019; Drake *et al.*, 2019; 2020; MacDonald *et al.*, 2019), palaeoclimate reconstructions (Nicholson *et al.*, 2020), and hominin dispersion (Scardia *et al.*, 2019), for example.

30 All these *in situ* studies have targeted relatively young samples (younger than *ca.* 465 Ma), with two-third of dates younger
than 50 Ma, and all but ~~two~~ **three** younger than 300 Ma (Supplementary Table S1). This is consistent with the accepted idea
that carbonates are not very resistant to resetting of their U-Pb isotope systematics when thermal- and/or fluid-related alteration
events take place after their formation (e.g., Roberts *et al.*, 2020). However, a few studies have focused on dating older
35 carbonate samples, using wet chemistry to isolate Pb isotopes, and obtained Pb/Pb isochron dates ranging between *ca.* 1.60
and 2.84 Ga (Moorbath *et al.*, 1987; Bau *et al.*, 1999; Ray *et al.*, 2003; Sarangi *et al.*, 2004; Farey *et al.*, 2013). These Pb/Pb
dates have been interpreted as dating the deposition of these carbonates, suggesting that in some settings the Pb isotope
systematics of carbonates can remain undisturbed for billions of years. To further assess the robustness of the carbonate U-Pb
chronometer in deep time, we decided to attempt *in situ* LA-ICP-MS U-Pb dating of carbonates in the martian meteorite Allan
Hills 84001 (ALH 84001). The formation of these carbonates has been dated at 3.94 ± 0.02 Ga (2σ) using Rb-Sr analyses on
40 acid leachates via thermal ionisation mass spectrometry (Borg *et al.*, 1999; Beard *et al.*, 2013; date recalculated using a ^{87}Rb
decay constant of 1.3972×10^{-11} yr $^{-1}$; Villa *et al.*, 2015).

The meteorite ALH 84001 is an orthopyroxenite, a cumulate rock mostly comprising orthopyroxene, olivine, and chromite
(e.g., Mittlefehldt, 1994), which formed ~~ca. 4.1~~ **4.09 ± 0.03** Ga ago as suggested by Lu-Hf and Pb/Pb dating (Bouvier *et al.*,
45 2009; Lapen *et al.*, 2010). Carbonate-rich areas are irregularly scattered throughout ALH 84001, appearing as spherical or
hemispherical globules, discs along fractures, and irregular fillings in orthopyroxene (see review by Treiman (2021), and
references therein). The patches of carbonates show strong compositional zoning ranging from calcite-rich to magnesite-
siderite solid solution compositions (e.g., Corrigan and Harvey, 2004; Holland *et al.*, 2005). These carbonates likely formed
at low temperature (*ca.* 10-20 °C; Halevy *et al.*, 2011; del Real *et al.*, 2016) during fluid-rock interactions that were broadly
50 contemporaneous with the main shock event recorded in ALH 84001 (e.g., Treiman, 2021). **In order to test the robustness of
the U-Pb system in *ca.* 4 Gyr-old carbonates, it is essential to understand the geological history of those carbonates to assess
whether there have been affected by any hydrous or other alteration event since their formation.**

Dating carbonate-rich fractions in ALH 84001 using the Rb-Sr system yielded a precise formation age of 3.94 ± 0.02 Ga (2σ ;
55 **Borg *et al.*, 1999; Beard *et al.*, 2013), which is consistent with a less precise Pb/Pb isochron corresponding to a date of 4.045
 **± 0.090 Ga (2σ ; Borg *et al.*, 1999). The formation of these carbonates occurred at low temperature (<20 °C) and likely involved
mixing of two different water sources, one rich in Ca, the other rich in Fe (e.g., Halevy *et al.*, 2011, del Real *et al.*, 2016;
Bridges *et al.*, 2019, Treiman, 2021). These carbonate-forming fluids equilibrated with the atmosphere at the time, unlike
igneous minerals in the matrix (Shaheen *et al.*, 2015). A contemporaneous impact event raised the temperature of surrounding
60 plagioclase to *ca.* 1400°C, melting it to produce glass, and faulting carbonate globules (Mittlefehldt, 1994). Following this,
Treiman (2021) states “*there is little evidence that anything had happened to ALH84001 since 3.9Ga*”, until another impact
event at *ca.* 14 Ma, which caused the progenitor material that formed ALH84001 to be ejected from Mars (Eugster *et al.*,
1997). After *ca.* 14 Ma in space, ALH 84001 parent meteoroid fell to Antarctica *ca.* 13,000 years ago (Eugster *et al.*, 1997)****

and remained buried deep in the ice for millennia, only emerging at the surface of the Allan Hills ice field probably no more than 500 years ago (Krähenbühl *et al.*, 1998). Finally, there is extensive olivine and glass of plagioclase composition in ALH 84001 with no evidence of any alteration to clays or phyllosilicates. Modelling of Ar diffusion within the constituent minerals implies that the progenitor material for ALH 84001 was not subjected to temperatures >30 °C for any 'long duration' (Cassata *et al.*, 2010, Shuster and Weiss, 2005). All these lines of evidence indicate that the minerals in ALH 84001 were not exposed to hydrous fluids or temperatures >25-30 °C during the last 3.9 Ga (e.g., Treiman, 2021). Dating carbonate rich fractions using the Rb-Sr system yielded a precise formation age of 3.94 ± 0.02 Ga (2 σ ; Borg *et al.*, 1999; Beard *et al.*, 2013), which is consistent with a less precise Pb/Pb isochron corresponding to a date of 4.045 ± 0.090 Ga (2 σ ; Borg *et al.*, 1999). As stated by Treiman (2021), "there is little evidence that anything had happened to ALH 84001 since 3.9 Ga" and the time of carbonate formation until it was launched from Mars *ca.* 14 Ma ago (Eugster *et al.*, 1997), which is crucial with respect to the U-Pb system having remained closed since ALH 84001 carbonates formed.

75 2 Studied sample

The studied polished section (Fig. 1) was derived from a chip of the ALH84001,287 allocation from the NASA Ancient Mars Meteorite Program. The section contains patches of carbonates, associated with chromite, in between larger orthopyroxene grains (Fig. 1). The carbonates display the range of compositions typical for ALH 84001 carbonates, from Mg-rich magnesite to Ca-rich calcite, with intermediate Fe-rich ankerite areas (Fig. 1). The Mg- and Fe-rich carbonates seem to be part of broken rosettes, while Ca-rich carbonates appear associated with maskelynite (Fig. 1).

3 Laser Ablation – Inductively Coupled Plasma Mass Spectrometer analyses

U-Pb analyses were carried out at the University of Manchester using a Teledyne Photon Machines Analyte Excite+ 193 nm ArF excimer laser ablation system equipped with a HelEx II active 2-volume ablation cell, coupled to an Agilent 8900 triple quadrupole Inductively Coupled Plasma Mass Spectrometer (ICP-MS) using a signal-smoothing device (see Supplementary Table S2 for a summary of the analytical setup and data processing procedure).

The material ablated from target carbonates was carried to the ICP-MS by high purity He, which was mixed with Ar before injection into the plasma source. High purity N₂ was added to the He stream at a flow rate of 2 mL/min to enhance sensitivity. Tuning of the ICP-MS and mass calibration were performed at the start of the analytical session by optimising the ion signals during ablation of the NIST SRM 612 reference glass, while maintaining $^{238}\text{U}^{+}/^{232}\text{Th}^{+}$ close to unity and minimising the $^{232}\text{Th}^{16}\text{O}^{+}/^{232}\text{Th}^{+}$ ratio (*ca.* 0.3%). Glass and carbonates were ablated using a 25 μm laser beam size, a fluence of 4 J/cm², and a repetition rate of 5 Hz. Each analyses lasted 50 s and was preceded by 30 s counting time of the gas blank (background). The masses analysed and corresponding dwell times are reported in Supplementary Table S2.

95

The reference glass NIST614 (0.823 $\mu\text{g/g}$ U, 2.32 $\mu\text{g/g}$ Pb; Jochum *et al.*, 2011) was used to correct for $^{207}\text{Pb}/^{206}\text{Pb}$ fractionation, while mass bias correction of the measured $^{238}\text{U}/^{206}\text{Pb}$ ratios was carried out using repeated analyses of the reference calcite WC-1, which has a thermal ionisation mass spectrometry (TIMS) age of 254.4 ± 6.4 Ma (Roberts *et al.*, 2017). To ensure accuracy, the Duff Brown Tank (DBT) calcite (64.0 ± 0.7 Ma; Hill *et al.*, 2016) and AUG-B6 calcite (43.0 ± 1.0 Ma; Pagel *et al.*, 2018) were also analysed and used as secondary reference materials. Since there is no U/Pb magnesite-ankerite carbonate reference material, we have to rely on using a calcite reference material to correct for U/Pb fractionation in ALH 84001 carbonates.

Data processing was carried out using Iolite v4.5, using the NIST614 glass as primary reference material to remove instrument baseline contributions, mass bias of Pb isotopes, and downhole fractionation and instrumental drift of $^{206}\text{Pb}/^{238}\text{U}$ ratios (Paton *et al.*, 2011). The reproducibility obtained on NIST614 for $^{207}\text{Pb}/^{206}\text{Pb}$ ($\pm 1.7\%$, $n = 9$, 95% confidence level) and $^{206}\text{Pb}/^{238}\text{U}$ ($\pm 1.6\%$, $n = 9$, 95% confidence level) ratios were propagated by quadrature addition into each analysis $^{207}\text{Pb}/^{206}\text{Pb}$ and $^{206}\text{Pb}/^{238}\text{U}$ individual uncertainties. Repeated analyses of the NIST612 glass yielded an average $^{207}\text{Pb}/^{206}\text{Pb}$ ratio of 0.871 ± 0.044 ($n = 8$, 2 standard deviation), which is within error of its known $^{207}\text{Pb}/^{206}\text{Pb}$ ratio of 0.90745 ± 0.00004 (Baker *et al.*, 2004).

The data obtained for the reference calcite WC-1 were then plotted in a Tera-Wasserburg diagram using IsoplotR (Vermeesch, 2018), and yielded a lower intercept uncorrected date of 255.2 ± 5.9 Ma (95% confidence level, MSWD = 1.1, $n = 10$) for a discordia anchored at a the common $^{207}\text{Pb}/^{206}\text{Pb}$ ratio of 0.85 ± 0.04 determined by Roberts *et al.* (2017). To obtain the known intercept age of 254.4 ± 6.4 Ma for the WC-1 calcite, we applied a linear correction factor of 1.0031 to the measured $^{206}\text{Pb}/^{238}\text{U}$ ratios, which we also applied to all the samples analysed in the session, as is commonly done for carbonate U-Pb dating by LA-ICP-MS (e.g., Roberts *et al.*, 2017; Drost *et al.*, 2018; Kylander-Clark, 2020). All calculated dates are associated with two uncertainties, the first one including the random uncertainties for each analysis (internal uncertainties on measured $^{207}\text{Pb}/^{206}\text{Pb}$ and $^{206}\text{Pb}/^{238}\text{U}$ ratios and reproducibility on repeated NIST614 analyses), while systematic uncertainties (2.5% uncertainty on the WC-1 age, and 0.14% and 0.11% on the ^{235}U and ^{238}U decay constants, respectively; Jaffey *et al.*, 1971) are propagated by quadratic addition in the second one.

The data obtained on the DBT calcite yielded a lower intercept date of $64.9 \pm 2.2/2.8$ Ma (95% confidence level, MSWD = 2.9, $n = 12$) for a discordia anchored at a common $^{207}\text{Pb}/^{206}\text{Pb}$ ratio of 0.74 ± 0.02 calculated based on isotope dilution and multi-collector ICP-MS analyses (Hill *et al.*, 2016) (Supplementary Fig. S1). Because of its lower U abundance and younger age, the data obtained on the AUG-B6 calcite are less precise, yielding a concordia date of $40.8 \pm 2.0/2.2$ Ma (95% confidence level, MSWD concordance + equivalence = 2.2, $n = 8$) (Supplementary Fig. S1), which is identical to consistent with a $^{238}\text{U}/^{206}\text{Pb}$ weighted average date of $41.6 \pm 2.1/2.3$ Ma (95% confidence level, MSWD = 0.4, $n = 8$). All results are available in Supplementary Table S3.

130 4 Results

The carbonates analysed in ALH 84001 contain *ca.* 0.1-0.4 $\mu\text{g/g}$ U and 0.1-0.5 $\mu\text{g/g}$ Th (Table 1). When plotted in a Tera-Wasserburg $^{207}\text{Pb}/^{206}\text{Pb}$ vs. $^{238}\text{U}/^{206}\text{Pb}$ diagram, ALH 84001 carbonates yield a concordant date of $3941 \pm 49/110$ Ma ($n = 14$, MSWD = 2.0; Fig. 2A), which is identical to a weighted mean $^{207}\text{Pb}/^{206}\text{Pb}$ date of $3967 \pm 56/113$ Ma ($n = 14$, MSWD = 1.9; Fig. 2B). In detail, the Mg-rich and Ca-rich carbonate analyses yield concordia dates of 3890 ± 72 Ma (2σ , MSWD = 1.5, $n =$
135 8) and 3995 ± 69 Ma (2σ , MSWD = 2.4, $n = 6$), respectively; U-Pb dates for these two carbonate compositions are, therefore, indistinguishable when uncertainties are considered. Carbonate analyses plot on the concordia curve, indicating that they do not contain appreciable amount of common Pb. This is consistent with the measured ^{204}Pb intensities that are within error of 0 counts per second, although it is fair to point out that this observation is qualitative, considering the large uncertainties associated with the ^{202}Hg , $^{204}(\text{Hg}+\text{Pb})$, and calculated ^{204}Pb count rates (Table 1 and Supplementary Table S3). It is noteworthy
140 that ^{202}Hg intensities in ALH 84001 carbonates are about an order of magnitude higher than in the terrestrial carbonate standards (Supplementary Table S3). This could indicate that martian carbonates contain higher Hg abundances than terrestrial carbonates. Alternatively, we believe that this extra Hg likely originates from contamination of the ALH 84001 section by Au coating applied in the past for secondary ion mass spectrometry studies.

5 Discussion and implications

145 5.1 Comparison with previous ALH 84001 carbonate dating studies

In situ U-Pb dating of carbonates in ALH 84001 using LA-ICP-MS yields a concordia date of $3941 \pm 49/110$ Ma, which is identical to the carbonate step-leaching Rb-Sr isochron date of 3.94 ± 0.02 Ga (Borg *et al.*, 1999; Beard *et al.*, 2013) and the less precise Pb/Pb isochron date of 4.045 ± 0.090 Ga (Borg *et al.*, 1999). Before propagating the $\pm 2.5\%$ uncertainty associated with the age of the primary U-Pb reference carbonate WC-1, and the uncertainties associated with the ^{238}U and ^{235}U decay
150 constants, the carbonate U-Pb concordia date is accompanied associated with a fairly precise 2σ uncertainty of $\pm 1.2\%$, which increases to $\pm 2.8\%$ when all uncertainties are propagated. This suggests that *in situ* U-Pb dating of carbonates has the potential to yield precise dates, but also highlights the need to reduce uncertainties on reference materials. Our LA-ICP-MS results also indicate that using a calcite primary reference material for correcting U/Pb fractionation in Mg- and Fe-rich carbonate matrices, such as magnesite and ankerite, produce accurate dates (within the obtained uncertainties). A final point worth highlighting is
155 the fact that carbonates in ALH 84001 do not contain appreciable amount of common Pb, as indicated by their concordant U-Pb date of *ca.* 3.94 Ga and their ^{204}Pb intensities within error of 0 count per second. This is unusual, as in most examples in terrestrial systems carbonates, do contain common Pb incorporated during their crystallisation (e.g., Roberts *et al.*, 2020). This observation suggests that the fluids from which ALH 84001 carbonates formed contained very little Pb.

5.2 Robustness of the carbonate U-Pb chronometer and further applications

160 Our *in situ* LA-ICP-MS analyses confirm that carbonates in ALH 84001 formed *ca.* 3.94 Gyr-ago, and that the U-Pb
chronometer in these carbonates has remained closed to any disturbance event since they formed. This is consistent with the
suggestion that not much happened to ALH 84001 between 3.9 Ga and its launch from Mars 14 Myr-ago (Treiman, 2021), and
indicates that this latter event did not reset the carbonate U-Pb chronometer. From the evidence summarised in section 1,
165 Treiman (2021) concluded that minerals in ALH 84001, including the carbonates, have neither experienced temperatures in
excess of *ca.* 25°C nor exposure to any hydrous fluids since 3.9 Ga, leading to the conclusion that the climate of Mars has
remained globally cold and dry since then. Any aqueous events occurring on Mars due to a globally warmer and wetter climate
therefore took place before 3.9 Ga. Heating and aqueous alteration events due to meteoroid impacts could have occurred at
any time but these would have been strictly localised. Our U-Pb data back-up the view that the carbonates in ALH 84001 have
170 not experienced any alteration since they formed, supporting the conclusion that the progenitor material of ALH 84001 was
not exposed to hydrous fluids or high temperature events for the last 3.9 Ga, suggesting a globally cold and dry Mars since
then.

On the other hand, Carbonates carbonates in terrestrial Archean samples are probably not the best suited for U-Pb dating as
most Archean formations would have been heated up to at least low greenschist metamorphic conditions and/or been affected
175 by hydrothermal alteration, because the Earth is geologically active and harbours a complex hydrological cycle. On the other
hand, results of this study open up opportunities for dating old carbonates in samples that have had a relatively simple history
post-carbonate formation. For example, volatile-rich carbonaceous chondrites (e.g., CI and CM chondrites) typically contain
carbonates formed during fluid-rock interactions on their parent-asteroids *ca.* 4563-4561 Myr-ago (e.g., Lee *et al.*, 2014; Jilly-
Rehak *et al.*, 2017, and references therein), which is within 10 Myr of the formation of the first solids in the Solar System.
180 Bulk CI and CM chondrites contain ~10 ng/g U (e.g., Braukmüller *et al.*, 2018; Turner *et al.*, 2021), a significant proportion
of which being potentially hosted in labile phases such as carbonates (e.g., Burkhardt *et al.*, 2019; Turner *et al.*, 2021). As they
make up ~1-2 vol.% of CM chondrites (e.g., Lee *et al.*, 2014), these carbonates could host a few 100's ng/g U, which is similar
to the U abundance in the ALH 84001 carbonates analysed here. After this phase of early hydrothermal alteration during which
carbonates formed, it is thought that not much happens to these samples volatile-rich carbonaceous chondrites on their parent
185 asteroids until they end up on the Earth as meteorite fragments. Carbonates in carbonaceous chondrites could thus be prime
targets to further constrain the timing of hydrothermal alteration on volatile-rich asteroids using the U-Pb dating chronometer.

Data availability

Raw LA-ICP-MS data and metadata are provided in Supplementary Material.

Author contribution

190 ICL acquired and prepared the sample. RT designed the experiment, performed the analysis, and interpreted the results. RT prepared the manuscript with contributions from ICL.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgments

195 We acknowledge the provision of the sample of ALH84001 from the NASA Ancient Mars Meteorite Programme, and funding from STFC (#ST/P005225/1 to RT), and The University of Manchester and STFC (grant #ST/S002170/1) for funding the LA-ICP-MS facility. Many thanks to Nick Roberts for providing us with the WC-1 calcite, Benjamin Brigaud and Thomas Blaise for the AUG-B6 calcite, and Victor Polyak for the Duff Brown Tank calcite. We also thank Graham Edwards and an anonymous reviewer for their constructive suggestions.

200 References

- Baker, J., Peate, D., Waight, T., and Meyzen, C.: Pb isotopic analysis of standards and samples using a ^{207}Pb – ^{204}Pb double spike and thallium to correct for mass bias with a double-focusing MC-ICP-MS, *Chem. Geol.*, 211, 275-303, 2004.
- Bau, M., Romer, R. L., Lüders, V., and Beukes, N. J.: Pb, O, and C isotopes in silicified Mooidraai dolomite (Transvaal Supergroup, South Africa): implications for the composition of Paleoproterozoic seawater and ‘dating’ the increase of oxygen in the Precambrian atmosphere, *Earth Planet. Sci. Lett.*, 174, 43-57, 1999.
- 205 Beard, B. L., Ludois, J. M., Lapen, T. J., and Johnson, C. M.: Pre-4.0 billion year weathering on Mars constrained by Rb–Sr geochronology on meteorite ALH84001, *Earth Planet. Sci. Lett.*, 361, 173-182, 2013.
- Beaudoin, N., Lacombe, O., Roberts, N. M., and Koehn, D.: U-Pb dating of calcite veins reveals complex stress evolution and thrust sequence in the Bighorn Basin, Wyoming, USA, *Geology*, 46, 1015-1018, 2018.
- 210 Bertok, C., Barale, L., d’Atri, A., Martire, L., Piana, F., Rossetti, P., and Gerdes, A.: Unusual marbles in a non-metamorphic succession of the SW Alps (Valdieri, Italy) due to early Oligocene hydrothermal flow, *Int. J. Earth Sci.*, 108, 693-712, 2019.
- Borg, L. E., Connelly, J. N., Nyquist, L. E., Shih, C. Y., Wiesmann, H., and Reese, Y.: The age of the carbonates in martian meteorite ALH84001, *Science*, 286, 90-94, 1999.
- Bouvier, A., Blichert-Toft, J., and Albarede, F.: Martian meteorite chronology and the evolution of the interior of Mars, *Earth Planet. Sci. Lett.*, 280, 285-295, 2009.
- Braukmüller, N., Wombacher, F., Hezel, D. C., Escoube, R., and Münker, C.: The chemical composition of carbonaceous chondrites: Implications for volatile element depletion, complementarity and alteration, *Geochim. Cosmochim. Ac.*, 239, 17-48, 2018.
- 220 Bridges, J. C., Hicks, L. J., and Treiman, A. H.: Carbonates on Mars, in: *Volatiles in the Martian Crust*, edited by: Filiberto, J., and Schwenzer, S. P., Elsevier, Amsterdam, 89-118, 2019.

- Brigaud, B., Andrieu, S., Blaise, T., Haurine, F., and Barbarand, J.: Calcite uranium–lead geochronology applied to hardground lithification and sequence boundary dating, *Sedimentology*, 68, 168-195, 2021.
- Burisch, M., Gerdes, A., Walter, B. F., Neumann, U., Fettel, M., and Markl, G.: Methane and the origin of five-element veins: mineralogy, age, fluid inclusion chemistry and ore forming processes in the Odenwald, SW Germany, *Ore Geol. Rev.*, 81, 42-61, 2017.
- Burisch, M., Walter, B. F., Gerdes, A., Lanz, M., and Markl, G.: Late-stage anhydrite-gypsum-siderite-dolomite-calcite assemblages record the transition from a deep to a shallow hydrothermal system in the Schwarzwald mining district, SW Germany, *Geochim. Cosmochim. Ac.*, 223, 259-278, 2018.
- Burkhardt, C., Dauphas, N., Hans, U., Bourdon, B., and Kleine, T.: Elemental and isotopic variability in solar system materials by mixing and processing of primordial disk reservoirs, *Geochim. Cosmochim. Ac.*, 261, 145-170, 2019.
- Cassata, W. S., Shuster, D. L., Renne, P. R., and Weiss, B. P.: Evidence for shock heating and constraints on martian surface temperatures revealed by $^{40}\text{Ar}/^{39}\text{Ar}$ thermochronometry of martian meteorites, *Geochim. Cosmochim. Acta*, 74, 6900-6920, 2010.
- Coogan, L. A., Parrish, R. R., and Roberts, N. M.: Early hydrothermal carbon uptake by the upper oceanic crust: Insight from in situ U-Pb dating, *Geology*, 44, 147-150, 2016.
- Corrigan, C. M. and Harvey, R. P.: Multi-generational carbonate assemblages in martian meteorite Allan Hills 84001: implications for nucleation, growth, and alteration, *Meteorit. Planet. Sci.*, 39, 17-30, 2004.
- del Real, P. G., Maher, K., Kluge, T., Bird, D. K., Brown, G. E., and John, C. M.: Clumped-isotope thermometry of magnesium carbonates in ultramafic rocks, *Geochim. Cosmochim. Acta*, 193, 222-250, 2016.
- Drake, H., Roberts, N. M. W., Heim, C., Whitehouse, M. J., Siljeström, S., Kooijman, E., Broman, C., Ivarsson, M., and Åström, M. E.: Timing and origin of natural gas accumulation in the Siljan impact structure, Sweden, *Nat. Commun.*, 10, 1-14, 2019.
- Drake, H., Roberts, N. M., and Whitehouse, M. J.: Geochronology and Stable Isotope Analysis of Fracture-fill and Karst Mineralization Reveal Sub-Surface Paleo-Fluid Flow and Microbial Activity of the COSC-1 Borehole, Scandinavian Caledonides, *Geosciences*, 10, 56, <https://doi.org/10.3390/geosciences10020056>, 2020.
- Drost, K., Chew, D., Petrus, J. A., Scholze, F., Woodhead, J. D., Schneider, J. W., and Harper, D. A.: An Image Mapping Approach to U-Pb LA-ICP-MS Carbonate Dating, and Applications to Direct Dating of Carbonate Sedimentation, *Geochem. Geophys. Geosy.*, 19, 4631-4648, 2018.
- Eugster, O., Weigel, A., and Polnau, E.: Ejection times of martian meteorites, *Geochim. Cosmochim. Acta*, 61, 2749-2757, 1997.
- Fairey, B., Tsikos, H., Corfu, F., and Polteau, S.: U–Pb systematics in carbonates of the Postmasburg Group, Transvaal Supergroup, South Africa: primary versus metasomatic controls, *Precambrian Res.*, 231, 194-205, 2013.
- Godeau, N., Deschamps, P., Guihou, A., Leonide, P., Tendil, A., Gerdes, A., Hamelin, B., and Girard, J. P.: U-Pb dating of calcite cement and diagenetic history in microporous carbonate reservoirs: Case of the Urgonian Limestone, France, *Geology*, 46, 247-250, 2018.
- Goodfellow, B. W., Viola, G., Bingen, B., Nuriel, P., and Kylander-Clark, A. R.: Palaeocene faulting in SE Sweden from U–Pb dating of slickenfibres calcite, *Terra Nova*, 29, 321-328, 2017.

- Halevy, I., Fischer, W. W., and Eiler, J. M.: Carbonates in the martian meteorite Allan Hills 84001 formed at $18 \pm 4^\circ\text{C}$ in a near-surface aqueous environment, *Proc. Natl. Acad. Sci. USA*, 108, 16895-16899, 2011.
- 260 Hansman, R. J., Albert, R., Gerdes, A., and Ring, U.: Absolute ages of multiple generations of brittle structures by U-Pb dating of calcite, *Geology*, 46, 207-210, 2018.
- Hill, C. A., Polyak, V. J., Asmerom, Y., and Provencio, P.: Constraints on a Late Cretaceous uplift, denudation, and incision of the Grand Canyon region, southwestern Colorado Plateau, USA, from U-Pb dating of lacustrine limestone: U-Pb age of lacustrine limestone, *Tectonics*, 35, 896-906, 2016.
- 265 Holdsworth, R. E., McCaffrey, K. J. W., Dempsey, E., Roberts, N. M. W., Hardman, K., Morton, A., Feely, M., Hunt, J., Conway, A., and Robertson, A.: Natural fracture propping and earthquake-induced oil migration in fractured basement reservoirs, *Geology*, 47, 700-704, 2019.
- Holland, G., Saxton, J., Lyon, I., and Turner, G.: Negative $\delta^{18}\text{O}$ values in Allan Hills 84001 carbonate: possible evidence for water precipitation on Mars, *Geochim. Cosmochim. Acta*, 69, 1359-1370, 2005.
- 270 Jaffey A. H., Flynn, K. F., Glendenin, L. E., Bentley, W. C., and Essling, A.M.: Precision measurement of half-lives and specific activities of ^{235}U and ^{238}U , *Physical Review C*, 4, 1889-1906, 1971.
- Jilly-Rehak, C. E., Huss, G. R., and Nagashima, K.: ^{53}Mn - ^{53}Cr radiometric dating of secondary carbonates in CR chondrites: Timescales for parent body aqueous alteration, *Geochim. Cosmochim. Acta*, 201, 224-244, 2017.
- Jochum, K. P., Weis, U., Stoll, B., Kuzmin, D., Yang, Q., Raczek, I., Jacob, D. E., Stracke, A., Birbaum, K., Frick, D. A.,
 275 Gunther, D., and Enzweiler, J.: Determination of reference values for NIST SRM 610–617 glasses following ISO guidelines, *Geostand. Geoanal. Res.*, 35, 397-429, 2011.
- Krähenbühl, U., Noll, K., Dobeli, M., Grambole, D., Herrmann, F., and Tobler, L.: Exposure of Allan Hills 84001 and other achondrites on the Antarctic Ice, *Meteorit. Planet. Sci.*, 33, 665-670, 1998.
- Kylander-Clark, A. R. C.: Expanding the limits of laser-ablation U–Pb calcite geochronology, *Geochronology*, 2, 343-354,
 280 2020.
- Lapen, T., Richter, M., Brandon, A., Debaille, V., Beard, B., Shafer, J., and Peslier, A.: A younger age for ALH84001 and its geochemical link to shergottite sources in Mars, *Science*, 328, 347-351, 2010.
- Lee, M. R., Lindgren, P., and Sofo, M. R.: Aragonite, breunnerite, calcite and dolomite in the CM carbonaceous chondrites: High fidelity recorders of progressive parent body aqueous alteration, *Geochim. Cosmochim. Ac.*, 144, 126-156, 2014.
- 285 Li, Q., Parrish, R. R., Horstwood, M. S. A., and McArthur, J. M.: U–Pb dating of cements in Mesozoic ammonites, *Chem. Geol.*, 376, 76-83, 2014.
- MacDonald, J. M., Faithfull, J. W., Roberts, N. M. W., Davies, A. J., Holdsworth, C. M., Newton, M., Williamson, S., Boyce, A., and John, C. M.: Clumped-isotope palaeothermometry and LA-ICP-MS U–Pb dating of lava-pile hydrothermal calcite veins, *Contrib. Mineral. Petr.*, 174, 63, <https://doi.org/10.1007/s00410-019-1599-x>, 2019.
- 290 Mittlefehldt, D. W.: ALH84001, a cumulate orthopyroxenite member of the Martian meteorite clan. *Meteoritics*, 29, 214-221, 1994.
- Moorbath, S., Taylor, P. N., Orpen, J. L., Treloar, P., and Wilson, J. F.: First direct radiometric dating of Archaean stromatolitic limestone, *Nature*, 326, 865-867, 1987.

- 295 Mueller, M., Igbokwe, O. A., Walter, B., Pederson, C. L., Riechelmann, S., Richter, D. K., Albert, R., Gerdes, A., Buhl, D., Neuser, R. D., Bertotti, G., and Immenhauser, A.: Testing the preservation potential of early diagenetic dolomites as geochemical archives, *Sedimentology*, 67, 849-881, 2020.
- Nicholson, S. L., Pike, A. W., Hosfield, R., Roberts, N. M. W., Sahy, D., Woodhead, J., Cheng, H., Edwards, R. L., Affolter, S., Leuenberger, M., and Burns, S. J.: Pluvial periods in Southern Arabia over the last 1.1 million-years, *Quaternary Sci. Rev.*, 229, 106112, <https://doi.org/10.1016/j.quascirev.2019.106112>, 2020.
- 300 Nuriel, P., Craddock, J., Kylander-Clark, A. R., Uysal, T., Karabacak, V., Dirik, R. K., Hacker, B. R., and Weinberger, R.: Reactivation history of the North Anatolian fault zone based on calcite age-strain analyses, *Geology*, 47, 465-469, 2019.
- Nuriel, P., Weinberger, R., Kylander-Clark, A. R. C., Hacker, B. R., and Craddock, J. P.: The onset of the Dead Sea transform based on calcite age-strain analyses, *Geology*, 45, 587-590, 2017.
- 305 Pagel, M., Bonifacie, M., Schneider, D. A., Gautheron, C., Brigaud, B., Calmels, D., Cros, A., Saint-Bezar, B., Landrein, P., Sutcliffe, C., Davis, D., and Chaduteau, C.: Improving paleohydrological and diagenetic reconstructions in calcite veins and breccia of a sedimentary basin by combining Δ_{47} temperature, $\delta^{18}\text{O}_{\text{water}}$ and U-Pb age, *Chem. Geol.*, 481, 1-17, 2018.
- Parrish, R. R., Parrish, C. M., and Lasalle, S.: Vein calcite dating reveals Pyrenean orogen as cause of Paleogene deformation in southern England, *J. Geol. Soc.*, 175, 425-442, 2018.
- 310 Paton C., Hellstrom J., Paul B., Woodhead J., and Hergt J.: Iolite: Freeware for the visualisation and processing of mass spectrometric data, *J. Anal. Atom. Spectro.*, 26, 2508-2518, 2011.
- Ray, J. S., Veizer, J., and Davis, W. J.: C, O, Sr and Pb isotope systematics of carbonate sequences of the Vindhyan Supergroup, India: age, diagenesis, correlations and implications for global events, *Precambrian Res.*, 121, 103-140, 2003.
- Ring, U. and Gerdes, A.: Kinematics of the Alpenrhein-Bodensee graben system in the Central Alps: Oligocene/Miocene transtension due to formation of the Western Alps arc, *Tectonics*, 35, 1367-1391, 2016.
- 315 Roberts, N. M. W. and Walker, R. J.: U-Pb geochronology of calcite-mineralized faults: Absolute timing of rift-related fault events on the northeast Atlantic margin, *Geology*, 44, 531-534, 2016.
- 320 Roberts, N. M. W., Drost, K., Horstwood, M. S. A., Condon, D. J., Chew, D., Drake, H., Milodowski, A. E., McLean, N. M., Smye, A. J., Walker, R. J., Haslam, R., Hodson, K., Imber, J., Beaudoin, N., and Lee, J. K.: Laser ablation inductively coupled plasma mass spectrometry (LA-ICP-MS) U-Pb carbonate geochronology: strategies, progress, and limitations, *Geochronology*, 2, 33-61, 2020.
- Roberts, N. M. W., Rasbury, E. T., Parrish, R. R., Smith, C. J., Horstwood, M. S. A., and Condon, D. J.: A calcite reference material for LA-ICP-MS U-Pb geochronology, *Geochem. Geophys. Geosy.*, 18, 2807-2814, 2017.
- 325 Sarangi, S., Gopalan, K., and Kumar, S.: Pb-Pb age of earliest megascopic, eukaryotic alga bearing Rohtas Formation, Vindhyan Supergroup, India: implications for Precambrian atmospheric oxygen evolution, *Precambrian Res.*, 132, 107-121, 2004.
- Scardia, G., Parenti, F., Miggins, D. P., Gerdes, A., Araujo, A. G., and Neves, W. A.: Chronologic constraints on hominin dispersal outside Africa since 2.48 Ma from the Zarqa Valley, Jordan, *Quaternary Sci. Rev.*, 219, 1-19, 2019.
- Shaheen, R., Niles, P. B., Chong, K., Corrigan, C. M., and Thiemens, M. H.: Carbonate formation events in ALH 84001 trace the evolution of the martian atmosphere, *Proc. Natl. Acad. Sci. USA*, 112, 336-341, 2015.
- 330 Shuster, D. L., and Weiss, B. P.: Martian surface paleotemperatures from thermochronology of meteorites, *Science*, 309, 594-600, 2005.

- Smeraglia, L., Aldega, L., Billi, A., Carminati, E., Di Fiore, F., Gerdes, A., Albert, R., Rossetti, F., and Vignaroli, G.: Development of an intra-wedge tectonic mélangé by out-of-sequence thrusting, buttressing, and intraformational rheological contrast, Mt. Massico ridge, Apennines, Italy, *Tectonics*, 38, 1223-1249, 2019.
- 335 Treiman, A. H.: Uninhabitable and Potentially Habitable Environments on Mars: Evidence from Meteorite ALH 84001, *Astrobiology*, 21, 940-953, 2021.
- Turner, S., McGee, L., Humayun, M., Creech, J., and Zanda, B.: Carbonaceous chondrite meteorites experienced fluid flow within the past million years, *Science*, 371, 164-167, 2021.
- Vermeesch, P.: IsoplotR: a free and open toolbox for geochronology, *Geosci. Frontiers*, 9, 1479-1493, 2018.
- 340 Villa, I. M., De Bievre, P., Holden, N. E., and Renne, P. R.: IUPAC–IUGS recommendation on the half-life of ^{87}Rb , *Geochim. Cosmochim. Acta*, 164, 382-385, 2015.
- Walter, B. F., Gerdes, A., Kleinhanns, I. C., Dunkl, I., von Eynatten, H., Kreissl, S., and Markl, G.: The connection between hydrothermal fluids, mineralization, tectonics and magmatism in a continental rift setting: Fluorite Sm-Nd and hematite and carbonates U-Pb geochronology from the Rhinegraben in SW Germany, *Geochim. Cosmochim. Ac.*, 240, 11-42, 2018.




Figure 1: Backscattered electron image (A) and composite X-ray map (B) of the target carbonate patches in ALH 84001. Mineral abbreviations are ank = ankerite, cal = calcite, chr = chromite, mgs = magnesite, msk = maskelynite, opx = orthopyroxene.




Figure 2: Tera-Wasserburg $^{207}\text{Pb}/^{206}\text{Pb}$ vs. $^{238}\text{U}/^{206}\text{Pb}$ diagram (A) and $^{207}\text{Pb}/^{206}\text{Pb}$ dates (B) obtained on ALH 84001 carbonates. Ellipses (A) and error bars (B) correspond to 2σ uncertainties standard errors. The bold black ellipse in (A) corresponds to the calculated concordia date and associated uncertainty.

355 **Table 1: LA-ICP-MS results for ALH 84001 carbonates.**

Analysis #	²⁰⁴ Pb (cps) ¹	±	Pb (µg/g)	U (µg/g)	Th (µg/g)	Th/ U	Ratios ± 2σ (%)				conc. (%) ²	Dates ± 2σ (Ma)			
							²³⁸ U/ ²⁰⁶ Pb	±	²⁰⁷ Pb/ ²⁰⁶ Pb	±		²⁰⁷ Pb/ ²⁰⁶ Pb	±	²⁰⁶ Pb / ²³⁸ U	±
MgRich_1	100	110	0.11	0.26	0.20	0.76	1.06	29.2	0.4037	28.3	109	3923	370	4270	853
MgRich_3	-123	82	0.12	0.14	0.08	0.57	1.17	23.9	0.5356	25.2	92	4342	327	3976	671
MgRich_4	10	50	0.13	0.19	0.12	0.64	1.23	18.1	0.4295	24.9	96	4016	329	3835	504
MgRich_5	-4	42	0.20	0.41	0.50	1.21	1.29	8.8	0.3570	12.5	99	3737	178	3698	244
MgRich_6	27	49	0.10	0.18	0.13	0.74	1.06	13.0	0.3850	20.1	111	3852	275	4274	395
MgRich2_1	1	36	0.25	0.39	0.48	1.25	1.23	9.0	0.4049	9.8	98	3927	140	3840	254
MgRich2_2	-18	70	0.12	0.24	0.22	0.92	1.09	15.3	0.4378	21.0	104	4044	282	4197	456
MgRich2_4	-34	41	0.11	0.26	0.27	1.02	1.32	12.4	0.4347	12.9	90	4034	180	3642	335
CaRich_1	50	93	0.06	0.22	0.09	0.41	1.14	24.0	0.4134	25.8	103	3958	340	4070	686
CaRich_2	-2	33	0.10	0.26	0.18	0.68	1.04	9.9	0.3834	9.8	113	3845	140	4346	306
CaRich_3	20	100	0.19	0.43	0.42	0.98	1.33	14.0	0.3767	19.2	95	3819	264	3618	376
CaRich_4	-5	40	0.16	0.27	0.27	0.98	1.17	9.8	0.4042	12.6	101	3925	177	3983	285
CaRich_5	-7	52	0.06	0.38	0.09	0.24	1.13	10.2	0.4742	10.9	98	4163	152	4082	300
CaRich_6	19	45	0.03	0.37	0.08	0.21	1.04	9.3	0.4646	12.1	105	4132	169	4347	288

¹Background-corrected ²⁰⁴Pb intensity, calculated using the measured ²⁰⁴(Pb+Hg) and ²⁰²Hg, and a ²⁰⁴Hg/²⁰²Hg of 0.22932.

²Concordance (100 × ²⁰⁶Pb/²³⁸U date / ²⁰⁷Pb/²⁰⁶Pb date).