



***In situ* U-Pb dating of 4 billion year old carbonates in martian meteorite Allan Hills 84001**

Romain Tartèse¹, Ian C. Lyon¹

¹Department of Earth and Environmental Sciences, The University of Manchester, Manchester, M13 9PL, UK

5 *Correspondence to:* Romain Tartèse (romain.tartese@manchester.ac.uk)

Abstract. *In situ* carbonate U-Pb dating studies have proliferated dramatically in recent years. Almost all these studies have targeted relatively young calcite up to Carboniferous in age. To assess the robustness of the carbonate U-Pb chronometer in deep-time, we carried out *in situ* U-Pb analyses in magnesite-ankerite-calcite carbonates in the martian meteorite Allan Hills (ALH) 84001. Carbonates in ALH 84001 formed at *ca.* 3.94 Ga, and there is little evidence that much happened to this rock 10 since then, making it an ideal sample to test the robustness of the U-Pb system in old carbonates. We obtained a concordant date of $3941 \pm 49/110$ Ma ($n = 14$, MSWD = 2.0), which is identical to the step-leaching Rb/Sr date determined previously. These results thus confirm that old carbonates are amenable to U-Pb dating in samples that have had a relatively simple history 15 post-carbonate formation.

1 Introduction

15 Analytical developments in laser ablation – inductively coupled plasma mass spectrometry (LA-ICP-MS) over the last decade have driven important progress in *in situ* dating of carbonates, and in particular of calcite and occasionally dolomite, using the radioactive decay of uranium (U) into lead (Pb) (see the recent review by Roberts *et al.*, 2020, for example). Indeed, biogenic, diagenetic, and vein carbonates can typically incorporate up to *ca.* 10-20 $\mu\text{g.g}^{-1}$ U, and up to *ca.* 100 $\mu\text{g.g}^{-1}$ U in speleothems (e.g., Roberts *et al.*, 2020). Carbonates also incorporate initial Pb, meaning that multiple analyses on carbonate samples often 20 yield linear arrays in a Tera-Wasserburg inverse concordia diagram, providing information on both the $^{207}\text{Pb}/^{206}\text{Pb}$ composition of the initial Pb and the age of formation of the carbonates. Recent applications of carbonate U-Pb dating using LA-ICP-MS include constraining the timing of sedimentation, lithification, and diagenesis (e.g., Drost *et al.*, 2018; Godeau *et al.*, 2018; Mueller *et al.*, 2020; Brigaud *et al.*, 2021), faulting (e.g., Ring and Gerdes, 2016; Roberts and Walker, 2016; Goodfellow *et al.*, 2017; Nuriel *et al.*, 2017; 2019; Hansman *et al.*, 2018; Beaudoin *et al.*, 2018; Holdsworth *et al.*, 2019; Smeraglia *et al.*, 25 2019), aragonite to calcite conversion in ammonites (Li *et al.*, 2014), alteration of oceanic crust (Coogan *et al.*, 2016), veining, hydrothermalism, and mineralisation (Burisch *et al.*, 2017; 2018; Parrish *et al.*, 2018; Walter *et al.*, 2018; Bertok *et al.*, 2019; Drake *et al.*, 2019; 2020; MacDonald *et al.*, 2019), palaeoclimate reconstructions (Nicholson *et al.*, 2020), and hominin dispersion (Scardia *et al.*, 2019), for example.



30 All these *in situ* studies have targeted relatively young samples (younger than *ca.* 465 Ma), with two-third of dates younger
than 50 Ma, and all but two younger than 300 Ma (Supplementary Table S1). This is consistent with the accepted idea that
carbonates are not very resistant to resetting of their U-Pb isotope systematics when thermal- and/or fluid-related alteration
events take place after their formation (e.g., Roberts *et al.*, 2020). However, a few studies have focused on dating older
35 carbonate samples, using wet chemistry to isolate Pb isotopes, and obtained Pb/Pb isochron dates ranging between *ca.* 1.60
and 2.84 Ga (Moorbath *et al.*, 1987; Bau *et al.*, 1999; Ray *et al.*, 2003; Sarangi *et al.*, 2004; Farey *et al.*, 2013). These Pb/Pb
dates have been interpreted as dating the deposition of these carbonates, suggesting that in some settings the Pb isotope
systematics of carbonates can remain undisturbed for billions of years. To further assess the robustness of the carbonate U-Pb
chronometer in deep time, we decided to attempt *in situ* LA-ICP-MS U-Pb dating of carbonates in the martian meteorite Allan
40 Hills 84001 (ALH 84001). The formation of these carbonates has been dated at 3.94 ± 0.02 Ga (2σ) using Rb-Sr analyses on
acid leachates via thermal ionisation mass spectrometry (Borg *et al.*, 1999; Beard *et al.*, 2013; date recalculated using a ^{87}Rb
decay constant of $1.3972 \times 10^{-11} \text{ yr}^{-1}$; Villa *et al.*, 2015).

The meteorite ALH 84001 is an orthopyroxenite, a cumulate rock mostly comprising orthopyroxene, olivine, and chromite
(e.g., Mittlefehldt, 1994), which formed *ca.* 4.1 Ga ago as suggested by Lu-Hf and Pb/Pb dating (Bouvier *et al.*, 2009; Lapan
45 *et al.*, 2010). Carbonate-rich areas are irregularly scattered throughout ALH 84001, appearing as spherical or hemispherical
globules, discs along fractures, and irregular fillings in orthopyroxene (see review by Treiman (2021), and references therein).
The patches of carbonates show strong compositional zoning ranging from calcite-rich to magnesite-siderite solid solution
compositions (e.g., Corrigan and Harvey, 2004; Holland *et al.*, 2005). These carbonates likely formed at low temperature (*ca.*
10-20 °C; Halevy *et al.*, 2011; del Real *et al.*, 2016) during fluid-rock interactions that were broadly contemporaneous with
50 the main shock event recorded in ALH 84001 (e.g., Treiman, 2021). Dating carbonate-rich fractions using the Rb-Sr system
yielded a precise formation age of 3.94 ± 0.02 Ga (2σ ; Borg *et al.*, 1999; Beard *et al.*, 2013), which is consistent with a less
precise Pb/Pb isochron corresponding to a date of 4.045 ± 0.090 Ga (2σ ; Borg *et al.*, 1999). As stated by Treiman (2021),
“*there is little evidence that anything had happened to ALH 84001 since 3.9 Ga*” and the time of carbonate formation until it
was launched from Mars *ca.* 14 Ma ago (Eugster *et al.*, 1997), which is crucial with respect to the U-Pb system having remained
55 closed since ALH 84001 carbonates formed.

2 Studied sample

The studied polished section (Fig. 1) was derived from a chip of the ALH84001,287 allocation from the NASA Ancient Mars
Meteorite Program. The section contains patches of carbonates, associated with chromite, in between larger orthopyroxene
grains (Fig. 1). The carbonates display the range of compositions typical for ALH 84001 carbonates, from Mg-rich magnesite
60 to Ca-rich calcite, with intermediate Fe-rich ankerite areas (Fig. 1). The Mg- and Fe-rich carbonates seem to be part of broken
rosettes, while Ca-rich carbonates appear associated with maskelynite (Fig. 1).



3 Laser Ablation – Inductively Coupled Plasma Mass Spectrometer analyses

U-Pb analyses were carried out at the University of Manchester using a Teledyne Photon Machines Analyte Excite+ 193 nm ArF excimer laser ablation system equipped with a HelEx II active 2-volume ablation cell, coupled to an Agilent 8900 triple quadrupole Inductively Coupled Plasma Mass Spectrometer (ICP-MS) using a signal-smoothing device (see Supplementary Table S2) for a summary of the analytical setup and data processing procedure).

The material ablated from target carbonates was carried to the ICP-MS by high purity He, which was mixed with Ar before injection into the plasma source. High purity N₂ was added to the He stream at a flow rate of 2 mL·min⁻¹ to enhance sensitivity. Tuning of the ICP-MS and mass calibration were performed at the start of the analytical session by optimising the ion signals during ablation of the NIST SRM 612 reference glass, while maintaining ²³⁸U⁺/²³²Th⁺ close to unity and minimising the ²³²Th¹⁶O⁺/²³²Th⁺ ratio (*ca.* 0.3%). Glass and carbonates were ablated using a 25 µm laser beam size, a fluence of 4 J·cm⁻², and a repetition rate of 5 Hz. Each analyses lasted 50 s and was preceded by 30 s counting time of the gas blank (background). The masses analysed and corresponding dwell times are reported in Supplementary Table S2.

The reference glass NIST614 (0.823 µg·g⁻¹ U, 2.32 µg·g⁻¹ Pb; Jochum *et al.*, 2011) was used to correct for ²⁰⁷Pb/²⁰⁶Pb fractionation, while mass bias correction of the measured ²³⁸U/²⁰⁶Pb ratios was carried out using repeated analyses of the reference calcite WC-1, which has a thermal ionisation mass spectrometry (TIMS) age of 254.4 ± 6.4 Ma (Roberts *et al.*, 2017). To ensure accuracy, the Duff Brown Tank (DBT) calcite (64.0 ± 0.7 Ma; Hill *et al.*, 2016) and AUG-B6 calcite (43.0 ± 1.0 Ma; Pagel *et al.*, 2018) were also analysed and used as secondary reference materials.

Data processing was carried out using Iolite v4.5, using the NIST614 glass as primary reference material to remove instrument baseline contributions, mass bias of Pb isotopes, and downhole fractionation and instrumental drift of ²⁰⁶Pb/²³⁸U ratios (Paton *et al.*, 2011). The reproducibility obtained on NIST614 for ²⁰⁷Pb/²⁰⁶Pb (±1.7%, *n* = 9, 95% confidence level) and ²⁰⁶Pb/²³⁸U (±1.6%, *n* = 9, 95% confidence level) ratios were propagated by quadrature addition into each analysis ²⁰⁷Pb/²⁰⁶Pb and ²⁰⁶Pb/²³⁸U individual uncertainties. Repeated analyses of the NIST612 glass yielded an average ²⁰⁷Pb/²⁰⁶Pb ratio of 0.871 ± 0.044 (*n* = 8, 2 standard deviation), which is within error of its known ²⁰⁷Pb/²⁰⁶Pb ratio of 0.90745 ± 0.00004 (Baker *et al.*, 2004).

The data obtained for the reference calcite WC-1 were then plotted in a Tera-Wasserburg diagram using IsoplotR (Vermeesch, 2018), and yielded a lower intercept uncorrected date of 255.2 ± 5.9 Ma (95% confidence level, MSWD = 1.1, *n* = 10) for a discordia anchored at a common ²⁰⁷Pb/²⁰⁶Pb ratio of 0.85 (Roberts *et al.*, 2017). To obtain the known intercept age of 254.4 ± 6.4 Ma for the WC-1 calcite, we applied a linear correction factor of 1.0031 to the ²⁰⁶Pb/²³⁸U ratios, which we also applied to all the samples analysed in the session. All calculated dates are associated with two uncertainties, the first one

including the random uncertainties for each analysis (internal uncertainties on measured $^{207}\text{Pb}/^{206}\text{Pb}$ and $^{206}\text{Pb}/^{238}\text{U}$ ratios and reproducibility on repeated NIST614 analyses), while systematic uncertainties (2.5% uncertainty on the WC-1 age, and 0.14% and 0.11% on the ^{235}U and ^{238}U decay constants, respectively; Jaffey *et al.*, 1971) are propagated by quadratic addition in the second one.

100

The data obtained on the DBT calcite yielded a lower intercept date of $64.9 \pm 2.2/2.8$ Ma (95% confidence level, MSWD = 2.9, $n = 12$) for a discordia anchored at a common $^{207}\text{Pb}/^{206}\text{Pb}$ ratio of 0.74 (Hill *et al.*, 2016) (Supplementary Fig. S1).

Because of its lower U abundance and younger age, the data obtained on the AUG-B6 calcite are less precise, yielding a concordia date of $40.8 \pm 2.0/2.2$ Ma (95% confidence level, MSWD concordance + equivalence = 2.2, $n = 8$)

105 (Supplementary Fig. S1), which is consistent with a $^{238}\text{U}/^{206}\text{Pb}$ weighted average date of $41.6 \pm 2.1/2.3$ Ma (95% confidence level, MSWD = 0.4, $n = 8$). All results are available in Supplementary Table S3.

4 Results

The carbonates analysed in ALH 84001 contain *ca.* 0.1-0.4 $\mu\text{g.g}^{-1}$ U and 0.1-0.5 $\mu\text{g.g}^{-1}$ Th (Table 1). When plotted in a Tera-Wasserburg $^{207}\text{Pb}/^{206}\text{Pb}$ vs. $^{238}\text{U}/^{206}\text{Pb}$ diagram, ALH 84001 carbonates yield a concordant date of $3941 \pm 49/110$ Ma ($n = 14$, 110 MSWD = 2.0; Fig. 2A), which is identical to a weighted mean $^{207}\text{Pb}/^{206}\text{Pb}$ date of $3967 \pm 56/113$ Ma ($n = 14$, MSWD = 1.9; Fig. 2B). Carbonate analyses plot on the concordia curve, indicating that they do not contain appreciable amount of common Pb. This is consistent with the measured ^{204}Pb intensities that are within error of 0 counts per second (Table 1).

5 Discussion and implications

5.1 Comparison with previous ALH 84001 carbonate dating studies

115 *In situ* U-Pb dating of carbonates in ALH 84001 using LA-ICP-MS yields a concordia date of $3941 \pm 49/110$ Ma, which is identical to the carbonate step-leaching Rb-Sr isochron date of 3.94 ± 0.02 Ga (Borg *et al.*, 1999; Beard *et al.*, 2013) and the less precise Pb/Pb isochron date of 4.045 ± 0.090 Ga (Borg *et al.*, 1999). Before propagating the $\pm 2.5\%$ uncertainty associated with the age of the primary U-Pb reference carbonate WC-1, and the uncertainties associated with the ^{238}U and ^{235}U decay constants, the carbonate U-Pb concordia date is accompanied with a fairly precise 2σ uncertainty of $\pm 1.2\%$, which increases 120 to $\pm 2.8\%$ when all uncertainties are propagated. This suggests that *in situ* U-Pb dating of carbonates has the potential to yield precise dates, but also highlights the need to reduce uncertainties on reference materials. Our LA-ICP-MS results also indicate that using a calcite primary reference material for correcting U/Pb fractionation in Mg- and Fe-rich carbonate matrices, such as magnesite and ankerite, produce accurate dates (within the obtained uncertainties). A final point worth highlighting is the fact that carbonates in ALH 84001 do not contain appreciable amount of common Pb, as indicated by their concordant U-Pb 125 date of *ca.* 3.94 Ga and their ^{204}Pb intensities within error of 0 count per second. This is unusual, as in most examples in



terrestrial systems carbonates, do contain common Pb incorporated during their crystallisation (e.g., Roberts *et al.*, 2020). This observation suggests that the fluids from which ALH 84001 carbonates formed contained very little Pb.

5.2 Robustness of the carbonate U-Pb chronometer and further applications

Our *in situ* LA-ICP-MS analyses confirm that carbonates in ALH 84001 formed *ca.* 3.94 Gyr-ago, and that the U-Pb 130 chronometer in these carbonates has remained closed to any disturbance event since they formed. This is consistent with the suggestion that not much happened to ALH 84001 between 3.9 Ga and its launch from Mars 14 Myr-ago (Treiman, 2021), and indicates that this latter event did not reset the carbonate U-Pb chronometer. Carbonates in terrestrial Archean samples are probably not the best suited for U-Pb dating as most Archean formations would have been heated up to at least low greenschist 135 metamorphic conditions and/or been affected by hydrothermal alteration, because the Earth is geologically active and harbours a complex hydrological cycle. On the other hand, results of this study open up opportunities for dating old carbonates in samples that have had a relatively simple history post-carbonate formation. For example, volatile-rich carbonaceous chondrites (e.g., CI and CM chondrites) typically contain carbonates formed during fluid-rock interactions on their parent-asteroids *ca.* 4563-4561 Myr-ago (e.g., Jilly-Rehak *et al.*, 2017, and references therein), which is within 10 Myr of the formation of the first 140 solids in the Solar System. After this phase of early hydrothermal alteration, it is thought that not much happens to these samples on their parent asteroids until they end up on the Earth as meteorite fragments. Carbonates in carbonaceous chondrites could thus be prime targets to further constrain the timing of hydrothermal alteration on volatile-rich asteroids using the U-Pb dating chronometer.

Data availability

Raw LA-ICP-MS data and metadata are provided in Supplementary Material.

145 Author contribution

ICL acquired and prepared the sample. RT designed the experiment, performed the analysis, and interpreted the results. RT prepared the manuscript with contributions from ICL.

Competing interests

The authors declare that they have no conflict of interest.

150 Acknowledgments

We acknowledge the provision of the sample of ALH84001 from the NASA Ancient Mars Meteorite Programme, and funding from STFC (#ST/P005225/1 to RT), and The University of Manchester and STFC (grant #ST/S002170/1) for funding the LA-



ICP-MS facility. Many thanks to Nick Roberts for providing us with the WC-1 calcite, Benjamin Brigaud and Thomas Blaise for the AUG-B6 calcite, and Victor Polyak for the Duff Brown Tank calcite.

155 References

Baker, J., Peate, D., Waught, T., and Meyzen, C.: Pb isotopic analysis of standards and samples using a ^{207}Pb – ^{204}Pb double spike and thallium to correct for mass bias with a double-focusing MC-ICP-MS, *Chem. Geol.*, 211, 275–303, 2004.

Bau, M., Romer, R. L., Lüders, V., and Beukes, N. J.: Pb, O, and C isotopes in silicified Moodraai dolomite (Transvaal Supergroup, South Africa): implications for the composition of Paleoproterozoic seawater and ‘dating’ the increase of oxygen in the Precambrian atmosphere, *Earth Planet. Sci. Lett.*, 174, 43–57, 1999.

160 Beard, B. L., Ludois, J. M., Lapen, T. J., and Johnson, C. M.: Pre-4.0 billion year weathering on Mars constrained by Rb–Sr geochronology on meteorite ALH84001, *Earth Planet. Sci. Lett.*, 361, 173–182, 2013.

Beaudoin, N., Lacombe, O., Roberts, N. M., and Koehn, D.: U–Pb dating of calcite veins reveals complex stress evolution and thrust sequence in the Bighorn Basin, Wyoming, USA, *Geology*, 46, 1015–1018, 2018.

165 Bertok, C., Barale, L., d'Atri, A., Martire, L., Piana, F., Rossetti, P., and Gerdes, A.: Unusual marbles in a non-metamorphic succession of the SW Alps (Valdieri, Italy) due to early Oligocene hydrothermal flow, *Int. J. Earth Sci.*, 108, 693–712, 2019.

Borg, L. E., Connelly, J. N., Nyquist, L. E., Shih, C. Y., Wiesmann, H., and Reese, Y.: The age of the carbonates in martian meteorite ALH84001, *Science*, 286, 90–94, 1999.

170 Bouvier, A., Blichert-Toft, J., and Albarede, F.: Martian meteorite chronology and the evolution of the interior of Mars, *Earth Planet. Sci. Lett.*, 280, 285–295, 2009.

Brigaud, B., Andrieu, S., Blaise, T., Haurine, F., and Barbarand, J.: Calcite uranium–lead geochronology applied to hardground lithification and sequence boundary dating, *Sedimentology*, 68, 168–195, 2021.

Burisch, M., Gerdes, A., Walter, B. F., Neumann, U., Fettel, M., and Markl, G.: Methane and the origin of five-element veins: mineralogy, age, fluid inclusion chemistry and ore forming processes in the Odenwald, SW Germany, *Ore Geol. Rev.*, 81, 42–61, 2017.

175 Burisch, M., Walter, B. F., Gerdes, A., Lanz, M., and Markl, G.: Late-stage anhydrite–gypsum–siderite–dolomite–calcite assemblages record the transition from a deep to a shallow hydrothermal system in the Schwarzwald mining district, SW Germany, *Geochim. Cosmochim. Ac.*, 223, 259–278, 2018.

Coogan, L. A., Parrish, R. R., and Roberts, N. M.: Early hydrothermal carbon uptake by the upper oceanic crust: Insight from in situ U–Pb dating, *Geology*, 44, 147–150, 2016.

180 Corrigan, C. M. and Harvey, R. P.: Multi-generational carbonate assemblages in martian meteorite Allan Hills 84001: implications for nucleation, growth, and alteration, *Meteorit. Planet. Sci.*, 39, 17–30, 2004.

del Real, P. G., Maher, K., Kluge, T., Bird, D. K., Brown, G. E., and John, C. M.: Clumped-isotope thermometry of magnesium carbonates in ultramafic rocks, *Geochim. Cosmochim. Acta*, 193, 222–250, 2016.

185 Drake, H., Roberts, N. M. W., Heim, C., Whitehouse, M. J., Siljeström, S., Kooijman, E., Broman, C., Ivarsson, M., and Åström, M. E.: Timing and origin of natural gas accumulation in the Siljan impact structure, Sweden, *Nat. Commun.*, 10, 1–14, 2019.



190 Drake, H., Roberts, N. M., and Whitehouse, M. J.: Geochronology and Stable Isotope Analysis of Fracture-fill and Karst
Mineralization Reveal Sub-Surface Paleo-Fluid Flow and Microbial Activity of the COSC-1 Borehole, Scandinavian
Caledonides, *Geosciences*, 10, 56, <https://doi.org/10.3390/geosciences10020056>, 2020.

195 Drost, K., Chew, D., Petrus, J. A., Scholze, F., Woodhead, J. D., Schneider, J. W., and Harper, D. A.: An Image Mapping
Approach to U-Pb LA-ICP-MS Carbonate Dating, and Applications to Direct Dating of Carbonate Sedimentation, *Geochem.
Geophys. Geosy.*, 19, 4631-4648, 2018

200 Eugster, O., Weigel, A., and Polnau, E.: Ejection times of martian meteorites, *Geochim. Cosmochim. Acta*, 61, 2749-2757,
1997.

Fairey, B., Tsikos, H., Corfu, F., and Polteau, S.: U-Pb systematics in carbonates of the Postmasburg Group, Transvaal
Supergroup, South Africa: primary versus metasomatic controls, *Precambrian Res.*, 231, 194-205, 2013.

205 Godeau, N., Deschamps, P., Guihou, A., Leonide, P., Tendil, A., Gerdes, A., Hamelin, B., and Girard, J. P.: U-Pb dating of
calcite cement and diagenetic history in microporous carbonate reservoirs: Case of the Urgonian Limestone,
France, *Geology*, 46, 247-250, 2018.

210 Goodfellow, B. W., Viola, G., Bingen, B., Nuriel, P., and Kylander-Clark, A. R.: Palaeocene faulting in SE Sweden from U-
Pb dating of slickenfibre calcite, *Terra Nova*, 29, 321-328, 2017.

Halevy, I., Fischer, W. W., and Eiler, J. M.: Carbonates in the martian meteorite Allan Hills 84001 formed at $18 \pm 4^\circ\text{C}$ in a
near-surface aqueous environment, *Proc. Natl. Acad. Sci. USA*, 108, 16895-16899, 2011.

215 Hansman, R. J., Albert, R., Gerdes, A., and Ring, U.: Absolute ages of multiple generations of brittle structures by U-Pb
dating of calcite, *Geology*, 46, 207-210, 2018.

Hill, C. A., Polyak, V. J., Asmerom, Y., and Provencio, P.: Constraints on a Late Cretaceous uplift, denudation, and incision
of the Grand Canyon region, southwestern Colorado Plateau, USA, from U-Pb dating of lacustrine limestone: U-Pb age of
lacustrine limestone, *Tectonics*, 35, 896-906, 2016.

220 Holdsworth, R. E., McCaffrey, K. J. W., Dempsey, E., Roberts, N. M. W., Hardman, K., Morton, A., Feely, M., Hunt, J.,
Conway, A., and Robertson, A.: Natural fracture propping and earthquake-induced oil migration in fractured basement
reservoirs, *Geology*, 47, 700-704, 2019.

Holland, G., Saxton, J., Lyon, I., and Turner, G.: Negative $\delta^{18}\text{O}$ values in Allan Hills 84001 carbonate: possible evidence for
water precipitation on Mars, *Geochim. Cosmochim. Acta*, 69, 1359-1370, 2005.

225 Jaffey A. H., Flynn, K. F., Glendenin, L. E., Bentley, W. C., and Essling, A.M.: Precision measurement of half-lives and
specific activities of ^{235}U and ^{238}U , *Physical Review C*, 4, 1889-1906, 1971.

Jilly-Rehak, C. E., Huss, G. R., and Nagashima, K.: ^{53}Mn - ^{53}Cr radiometric dating of secondary carbonates in CR chondrites:
Timescales for parent body aqueous alteration, *Geochim. Cosmochim. Acta*, 201, 224-244, 2017.

Jochum, K. P., Weis, U., Stoll, B., Kuzmin, D., Yang, Q., Raczek, I., Jacob, D. E., Stracke, A., Birbaum, K., Frick, D. A.,
220 Gunther, D., and Enzweiler, J.: Determination of reference values for NIST SRM 610-617 glasses following ISO guidelines,
Geostand. Geoanal. Res., 35, 397-429, 2011.

Lapen, T., Righter, M., Brandon, A., Debaille, V., Beard, B., Shafer, J., and Peslier, A.: A younger age for ALH84001 and
its geochemical link to shergottite sources in Mars, *Science*, 328, 347-351, 2010.

Li, Q., Parrish, R. R., Horstwood, M. S. A., and McArthur, J. M.: U-Pb dating of cements in Mesozoic ammonites, *Chem.
Geol.*, 376, 76-83, 2014.



MacDonald, J. M., Faithfull, J. W., Roberts, N. M. W., Davies, A. J., Holdsworth, C. M., Newton, M., Williamson, S., Boyce, A., and John, C. M.: Clumped-isotope palaeothermometry and LA-ICP-MS U–Pb dating of lava-pile hydrothermal calcite veins, *Contrib. Mineral. Petr.*, 174, 63, <https://doi.org/10.1007/s00410-019-1599-x>, 2019.

Mittlefehldt, D. W.: ALH84001, a cumulate orthopyroxenite member of the Martian meteorite clan. *Meteoritics*, 29, 214-221, 1994.

Moorbath, S., Taylor, P. N., Orpen, J. L., Treloar, P., and Wilson, J. F.: First direct radiometric dating of Archaean stromatolitic limestone, *Nature*, 326, 865-867, 1987.

Mueller, M., Igbokwe, O. A., Walter, B., Pederson, C. L., Riechelmann, S., Richter, D. K., Albert, R., Gerdes, A., Buhl, D., Neuser, R. D., Bertotti, G., and Immenhauser, A.: Testing the preservation potential of early diagenetic dolomites as geochemical archives, *Sedimentology*, 67, 849-881, 2020.

Nicholson, S. L., Pike, A. W., Hosfield, R., Roberts, N. M. W., Sahy, D., Woodhead, J., Cheng, H., Edwards, R. L., Affolter, S., Leuenberger, M., and Burns, S. J.: Pluvial periods in Southern Arabia over the last 1.1 million-years, *Quaternary Sci. Rev.*, 229, 106112, <https://doi.org/10.1016/j.quascirev.2019.106112>, 2020.

Nuriel, P., Craddock, J., Kylander-Clark, A. R., Uysal, T., Karabacak, V., Dirik, R. K., Hacker, B. R., and Weinberger, R.: Reactivation history of the North Anatolian fault zone based on calcite age-strain analyses, *Geology*, 47, 465-469, 2019.

Nuriel, P., Weinberger, R., Kylander-Clark, A. R. C., Hacker, B. R., and Craddock, J. P.: The onset of the Dead Sea transform based on calcite age-strain analyses, *Geology*, 45, 587-590, 2017.

Pagel, M., Bonifacie, M., Schneider, D. A., Gautheron, C., Brigaud, B., Calmels, D., Cros, A., Saint-Bezar, B., Landrein, P., Sutcliffe, C., Davis, D., and Chaduteau, C.: Improving paleohydrological and diagenetic reconstructions in calcite veins and breccia of a sedimentary basin by combining D47 temperature, $d^{18}\text{O}_{\text{water}}$ and U-Pb age, *Chem. Geol.*, 481, 1-17, 2018.

Parrish, R. R., Parrish, C. M., and Lasalle, S.: Vein calcite dating reveals Pyrenean orogen as cause of Paleogene deformation in southern England, *J. Geol. Soc.*, 175, 425-442, 2018.

Paton C., Hellstrom J., Paul B., Woodhead J., and Hergt J.: Iolite: Freeware for the visualisation and processing of mass spectrometric data, *J. Anal. Atom. Spectro.*, 26, 2508-2518, 2011.

Ray, J. S., Veizer, J., and Davis, W. J.: C, O, Sr and Pb isotope systematics of carbonate sequences of the Vindhyan Supergroup, India: age, diagenesis, correlations and implications for global events, *Precambrian Res.*, 121, 103-140, 2003.

Ring, U. and Gerdes, A.: Kinematics of the Alpenrhein-Bodensee graben system in the Central Alps: Oligocene/Miocene transtension due to formation of the Western Alps arc, *Tectonics*, 35, 1367-1391, 2016.

Roberts, N. M. W. and Walker, R. J.: U-Pb geochronology of calcite-mineralized faults: Absolute timing of rift-related fault events on the northeast Atlantic margin, *Geology*, 44, 531-534, 2016.

Roberts, N. M. W., Drost, K., Horstwood, M. S. A., Condon, D. J., Chew, D., Drake, H., Milodowski, A. E., McLean, N. M., Smye, A. J., Walker, R. J., Haslam, R., Hodson, K., Imber, J., Beaudoin, N., and Lee, J. K.: Laser ablation inductively coupled plasma mass spectrometry (LA-ICP-MS) U–Pb carbonate geochronology: strategies, progress, and limitations, *Geochronology*, 2, 33-61, 2020.

Roberts, N. M. W., Rasbury, E. T., Parrish, R. R., Smith, C. J., Horstwood, M. S. A., and Condon, D. J.: A calcite reference material for LA-ICP-MS U-Pb geochronology, *Geochem. Geophys.*, 18, 2807-2814, 2017.



Sarangi, S., Gopalan, K., and Kumar, S.: Pb–Pb age of earliest megascopic, eukaryotic alga bearing Rohtas Formation, Vindhyan Supergroup, India: implications for Precambrian atmospheric oxygen evolution, *Precambrian Res.*, 132, 107-121, 2004.

265 Scardia, G., Parenti, F., Miggins, D. P., Gerdes, A., Araujo, A. G., and Neves, W. A.: Chronologic constraints on hominin dispersal outside Africa since 2.48 Ma from the Zarqa Valley, Jordan, *Quaternary Sci. Rev.*, 219, 1-19, 2019.


Smeraglia, L., Aldega, L., Billi, A., Carminati, E., Di Fiore, F., Gerdes, A., Albert, R., Rossetti, F., and Vignaroli, G.: Development of an intra-wedge tectonic mélange by out-of-sequence thrusting, buttressing, and intraformational rheological contrast, Mt. Massico ridge, Apennines, Italy, *Tectonics*, 38, 1223-1249, 2019.

270 Treiman, A. H.: Uninhabitable and Potentially Habitable Environments on Mars: Evidence from Meteorite ALH 84001, *Astrobiology*, 21, 940-953, 2021.


Vermeesch, P.: IsoplotR: a free and open toolbox for geochronology, *Geosci. Frontiers*, 9, 1479-1493, 2018.

Villa, I. M., De Bievre, P., Holden, N. E., and Renne, P. R.: IUPAC–IUGS recommendation on the half-life of ^{87}Rb , *Geochim. Cosmochim. Acta*, 164, 382-385, 2015.

275 Walter, B. F., Gerdes, A., Kleinhanß, I. C., Dunkl, I., von Eynatten, H., Kreissl, S., and Markl, G.: The connection between hydrothermal fluids, mineralization, tectonics and magmatism in a continental rift setting: Fluorite Sm-Nd and hematite and carbonates U-Pb geochronology from the Rhinegraben in SW Germany, *Geochim. Cosmochim. Ac.*, 240, 11-42, 2018.



280 **Figure 1: Backscattered electron image (A) and composite X-ray map (B) of the target carbonate patches in ALH 84001. Mineral abbreviations are ank = ankerite, cal = calcite, chr = chromite, mgs = magnesite, msk = maskelynite, opx = orthopyroxene.**



285 **Figure 2: Tera-Wasserburg $^{207}\text{Pb}/^{206}\text{Pb}$ vs. $^{238}\text{U}/^{206}\text{Pb}$ diagram (A) and $^{207}\text{Pb}/^{206}\text{Pb}$ dates (B) obtained on ALH 84001 carbonates. Ellipses (A) and error bars (B) correspond to 2σ uncertainties.**



Table 1: LA-ICP-MS results for ALH 84001 carbonates.

Analysis #	^{204}Pb (cps) ¹	±	Pb ($\mu\text{g/g}$)	U ($\mu\text{g/g}$)	Th ($\mu\text{g/g}$)	Ratios $\pm 2\sigma$ (%)			Dates $\pm 2\sigma$ (Ma)						
						$^{238}\text{U}/$ ^{206}Pb	±	$^{207}\text{Pb}/$ ^{206}Pb	±	conc. (%) ²	$^{206}\text{Pb}/$ ^{238}U	±			
MgRich_1	100	110	0.11	0.26	0.20	0.76	1.06	29.2	0.4037	28.3	109	3923	370	4270	853
MgRich_3	-123	82	0.12	0.14	0.08	0.57	1.17	23.9	0.5356	25.2	92	4342	327	3976	671
MgRich_4	10	50	0.13	0.19	0.12	0.64	1.23	18.1	0.4295	24.9	96	4016	329	3835	504
MgRich_5	-4	42	0.20	0.41	0.50	1.21	1.29	8.8	0.3570	12.5	99	3737	178	3698	244
MgRich_6	27	49	0.10	0.18	0.13	0.74	1.06	13.0	0.3850	20.1	111	3852	275	4274	395
MgRich2_1	1	36	0.25	0.39	0.48	1.25	1.23	9.0	0.4049	9.8	98	3927	140	3840	254
MgRich2_2	-18	70	0.12	0.24	0.22	0.92	1.09	15.3	0.4378	21.0	104	4044	282	4197	456
MgRich2_4	-34	41	0.11	0.26	0.27	1.02	1.32	12.4	0.4347	12.9	90	4034	180	3642	335
CaRich_1	50	93	0.06	0.22	0.09	0.41	1.14	24.0	0.4134	25.8	103	3958	340	4070	686
CaRich_2	-2	33	0.10	0.26	0.18	0.68	1.04	9.9	0.3834	9.8	113	3845	140	4346	306
CaRich_3	20	100	0.19	0.43	0.42	0.98	1.33	14.0	0.3767	19.2	95	3819	264	3618	376
CaRich_4	-5	40	0.16	0.27	0.27	0.98	1.17	9.8	0.4042	12.6	101	3925	177	3983	285
CaRich_5	-7	52	0.06	0.38	0.09	0.24	1.13	10.2	0.4742	10.9	98	4163	152	4082	300
CaRich_6	19	45	0.03	0.37	0.08	0.21	1.04	9.3	0.4646	12.1	105	4132	169	4347	288

¹Background-corrected ^{204}Pb intensity, calculated using the measured $^{204}\text{Pb} + \text{Hg}$ and ^{202}Hg , and a $^{204}\text{Hg}/^{202}\text{Hg}$ of 0.22932.

²Concordance ($100 \times 206\text{Pb}/^{238}\text{U}$ date / $^{207}\text{Pb}/^{206}\text{Pb}$ date).