

1 **Bayesian age-depth modelling applied to varve and radiometric**  
2 **dating to optimize the transfer of an existing high-resolution**  
3 **chronology to a new composite sediment profile from Holzmaar**  
4 **(West-Eifel Volcanic Field, Germany)**

5 Stella Birlo<sup>1\*</sup>, Wojciech Tylmann<sup>2</sup>, Bernd Zolitschka<sup>1</sup>

6 1 - University of Bremen, Institute of Geography, GEOPOLAR, Bremen, Germany

7 2 - University of Gdańsk, Faculty of Oceanography and Geography, Gdańsk, Poland

8 \*Corresponding author: sbirlo@uni-bremen.de

## 9 Abstract

10 This study gives an overview of ~~different methods to integrate information from a varve~~  
11 ~~chronology and radiometric measurements in the Bayesian tool Bacon~~~~different varve~~  
12 ~~integration methods with Bacon~~. These techniques will become important for the future as  
13 technologies evolve with more sites being revisited for the application of new and high-resolution  
14 scanning methods. Thus, the transfer of existing chronologies will become necessary, because the  
15 recounting of varves will be too time consuming and expensive to be funded.

16 We introduce new sediment cores from Holzmaar (West-Eifel Volcanic Field, Germany), a volcanic  
17 maar lake with a well-studied varve<sup>d</sup> record. Four different age-depth models ~~(A-D)~~ have been  
18 calculated for the new composite sediment profile (HZM19) using Bayesian ~~statistics-modelling~~  
19 with Bacon. All models incorporate new Pb-210 and Cs-137 dates for the top of the record, the  
20 latest calibration curve (IntCal20) for radiocarbon ages as well as the new age estimation for the  
21 Laacher See Tephra. Model A is based on previously published radiocarbon measurements only,  
22 while Models B-D integrate the previously published varve chronology (VT-99) with different  
23 approaches. Model B rests upon radiocarbon data, while parameter settings are obtained from  
24 sedimentation rates derived from VT-99. Model C is based on radiocarbon dates and on VT-99 as  
25 several normal-distributed tie-points, while Model D is segmented into four sections: Sections 1  
26 and 3 are based on VT-99 only, whereas Sections 2 and 4 rely on Bacon age-depth models  
27 including additional information from VT-99. In terms of accuracy, the parameter-based  
28 integration Model B shows little improvement over the non-integrated approach, whereas the tie  
29 point-based integration Model C reflects the complex accumulation history of Holzmaar much  
30 better. Only the segmented and parameter-based age-integration approach of Model D adapts and  
31 improves VT-99 by replacing sections of higher counting errors with Bayesian modelling of  
32 radiocarbon ages and thus efficiently makes available the best possible and ~~most~~ precise age-  
33 depth model for HZM19. This approach will value all ongoing ~~and~~ high-resolution investigations  
34 for a better understanding of decadal-scale Holocene environmental and climatic variations.

35 Keywords: Lacustrine sediments, Varves, Bayesian age-depth modelling, Bacon, Radiometric  
36 dating

## 1. Introduction

Terrestrial archives from lakes have the potential to provide information about climate and the human history of its catchment area beyond instrumental and historical data (Berglund, 1986; Last and Smol, 2001a, b; Cohen, 2003). In the late 1980s, [gravity coring](#) (Kelts et al., 1986), piston coring (Nesje et al., 1987; Wright et al., 1984) and freeze coring techniques (Renberg and Hansson, 1993) for lacustrine sediment records have improved tremendously allowing a better quality of sediments to be recovered from modern lakes. Since then, the new fields of limnogeology and paleolimnology flourished with increasing demand of societies for documentation of natural background data related to questions around acid rain (e.g. Battarbee et al., 1990), environmental pollution (e.g. Renberg et al., 1994) and more and more with a focus on global climate change (e.g. Jenny et al., 2019).

To provide such information not only on local scales but also on larger regional to global scales, investigations from different sites need to be compared and linked. However, such correlations are only successful if the contributing archives are based on robust chronologies. Therefore, precise and reliable age-depth models are the basis for sedimentary investigations and reconstructions of environmental and climatic changes of the past, as [only](#) they ensure intra-site comparability and enable recognition of larger scale patterns. A reliable chronology ~~should~~[can](#) be based on a combination of different dating techniques (multiple dating approach) such as radiometric dating, well-known events [such as tephra layers](#) (Turkey and Lowe, 2001; Davies, 2015), ~~layers (e.g., tephrochronology)~~, historic data (e.g., flood events) or varve counting. The term “varve” (Swedish: layer) was first introduced by De Geer (1912) for outcrops with proglacial sediments and describes finely laminated sediment structures with annual origin. The alternating pale and dark layers are driven by seasonal changes in temperature and precipitation that cause different chemical and biological processes within the lake and its catchment area. When anoxic conditions at the sediment-water-interface are given at least seasonally, i.e. no bioturbation destroys laminations, varves are preserved and provide high-resolution and precise chronologies in calendar years (Zolitschka et al., 2015; Lamoureux, 2001).

Until the 1980s, varve chronologies were the only option for calendar-year chronologies ~~for~~[of](#) sediment records, while AMS radiocarbon dating was still in its infancy and calibration of radiocarbon ages was restricted to [tree rings of](#) the Middle and Late Holocene, if at all applied (Pearson et al., 1977; Olsson, 1986). First reviews about methodological advances in the study of annually laminated sediments appeared at the same time (Anderson and Dean, 1988; O’Sullivan, 1983; Saarnisto, 1986) and [first long and](#) varve-dated reconstructions were published for Elk Lake, USA (Dean et al., 1984) and Lake Valkiajärvi, Finland (Saarnisto, 1985).

Meerfelder Maar and Holzmaar were the first varve-dated lacustrine records covering the entire Holocene and the Late Glacial for Central Europe (Zolitschka, 1989, 1988), followed by records concentrating on the Late Glacial to Holocene transition at Soppensee, Switzerland (Lotter, 1991) and at Lake Gosciarz, Poland (Goslar et al., 1993). As such, the Holzmaar record became one of the best studied lacustrine records in Europe, if not world-wide.

To produce the chronology for HZM19 we test and compare different methods integrating varve counts with radiometric measurements using Bayesian age-depth modelling. The advantage of any modelling approach is that all possible calendar ages of calibrated radiocarbon dates and their probability density functions (PDFs) will be tested by using a repeated random sampling method (Blaauw, 2010; Telford et al., 2004). In addition, using the Bayes theorem allows to incorporate information of the accumulation history known prior to modelling. Thus, calendar ages, which are monotonic with depth and with positive accumulation rates in  $\text{yr cm}^{-1}$  (in sedimentological terms, accumulation rates as they are used for Bayesian age-depth modelling are equivalent to "sedimentation rates", as corroborated by the units used) are calculated (Lacourse and Gajewski, 2020; Trachsel and Telford, 2017). This is different and an advantage if compared to the "Classical Age-depth Modelling" carried out by CLAM (Blaauw, 2010).

Currently established programs that use Bayesian statistics are Oxcal (Bronk Ramsey, 2008), BChron (Haslett and Parnell, 2008) and Bacon (Blaauw and Christen, 2011), all of which differ in terms of parameter settings and handling of outliers. In this study, we focus on varve-counting integration methods using Bacon (rBacon version 2.5.7; Blaauw et al., 2021; Blaauw and Christen, 2011) for the R programming language (version 4.1.1; R Core Team, 2021), as it is one of the most often used software package in paleo studies and provides many different ways for implementing additional information. Bacon uses a Markov Chain Monte Carlo (MCMC) sampling strategy to model the accumulation history piecewise using a gamma autoregressive semi-parametric model (Blaauw and Christen, 2011). The accumulation rate of each segment depends on the accumulation rate of the previous segment. Dates are treated using a student's t-distribution. Although Bacon provides default values, the accumulation rate is controlled by two adjustable prior distributions (prior model), the accumulation rate as a gamma distribution and the memory, which describes the dependence of accumulation rates between neighbouring depths as a beta distribution. Both latter parameters are defined by a shape and a strength prior, respectively, in addition to a mean prior. Furthermore, we make use of the number of segments (thick-parameter) recommended by Bacon. The program also allows to incorporate information about hiati and slump events in the profile.

We concentrate on approaches using the Bacon package for the R statistical programming software (Blaauw and Christen, 2011), whereas literature also provides comparable methods for

alternative Bayesian age-depth modelling software, such as OxCal (Martin-Puertas et al., 2021; Bronk Ramsey, 2008; Vandergoes et al., 2018), which was also used to integrate varve counting and radiometric dating for the Holocene sediment record HZM96-4a,4b from Holzmaar (Prasad and Baier, 2014). As Bacon provides many different options to incorporate information into the age-depth model, in the literature only few approaches are provided integrating varve and radiocarbon ages (Bonk et al., 2021; Vandergoes et al., 2018; Shanahan et al., 2012). For that reason, we summarize these approaches and compare them directly with each other. This will lead to faster decisions for future studies facing a comparable situation. As Bacon provides many different possibilities to incorporate information in the model, literature provides not many but few very different approaches to integrate varve and radiocarbon ages. For that reason, we sum up those approaches in this study and compare them directly with each other. This might lead to faster decision making for future researchers facing a similar situation.

~~Since the first coring campaign in 1984, several sediment records have been recovered from Holzmaar and numerous studies were carried out with sedimentological, biological, geochemical and geophysical methods (e.g. Zolitschka, 1989; Lottermoser et al., 1993; Hajdas et al., 1995; Raubitschek et al., 1999; Leroy et al., 2000). However, the early sediment records from Holzmaar, although counted and corrected multiple times, still contain sections of high counting uncertainty and thus suffer from optimal core correlation as it is possible today by applying high-resolution scanning techniques and digital line-scan images. Moreover, independent time control of varve chronologies with AMS radiocarbon dating became available only in the 1990s (Hajdas-Skowronek, 1993), while Bayesian age-depth modelling established as a tool for optimizing dating efforts only during the last decade (Ramsey, 2009) and sediment scanning revolutionized limnogeology and paleolimnology over the last 20 years. Therefore, we revisited Holzmaar to obtain fresh sediment cores for the conduction of innovative and high-resolution (sub-millimetre-scale) sediment scanning techniques to be based on an improved age-depth model.~~

As chronologies are always a “running target”, especially as new scientific methods and approaches appear, it is no wonder that the varve chronology for Holzmaar sediments has developed from its first attempt as “Varve Time 1990” (VT-90) (Zolitschka, 1990) to VT-99 ten years later (Zolitschka et al., 2000). In the course of applying ultra-high (sub-mm-scaled) resolution scanning techniques to the new set of sediment cores from Holzmaar (HZM19), VT-99 was/is transferred to HZM19 making use of marker layers and radiocarbon ages for correlation as well as of Bayesian age-depth modelling for the creation of an updated varve chronology (VT-22).

Different to earlier studies, we make use of available radiocarbon dates from Holzmaar not only to correct the varve chronology but to combine it with the independent radiocarbon chronology

141 using Bayesian modelling. This integration approach is not commonly used for lacustrine records  
142 ~~yet~~. Here we select three different methods to integrate varve and radiometric dating and apply it  
143 to the Holzmaar data. ~~We concentrate on approaches using the Bacon package for the R statistical~~  
144 ~~programming software (Blaauw and Christen, 2011), whereas literature also provides~~  
145 ~~comparable methods for alternative Bayesian age depth modelling software, such as OxCal~~  
146 ~~(Martin Puertas et al., 2021; Ramsey, 2008; Vandergoes et al., 2018), which was also used to~~  
147 ~~integrate varve counting and radiometric dating for the Holocene sediment record HZM96-4a,4b~~  
148 ~~from Holzmaar (Prasad and Baier, 2014).~~

149 ~~In this study we discuss the possibilities to integrate and improve different chronologies by~~  
150 ~~combining a varve chronology with modelling approaches. This is accomplished by testing and~~  
151 ~~comparing integration methods with regard to accuracy and precision from the interpolated varve~~  
152 ~~chronology itself and for a Bayesian model without any varve information. With this integration~~  
153 ~~of all age information we produce the most reliable age estimations for the HZM19 record: VT-22.~~  
154 ~~Based on the best model outcome, this master chronology serves as the chronological base for~~  
155 ~~ongoing and future biological, geochemical and geophysical investigations conducted on the new~~  
156 ~~Holzmaar sediment cores (e.g. García et al., 2022).~~

157 The aim of our study is to transfer and optimize the existing varve chronology from HZM-B/C to  
158 the new sediment record HZM19. In addition, we offer an overview about different approaches  
159 for age-depth modelling and their effects on model outcomes to researchers who face comparable  
160 challenges, thus supporting their decision making.

161 For this reason, we discuss the possibilities of integrating and improving the chronology by  
162 combining the varve chronology with modelling approaches using Bacon. This is accomplished by  
163 testing and comparing integration methods with regard to accuracy and precision obtained from  
164 the interpolated varve chronology itself and from a Bayesian model without any varve  
165 information relying on radiocarbon dates only.

166 With this integration of all age information we produce the most reliable age estimations for the  
167 HZM19 record: VT-22. Based on the best model approach, this master chronology of VT-22 serves  
168 as the chronological backbone for ongoing and future biological, geochemical and geophysical  
169 investigations conducted with the new Holzmaar sediment cores (e.g. García et al., 2022).

## 2. Materials and Methods Regional settings and the Holzmaar sediment record

### 2.1 Regional Settings

The late Quaternary volcanic maar lake Holzmaar (425 m a.s.l., 50°7'8" N, 6°52' 45" E) is located in the western central part of the Rhenish Massif in the West-Eifel Volcanic Field (WEVF; Rhineland-Palatinate, Germany, Fig. 1). The WEVF consists of more than [one](#) hundred volcanic cones and maars, of which only nine are water-filled today (Meyer, 2013; Schmincke, 2014). The volcanism in the Eifel region was caused by uplift of the Rhenish Shield since 700 - 800 ka, which started in the NW near Ormont (Meyer and Stets, 2002; Schmincke, 2007). Volcanic activities reached a peak at ca 600 - 450 ka in the central WEVF and then decreased towards Bad Bertrich in the SE (Schmincke, 2007). The uplift is responsible for many eruptive centres at NW-SE trending tectonic faults, along which several phreatomagmatic maar explosions occurred (Büchel, 1993; Lorenz, 1984; Lorenz et al., 2020; Meyer, 1985). One of these eruptions formed the Holzmaar system ca. 40 - 70 ka ago (Büchel, 1993) consisting of three maars with the maar lake of Holzmaar, the raised bog of Dürres Maar and the dry Hetsche or Hitsche Maar (from SE to NW). With 100 m in diameter, the latter is the smallest maar of the WEVF (Fig. 1).

The catchment area of Holzmaar (2.06 km<sup>2</sup>) includes the Sammetbach, a creek that flows in and out of the lake. Due to the low erosive energy of the stream no delta formed in the lake (Scharf, 1987; Zolitschka, 1998a). The geology in the catchment area consists of Devonian metamorphic slates, greywackes and quartzites as well as Quaternary loess and volcanic rocks related to eruptions of the Holzmaar system (Meyer, 2013). Holzmaar is located within a conservation area since 1975 protecting the surrounding beech forest (*Fagus sylvatica* L.), while ca. 60% of the catchment area is in agricultural use (Kienel et al., 2005).

The lake of Holzmaar has a diameter of 300 m (water surface: 58,000 m<sup>2</sup>) and with a maximum water depth of 19-20 m shows a deep and steep-sided morphology typical for maar lakes. Only a small and shallow embayment in the SW interrupts the nearly circular and 1100 m long shoreline. This appendix-like bay developed due to an artificial damming in the late Middle Ages, which was constructed to supply a downstream water mill (Zolitschka, 1998a). For the last glacial, paleolimnological investigations indicate oligotrophic conditions, but eutrophication started already at the onset of the Late Glacial (García et al., 2022). During the Holocene, water quality is affected by human activities, which started during the Neolithic (around 6500 cal. BP) according to pollen analysis (Litt et al., 2009). Together with the inflow of the Sammetbach this caused a steady but slow process of eutrophication and today leads to meso- to eutrophic conditions (Lücke et al., 2003; Scharf and Oehms, 1992; Zolitschka, 1990). The lake is holo- and dimictic with an

anoxic hypolimnion during summer stratification (Scharf and Oehms, 1992). Altogether, this caused a high potential for varves to be formed and preserved.




Figure 1: ESRI Satellite image of the Holzmaar volcanic system and its catchment area (indicated by a white dashed line) with Holzmaar, Dürrer Maar, Hetsche Maar and Sammetbach (blue line, flow direction indicated by arrows). Upper left insert: Location of Holzmaar in Germany (red star). Upper right insert: Bathymetric map with isobaths in meter and coring locations (H2M19-07, -08, -10 and -11) marked by red stars.

The catchment area of Holzmaar (2.06 km<sup>2</sup>) includes the Sammetbach, a creek that flows in and out of the lake. Due to the low erosive energy of the stream no delta formed in the lake (Scharf, 1987; Zolitschka, 1998a). The geology in the catchment area consists of Devonian metamorphic clates, greywackes and quartzites as well as Quaternary loess and volcanic rocks related to eruptions of the Holzmaar system (Meyer, 2013). Holzmaar is located within a conservation area since 1975 protecting the surrounding beech forest (*Fagus sylvatica* L.), while ca. 60% of the catchment area is in agricultural use (Kienel et al., 2005).

Holzmaar has a diameter of 300 m (water surface: 58,000 m<sup>2</sup>) and with a maximum water depth of 19-20 m shows a deep and steep-sided morphology typical for maar lakes. Only a small and shallow embayment in the SW interrupts the nearly circular and 1100 m long shoreline. This appendix-like bay developed due to an artificial damming in the late Middle Ages, which was



~~constructed to supply a downstream water mill (Zolitschka, 1998a). For the last glacial, paleolimnological investigations indicate oligotrophic conditions, but eutrophication started already at the onset of the Late Glacial (García et al., 2022). During the Holocene, water quality is affected by human activities, which started during the Neolithic (around 6500 cal. BP) according to pollen analysis (Litt et al., 2009). Together with the inflow of the Sammetbach this caused a steady but slow process of eutrophication and today leads to meso- to eutrophic conditions (Lücke et al., 2003; Scharf and Oehms, 1992; Zolitschka, 1990). The lake is hol- and dimictic with an anoxic hypolimnion during summer stratification (Scharf and Oehms, 1992). Altogether, this caused a high potential for varves to be formed and preserved.~~

### 3.12.2 Holzmaar Lithology

~~The four parallel cores HZM19-07, -08, -10 and -11 were aligned and correlated to form the composite profile HZM19 (Fig. 2), which includes 24 core sections and reaches to a basal depth of 14.64 m (Table A3). One technical sediment gap exists at a composite depth of 10.90 m. To determine the precise length of this gap, we use core photographs from a previous Holzmaar core (HZM90-H5u) and determined the technical gap with a length of 12.9 cm (Fig. A1).~~

~~In 2019 new sediment cores have been retrieved from Holzmaar to compile the new record HZM19 (see chapter 3.1, Fig. 2). The lithological description of HZM19 follows the characterization of Zolitschka (1998a, 1998b), dividing the HZM84-B/C profile into 12 lithozones (H1 – H12). We added the sediment colours found in HZM19 to this description.~~

Except H1, all lithozones cover finely-laminated diatomaceous gyttja with varying minerogenic and organic content ~~and as well as~~ colour. All lithozone depths are summarized in Table A14. The transition from light greenish grey (10Y 8/1) and greyish brown (2.5Y 5/2) minerogenic, finely laminated, weakly carbonaceous silts and clays in H1 (12.9 – 14.6 m) to carbonaceous laminated gyttja in light olive brown (2.5Y 5/3), black (10YR 2/1) and light-yellow brown (2.5Y 6/3) with slightly higher organic content in H2 (11.3 – 12.9 m) indicates the transition from the Pleniglacial to the Late Glacial (Fig. 2).

Within H2, the distinct and almost 20 cm thick coarse-grained tephra from the Laacher See eruption (LST, 11.5 – 11.7 m) is deposited, a well-dated isochrone (Reinig et al., 2021) of European lake sediments (Fig. 2). The following lithozone H3 (10.9 – 11.3 m) shows a high minerogenic content and almost no organic components with colours of light greenish grey (5GY 7/1) and grey brown (10YR 5/2), representing the YD at the end of the Pleistocene. Unfortunately, almost one third (12.9 cm) of the YD lithozone H3 is missing due to a technical gap (Fig. 2).

The Holocene sediment shows a periodic change from sections with higher organic content in black (2.5Y 2.5/1) and light olive brown (2.5Y 5/3) (H4: 10.7 – 10.9 m, H6: 9.9 – 10.0 m) to sections

**Formatiert:** Mit Gliederung + Ebene: 2 + Nummerierungsformatvorlage: 1, 2, 3, ... + Beginnen bei: 1 + Ausrichtung: Links + Ausgerichtet an: 1.88 cm + Einzug bei: 3.15 cm

with high organic and clastic content in slightly brighter colours like grey (10YR 5/1) (H5: 10.0 – 10.7 m, H7: 9.3 – 9.9 m). The tephra of the Ulmener Maar eruption (UMT, ca. 3 mm thick) occurs in H5 at 10.24 m. The longest lithozone H8 (5.5 – 9.3 m) contains distinctly varved dark reddish brown (5YR 3/2) sediments with high organic content changing towards the top to very dark greyish brown (10YR 3/2) and brown (10YR 4/3) with several up to 5 mm thick lenses of authigenic vivianite. Also, a low carbonate content was recognized. Furthermore, turbidites are observed more frequently from H8 to the top of HZM19 (Fig. 2).

Above H8, the clastic content increases and brightens up to light olive brown (2.5 Y 5/3) and greyish brown (2.5Y 5/2) hues in H9 (4.3 – 5.5 m). In H10 (3.1 – 4.3 m) colours change to darker hues, e.g. olive grey (5Y 4/2) and black (5Y 2.5/2), while the organic content remains high and terrestrial macrofossils like pieces of wood or leaf remains occur more frequently towards the top. The organic content is decreasing slightly in H11 (1.1 – 3.1 m), which also contains clastic components and terrestrial plant material as well as turbidites with paler colours, e.g. olive brown (2.5Y 5/3) and grey (2.5Y 5/1). The uppermost lithozone H12 (1.1 m to the top of HZM19) shows unconsolidated organic sediment with a homogenous blackish (5Y 2.5/1) colour for the lower part and brighter dark olive grey (5Y 3/2) sediment at the very top (Fig. 2).

## 2.3 Evolution of the Holzmaar varve chronology

Formatiert

### Holzmaar chronology

First varve counts and documentation of the annual origin for the finely laminated sediments preserved in the Holzmaar record were carried out in the late 1980's (Zolitschka, 1990, 1991, 1992), presenting the initial Holocene and Late Glacial varve chronology VT-90. Varve Time (VT) refers to varve (calendar) years before 1950 CE (Common Era), which is equivalent to the commonly used reference timescale for radiocarbon dates provided in cal. BP (calibrated years before present, i.e. 1950 CE). The chronology of VT-90 was elaborated for the HZM84-B/C composite record recovered in 1984 and was counted back to the onset of the Late Glacial, i.e. to 12,794 VT-90. This varve chronology was subsequently extended by counting the deeper, periglacial section back to the Last Glacial Maximum, i.e. to an age of 22,500 VT-90 (Brauer, 1994; Brauer et al., 1994).

By including the new sediment cores of HZM90-E/-F/-H, VT-90 was modified resulting in VT-94. These overlapping sediment-core series as well as all other mentioned cores have been recovered from the deepest part of Holzmaar, i.e. from within the 20-m isobath (Fig. 1). This recounting revealed an underestimation of the youngest 5000 years, for which 555 years have been added. This initial underestimation was mainly caused by sections with very thin varves difficult to count (Zolitschka, 1998b). Another discrepancy occurred within the sediments of the Younger Dryas (YD), for which 245 years had to be added. Altogether, the difference from VT-90 to VT-94

comprises an addition of 800 years, shifting the basal age of the Late Glacial back to 13,594 VT-94 (Zolitschka, 1998b).

To crosscheck the varve chronology with an independent dating method, 41 samples of terrestrial macrofossils along the entire profile (Tab. A2) have been analysed using the AMS (Accelerator Mass Spectrometry) radiocarbon method (Hajdas et al., 1995 and one unpublished radiocarbon date). A comparison between VT-94 and the calibrated radiocarbon chronology shows a discrepancy of +346 years between 3500 and 4500 VT-94 (Hajdas et al., 1995; Hajdas-Skowronek, 1993). This correction factor was estimated by  $\chi^2$ -minimization and added by linear interpolation between 3500 and 4500 VT-94. The outcome was VT-95, which consists of three segments. Segment I is covered by an "absolute" chronology until 3500 VT-95, while segment II (3500 - 4846 VT-95) was extended based on the discrepancy detected between varve and calibrated radiocarbon chronologies. Segment III covers sediments from 4846 - 13,940 VT-95 and is considered as a floating chronology (Hajdas et al., 1995; Zolitschka, 1998b).

In 1996 new sediment cores (HZM96-4a, -4b) have been obtained from Holzmaar and VT-95 was transferred to this new record using 26 distinct marker layers with their related VT and error. The age-depth model was subsequently obtained by linear interpolation (Baier et al., 2004). At the same time, novel varve counts for the Meerfelder Maar sediment record established 1880 varve years between the two isochrones of Laacher See Tephra (LST, eruption ca 40 km NE from Holzmaar) and Ulmener Maar Tephra (UMT, eruption ca 13 km NE from Holzmaar) (Brauer et al., 1999), which both are also archived in the Holzmaar sediment record. However, this well-constrained time interval was only 1560 years long for the Holzmaar record. The obviously missing 320 years have been positioned and added to VT-95 based on pollen data from Holzmaar (Leroy et al., 2000), assuming a hiatus for the middle part of the YD biozone at 12,025 VT-95. This resulted in the latest version (VT-99) of the Holzmaar varve chronology (Zolitschka et al., 2000) with a basal age of 14,260 VT-99 for the Late Glacial.

Varve quality and error estimations were first discussed and described based on multiple counts of selected and representative thin sections (Zolitschka, 1991). Later, different varve quality classes have been described in more detail for VT-90 (Zolitschka et al., 1992) and for VT-95 (Zolitschka, 1998b) with error estimations in the  $1\sigma$  range (Table A1). Similar error margins were confirmed by counting more recent sediment profiles (HZM96-4a, 4b) from Holzmaar (Prasad and Baier, 2014). In this study, the uppermost part was discussed as showing even higher counting uncertainties. However, no alternative error margins were can be provided for this section. Thus, we use the data of Table A1 for further evaluations.

### 3. Materials and Methods

#### 3.1 Sediment core collection

In August 2019, Holzmaar was revisited and four parallel cores (HBM19-07, HBM19-08, HBM19-10, HBM19-11) have been retrieved from the centre of the lake in 19 m water depth (Fig. 1) using a UWITEC piston-corer with a diameter of 90 mm (HBM19-07, -08, -10) and 60 mm (HBM19-11) from a coring platform. The coring locations are distributed evenly along a 12 m-long transect with 4 to 4.4 m distance between coring locations. The recovered sediment cores have lengths of 2 m (HBM19-07, -08, -10) and 3 m (HBM19-11), which have been split in the field into 1 and 1.5 m-long sections, respectively. In total, HBM19-07 covers a sediment depth of 15.5 m (0-15.5 m), while the other sites provided different depth ranges: HBM19-08 (0.25 – 10 m), HBM19-10 (4 – 14 m) and HBM19-11 (1 – 19 m). The water-sediment interface was perfectly recovered with HBM19-07-01 as the piston stopped 15 cm above the sediment surface. At the GEOPOLAR lab (University of Bremen) the cores have been split in halves lengthwise, photographed and visually described using a Munsell colour chart and according to the description guide-line by Schnurrenberger et al. (2003). Cross correlation of all sediment-core sections was conducted macroscopically using 48 distinct layers (Table A3).

The four parallel cores HBM19-07, -08, -10 and -11 were aligned and correlated to form the composite profile HBM19 (Fig. 2), which includes 24 core sections and reaches to a basal depth of 14.64 m (Table A4). One technical sediment gap exists at a composite depth of 10.90 m. To determine the precise length of this gap, we use core photographs from a previous Holzmaar core (HBM90-H5u) and determined the technical gap with to have a length of 12.9 cm (Fig. A1).

**Formatiert:** Überschrift 1, Links, Zeilenabstand: einfach, Nummerierte Liste + Ebene: 1 + Nummerierungsformatvorlage: 1, 2, 3, ... + Beginnen bei: 1 + Ausrichtung: Links + Ausgerichtet an: 0.63 cm + Einzug bei: 1.27 cm, Vom nächsten Absatz trennen

**Formatiert:** Englisch (Vereinigte Staaten)

**Formatiert:** Links, Zeilenabstand: Mehrere 1.08 ze, Keine Aufzählungen oder Nummerierungen




Figure 2: Composite profile of HZM19 with (from left to right) lithozones H1 to H12 (cf., Table A14), lithological description, core photographs taken immediately after core splitting, positions of marker layers and radiometric samples (cf., Tables A52, A75) and core sections used for the composite profile (cf., Table A43).

## 2.3 3.2 Chronology

### 2.3.1 3.2.1 Evolution of the Holzmaar varve chronology

First varve counts and documentation of the annual origin for the finely laminated sediments preserved in the Holzmaar record were carried out in the late 1980's (Zolitschka, 1990, 1991, 1992), presenting the initial Holocene and Late Glacial varve chronology VT 90. Varve Time (VT) refers to varve (calendar) years before 1950 CE (Common Era), which is equivalent to the commonly used reference timescale for radiocarbon dates provided in cal. BP (calibrated years before present, i.e. 1950 CE). The chronology of VT 90 was elaborated for the HZM84 B/C composite record recovered in 1984 and was counted back to the onset of the Late Glacial, i.e. to 12,794 VT 90. This varve chronology was subsequently extended by counting the deeper, periglacial section back to the Last Glacial Maximum, i.e. to an age of 22,500 VT 90 (Brauer, 1994; Brauer et al., 1994).

By including the new sediment cores of HZM90 E/ F/ H, VT 90 was modified resulting in VT 94. These overlapping sediment core series as well as all other mentioned cores have been recovered from the deepest part of Holzmaar, i.e. from within the 20 m isobath (Fig. 1). This recounting revealed an underestimation of the youngest 5000 years, for which 555 years have been added. This initial underestimation was mainly caused by sections with very thin varves difficult to count (Zolitschka, 1998b). Another discrepancy occurred within the sediments of the Younger Dryas (YD), for which 245 years had to be added. Altogether, the difference from VT 90 to VT 94 comprises an addition of 900 years, shifting the basal age of the Late Glacial back to 12,594 VT 94 (Zolitschka, 1998b).

To crosscheck the varve chronology with an independent dating method, 41 samples of terrestrial macrofossils along the entire profile (Tab. A2) have been analysed using the AMS (Accelerator Mass Spectroscopy) radiocarbon method (Hajdas et al., 1995 and one unpublished radiocarbon date). A comparison between VT 94 and the calibrated radiocarbon chronology shows a discrepancy of +346 years between 3500 and 4500 VT 94 (Hajdas et al., 1995; Hajdas-Skowronek, 1993). This correction factor was estimated by  $\chi^2$  minimization and added by linear interpolation between 3500 and 4500 VT 94. The outcome was VT 95, which consists of three segments. Segment I is covered by an "absolute" chronology until 3500 VT 95, while segment II (3500–4846 VT 95) was extended based on the discrepancy detected between varve and calibrated radiocarbon chronologies. Segment III covers sediments from 4846–12,940 VT 95 and is considered as a floating chronology (Hajdas et al., 1995; Zolitschka, 1998b).

In 1996 new sediment cores (HZM96 4a, 4b) have been obtained from Holzmaar and VT 95 was transferred to this new record using 26 distinct marker layers with their related VT and error. The age depth model was subsequently obtained by linear interpolation (Baier et al., 2004). At the same time, novel varve counts for the Meerfelder Maar sediment record established 1880 varve years between the two isochrones of Laacher See Tephra (LST, eruption ca 40 km NE from Holzmaar) and Ulmener Maar Tephra (UMT, eruption ca 13 km NE from Holzmaar) (Brauer et al., 1999), which both are also archived in the Holzmaar sediment record. However, this well constrained time interval was only 1560 years long for the Holzmaar record. The obviously missing 320 years have been positioned and added to VT 95 based on pollen data from Holzmaar (Leroy et al., 2000), assuming a hiatus for the middle part of the YD biozone at 12,025 VT 95. This results in the latest version (VT 99) of the Holzmaar varve chronology (Zolitschka et al., 2000) with a basal age of 14,260 VT 99 for the Late Glacial.

Varve quality and error estimations were first discussed and described based on multiple counts of selected and representative thin sections (Zolitschka, 1991). Later, different varve quality

**Formatiert:** Schriftart: (Standard) +Überschriften (Cambria), 14 Pt., Fett, Schriftfarbe: Text 1, Englisch (Vereinigte Staaten)

**Formatiert:** Links, Zeilenabstand: Mehrere 1.08 ze, Keine Aufzählungen oder Nummerierungen

**Formatiert:** Englisch (Vereinigte Staaten)

**Formatiert:** Überschrift 3, Links, Zeilenabstand: einfach, Keine Aufzählungen oder Nummerierungen

classes have been described in more detail for VT 99 (Zolitschka et al., 1992) and for VT 95 (Zolitschka, 1998b) with error estimations in the 1 $\sigma$  range (Table A1). Similar error margins were confirmed by counting more recent sediment profiles (H2M96 4a, 4b) from Holmaar (Prasad and Baier, 2014). In this study, the uppermost part was discussed as showing even higher counting uncertainties. However, no alternative error margins were provided for this section. Thus, we use the data of Table A1 for further evaluations.

### 2.3.2 Transfer of VT 99 to H2M19

The varve chronology VT 99 (Zolitschka et al., 2000) was transferred to H2M19 by using predefined marker layers and 41 radiocarbon sampling positions analysed by Hajdas et al. (1995, 2000) with their specific VT 99 ages and errors (Tables A1, A2). Both, marker layers and radiocarbon sampling positions have been identified and justified by comparison with documents describing the samples as well as core photographs from previous studies and sediment profiles, such as H2M90 E, F, H and H2M96 4a, 4b. All marker layers cover an age range from 141 to 14,158 VT 99. After assignment, the ages of the marker layers have been linearly interpolated and cumulative counting errors were calculated based on the 1 $\sigma$  errors provided with Table A1.

### 2.3.3 Pb-210 and Cs-137 dating

The isotopes Pb-210 and Cs-137 have been used to radiometrically date the uppermost part of H2M19 at the University of Gdansk. In total, 61 samples were taken with a thickness of 2 cm. The activity of Cs-137 was determined directly by gamma-ray spectrometry from freeze-dried and homogenized samples. Gamma measurements were carried out using a HPGe well-type detector (GCW 2021) with a relative efficiency of 27% and full width at half maximum (FWHM) of 1.9 at the energy of 1333 keV (Canberra). Energy and efficiency calibration were done using reference material CBSS-2 (Eurostandard CZ) in the same measurement geometry like the samples. The counting time for each sediment sample was 24 hours.

Activity of total Pb-210 was determined indirectly by measuring Po-210 using alpha spectrometry. Dry and homogenized sediment samples of 0.2 g were spiked with a Po-209 yield tracer and digested with concentrated HNO<sub>3</sub>, HClO<sub>4</sub> and HF at a temperature of 100 °C using a CEM Mars 6 microwave digestion system. The solution obtained was evaporated with 6M HCl to dryness and then dissolved in 0.5M HCl. Polonium isotopes were spontaneously deposited within four hours on silver discs. Activities were measured using a 7200-04 APEX Alpha Analyst integrated alpha-spectroscopy system (Canberra) equipped with PIPS A450-18AM detectors. Samples were counted for 24 hours. A certified mixed alpha source (U-234, U-238, Pu-239 and Am-241; SRS 73833-121, Analytics, Atlanta, USA) was used to check the detector counting efficiencies.

**Formatiert:** Standard, Links, Zeilenabstand: einfach, Keine Aufzählungen oder Nummerierungen

**Formatiert:** Überschrift 3, Links, Zeilenabstand: einfach, Keine Aufzählungen oder Nummerierungen

**Formatiert:** Schriftart: (Standard) +Textkörper (Cambria), 11 Pt., Nicht Fett, Schriftfarbe: Automatisch, Englisch (Vereinigte Staaten), Nicht Ausgeblendet

**Formatiert:** Standard, Keine, Abstand Vor: 0 Pt., Keine Aufzählungen oder Nummerierungen, Vom nächsten Absatz trennen, Diesen Absatz nicht zusammenhalten, Nicht Überschriften beim Öffnen standardmäßig reduziert anzeigen

### 2.3.4 3.2.2 Bayesian age-depth modelling

To produce the chronology for HZM19 we test and compare different methods integrating varve counts with radiometric measurements using Bayesian age depth modelling. The advantage of any modelling approach is that all possible calendar ages of calibrated radiocarbon dates and their probability density functions (PDFs) will be tested by using a repeated random sampling method (Blaauw, 2010; Telford et al., 2004). In addition, using the Bayes theorem allows to incorporate information of the accumulation history known prior to modelling. Thus, calendar ages, which are monotonic with depth and with positive accumulation rates in  $\text{yr cm}^{-1}$  (in sedimentological terms, accumulation rates as they are used for Bayesian age depth modelling are equivalent to "sedimentation rates", as corroborated by the units used) are calculated (Lacourse and Gajewski, 2020; Trachsel and Telford, 2017). This is different if compared to the "Classical Age depth Modelling" carried out by CLAM (Blaauw, 2010).

Currently established programs that use Bayesian statistics are Oxcal (Ramsey, 2008), BChron (Haslett and Parnell, 2008) and Bacon (Blaauw and Christen, 2011), all of which differ in terms of parameter settings and handling of outliers. In this study, we focus on varve counting integration methods using Bacon (rBacon version 2.5.7; Blaauw et al., 2021; Blaauw and Christen, 2011) for the R programming language (version 4.1.1; R Core Team, 2021). Bacon uses a Markov Chain Monte Carlo (MCMC) sampling strategy to model the accumulation history piecewise using a gamma autoregressive semi-parametric model (Blaauw and Christen, 2011). The accumulation rate of each segment depends on the accumulation rate of the previous segment. Dates are treated using a student's *t* distribution. Although Bacon provides default values, the accumulation rate is controlled by two adjustable prior distributions (prior model), the accumulation rate as a gamma distribution and the memory, which describes the dependence of accumulation rates between neighbouring depths as a beta distribution. Both latter parameters are defined by a shape and a strength prior, respectively, in addition to a mean prior. Furthermore, we make use of the number of segments (thick parameter) recommended by Bacon. The program also allows to incorporate information about hiatus and slump events in the profile.

Only few studies use the Bayesian approach that integrates varve counting information with radiocarbon dates (Bonk et al., 2021; Vandergoes et al., 2018; Shanahan et al., 2012; Fortin et al., 2019). We extracted three different methods and for comparison include one model only with radiocarbon data, i.e. excluding any VT-99 information. Thus, four different age-depth models (A-D) are compared and discussed:

A) Model based only on radiocarbon dates.

B) This parameter-based varve integration method introduced by Vandergoes et al. (2018) compares several varve integration techniques for sediments from Lake Ohau (New Zealand)

Formatiert: Überschrift 3, Links, Zeilenabstand: einfach, Keine Aufzählungen oder Nummerierungen



using both OxCal and Bacon. Here, we select the integration approach with Bacon, where the “varve counts function” is the source for the prior-parameter of mean accumulation rate. Major changes in accumulation history recorded by the varve data are derived by using the R package “segmented” (Muggeo, 2022). It dissects the sediment sequence and for each resulting segment an individual mean accumulation-rate prior is defined.

C) The tie point-based integration used by Shanahan et al. (2012) integrates the varve chronology from Lake Bosumtwi (Ghana) based on certain tie points with normally distributed age uncertainties of the cumulative error. They address the problem of integrating all individual varve counts, as they cannot be considered as independent chronological datapoints. Thus, they would be weighted too strongly in the model. The compromise we have chosen for this study, is placing one varve tie-point every 100 years. As there is no varve counting available for HZM19 but VT-99 ages based on marker layers, we implement them with cumulative errors as tie points instead.

D) The segmented and parameter-based integration introduced by Bonk et al. (2021) provides the most complex method for varve integration. The problem of not or poorly varved sections in the sediment profile of Lake Goszcz (Poland) is compensated by dividing the profile into three sections and interpolating the section with low-quality varves using Bayesian modelling. For the Holzmaar record, we define four sections: sections 2 and 4 are based on Bayesian modelling, while sections 1 and 3 rely on VT-99. Section 3 is treated as a floating chronology and placed based on the sum of calibrated radiocarbon probabilities lying within this section. To tighten the two Bayesian modelled sections to the following varved sections, an anchor tie-point based on the oldest age of the younger sections is implemented.

For each model we use [varve chronology published by Hajdas et al., 1995](#) and Hajdas et al., 2000 (Table 5), the calibration curve IntCal20 (Reimer et al., 2020) and make use of the default accumulation strength and memory priors. We also implement a surface age of  $-69 \pm 1$  cal. BP as tie point with a normal distributed error to anchor the chronology to present-day.

## 4. Results and Interpretation

### 4.1 Transfer of VT-99 to HZM19

The varve chronology VT-99 (Zolitschka et al., 2000) was transferred to HZM19 by using predefined marker layers and 41 radiocarbon sampling positions analysed by Hajdas et al. (1995, 2000) (Fig. A2) with their specific VT-99 ages and errors (Tables A2, A5). Both, marker layers and radiocarbon sampling positions have been identified and justified by comparison with documents describing the samples as well as core photographs from previous studies and sediment profiles, such as HZM90-E, -F, -H and HZM96-4a, 4b. All marker layers cover an age range from 141 to 14,158 VT-99. After assignment, the ages of these marker layers have been linearly

**Formatiert:** Links, Einzug: Links: 0.63 cm, Zeilenabstand: Mehrere 1.08 ze, Keine Aufzählungen oder Nummerierungen

**Formatiert:** Englisch (Vereinigte Staaten)

interpolated and cumulative counting errors were calculated based on the 1 $\sigma$  errors provided with Table A21.

All 84 marker layers distribute in HZM19 from 1.16 - 12.93 m and cover the entire VT-99 age range ~~from 141 to 14,158 VT-99~~ (Table A52). During the transfer of marker layers to HZM19 and comparison between HZM19 and previous Holzmaar sediment cores (HZM84-B/C, HZM92-E/-F/-H, HZM96-4a/4b) differences in position of the lowermost marker layers occurred (Fig. ~~ure~~ A2). All records show differences in distances between marker layers (ML) 1 (14,156 VT-99), ML-2 (14,152 VT-99) and ML-3 (13,646 VT-99) making a clear assignment of these layers difficult. Thus, we excluded these three marker layers for the transfer of VT-99 to HZM19. The lowermost applied marker layer is therefore ML-4 with a varve age of 13,087 VT-99 at a depth of 11.86 m. Because of inconsistencies in documentation, we excluded two more VT-99 ages, i.e. those related to the radiocarbon ages HZM-46 and HZM-10.1 (Table A52).

The marker layer density reaches a mean value of 5.5 dpm (dates per millennium) being most frequent before 10,000 and after 6000 cal. BP (Fig. 34). We use a linear interpolation to receive an age-depth model based only on VT-99 with a resulting accuracy of 282 years as a mean age range and a maximum age range of 744 years (Table A6).

The radiocarbon dating density of HZM reaches an overall mean value of 2.7 dpm (Fig. 34), which is 35% higher than the 2 dpm recommended for Bayesian modelling by Blaauw et al. (2018). However, their distribution is uneven. Radiocarbon dates are most frequent for ages >10,000 cal. BP with 3-7 dpm (mean: 5 dpm) (Fig. 34). A minimum density of radiocarbon dates (0-1 dpm) is obtained from 10,000-6000 cal. BP (mean: 0.5 dpm). Therefore, a chronology based on the available radiocarbon data within this section should be interpreted with caution. Dating density for the uppermost 6000 years is higher and varies between 1 and 4 dpm (mean: 2.2 dpm).

When we compare VT-99 with radiocarbon ages calibrated with the latest calibration curve IntCal20 (Reimer et al., 2020), an overall agreement with marker layers is observed. Only for the lowermost part below approximately 10.64 m, we observe an increasing underestimation of VT-99 in relation to IntCal20 calibrated radiocarbon ages (Fig. A32, Table A52). This was already observed by Hajdas et al. (2000) in comparison to Intcal98 but has not been corrected yet.




Figure 3: Number of dating points per millennium (dpm) of HZM19 for marker layers (n: 84, mean: 5.5 dpm) and radiocarbon dates (n: 41, mean: 2.7 dpm). Red dotted line marks the recommended threshold of 2 dpm for Bayesian modelling suggested by Blaauw et al. (2018). Surface age and three ages estimated by Cs-137 are excluded.

### 3.

## 3.2 — 4.2 Chronology

### New chronological information

#### 3.2.1 4.2.1 Pb-210 and Cs-137 dating

The profile of unsupported Pb-210 activity concentration shows a gradual rather than an exponential decrease within the first meter of HZM19 (Fig. 43). Additionally, a plateau from 8 to 30 cm is interpreted as a section with rapid deposition of homogenous material and will be treated for further analyses as a slump event. Despite this irregularity, the gradual decrease in unsupported Pb-210 activity with depth indicates high sedimentation rates. We use the CFCS (Constant Flux Constant Sedimentation) model to estimate mean sedimentation rates of  $1.09 \pm 0.13 \text{ cm yr}^{-1}$ . This value should be treated with caution but suggests that the uppermost meter (including a 22 cm-thick slump) was deposited in ca. 70 years.

**Formatiert:** Standard, Links, Zeilenabstand: einfach, Keine Aufzählungen oder Nummerierungen

**Formatiert:** Links, Einzug: Links: 1.25 cm, Zeilenabstand: Mehrere 1.08 ze, Keine Aufzählungen oder Nummerierungen

**Formatiert:** Einzug: Links: 1.25 cm

**Formatiert:** Links, Einzug: Links: 1.25 cm, Zeilenabstand: Mehrere 1.08 ze, Keine Aufzählungen oder Nummerierungen

**Formatiert:** Block




Figure 43: Results of unsupported Pb-210 (left) and Cs-137 (right) measurements with error bars for the uppermost 110 cm of HZM19. Shaded areas indicate the plateau shown by Pb-210 data, black arrows mark peaks assigned to radiochronological events (given numbers are ages in years CE).

The variability of Cs-137 activity concentrations delivers potentially three historical markers (Fig. 43). The Cs-137 profile is smooth lacking sharp peaks due to high sedimentation rates and likely sediment focusing. First traces of Cs-137 are recognizable at 101.2 cm and indicate atomic bomb testing in the early 1950's. At 89.2 cm, there is a significant increase signalling atmospheric fallout in the early 1960's in response to peak atomic bomb testing. Finally, at 69.2 cm a strong increase in Cs-137 documents the 1986 Chernobyl accident (Fig. 43, Table A75). This interpretation is generally in line with the results of Pb-210 dating. The shape of the Cs-137 record also corresponds nicely to the results of Sirocko et al. (2013), who measured Cs-137 on sediments from Schalkenmehrener Maar and Ulmener Maar (both WEVF). For both of these cases, the 1986 Chernobyl peak is also much larger than the one related to the start of atomic bomb tests in 1963.

### 3.2.2 4.3 Age-depth modelling

Four different Bayesian age-depth models are calculated, of which three include varve ages (Model B-D) and one only radiocarbon ages (Model A). In common for all model runs are the default memory priors and the use of the IntCal20 calibration curve (Reimer et al., 2020). Furthermore, based on the Pb-210 and Cs-137 dating analysis, a slump at a composite depth of 8-30 cm was implemented, as well as the LST from 11.52 – 11.71 m. As known from previous varve

Formatiert: Links, Einzug: Links: 0.63 cm, Zeilenabstand: Mehrere 1.08 ze, Keine Aufzählungen oder Nummerierungen

and pollen studies of the Holzmaar record (Brauer et al., 1999; Leroy et al., 2000), 320 years are missing during the YD and have been included into VT-99 at 12,025 VT-99. Based on the study of Leroy et al. (2000), we were able to locate the position of the YD hiatus to a depth of 11.09 m, which we implemented for each model with a maximum duration of 320 years. In addition to marker layers and radiocarbon dates, we included the surface age of  $-69 \pm 1$  cal. BP and three events dated by Cs-137 (Table A75).

Preliminary test runs reveal two necessary changes to be made for the calculations: 1) The default number of iterations is too low to produce a robust model for the entire HZM19 sediment sequence. Thus, we use the *Baconvergence()*-function of Bacon to estimate the number of iterations needed. This function repeats the calculations and tests if the MCMC mixing of the core results in a robust model by calculating the “Gelman and Rubin Reduction Factor” (Brooks and Gelman, 1998). Good mixing is indicated by a threshold of  $<1.05$ , which in our case was reached after three iterations when the number of iterations was increased to 40,000. This results in a better mix of MCMC iterations but also in long calculation times ( $> 5$  hours). 2) For each test run, Bacon predicted ages consistently too old for the LST, which is probably caused by slightly too old ages of the surrounding radiocarbon dates (Table A52). To gain a better comparability with studies from other sites, we decided to include the latest LST age of  $13,006 \pm 9$  cal. BP (Reinig et al., 2021, Table A75).

In addition, we extended the age-depth model to a maximum depth of 14.64 m, as ongoing analyses exceed the lowermost dated level. However, in the following chapters we only discuss the model output between the first (ML36/1) and the last (HZM-19) marker layer at 12.93 m (Table A52) and compare it with the interpolated varve chronology (VT-99).

After each calculation and if the Bacon output indicates a highly variable log of objectives or MCMC iterations, we made use of the *scissor()*-command to achieve a better mixing of the output. All Bacon model outputs with their settings and additional information are shown in Figure 5A3 and related ages are listed in Table A6.

The **model without varve integration (Model A)** is based on the year of sediment recovery (surface age), three dates estimated by Cs-137 analyses, the age for the LST (Reinig et al., 2021) and 41 calibrated radiocarbon probability density functions (Fig. 5A3A). Different to Hajdas et al. (1995), this model includes the outlier of HZM-23, but excludes HZM-24 and other described outliers (Table A52).




Figure 5: Bacon output for Model A, B, C and D (sections 2 and 4). Each output with indicator panels from top left to right: MCMC iterations, prior (green) and posterior (grey) for accumulation rate distribution, memory and hiatus with defined settings in red. Main panel: model with calibrated radiocarbon date probabilities (blue), tie-points with normal distribution (orange) and the posterior age-depth model with mean (red dotted line) and 95% confidence intervals (gray dotted line). Vertical gray lines (from left to right): slump event, defined hiatus and Laacher See Tephra. In additional panels of Models B and D2 boundaries indicating major changes in accumulation rate are provided as vertical dotted lines.

Model A results in an age of 14,615 [minimum: 14,339, maximum: 14,926] cal. BP at the lowermost dated depth of 12.93 m with a mean age uncertainty of 468 yrs. The maximum age uncertainty of approx. 1056 years occurs at a depth of 8.86 m within lithozone H8 (Table A6), where radiocarbon dating density is <1 dpm (Fig. 34).

The **parameter-based integration (Model B)** integrates VT-99 using all dates as in Model A and adjusts the prior information given for the calculation based on the varve accumulation-history. We follow the procedure presented by Vandergoes et al. (2018) and calculate a breakpoint based on ages and depths of the marker layers at 4.43 m, i.e. at 1312 VT-99 (Fig. 5A3B). This boundary is implemented as an additional hiatus to the Bacon code with a duration of 1 year. The accumulation rate prior is set based on published sedimentation rates (Zolitschka et al., 2000). We calculate with a mean of 0.49 yr/mm for the uppermost part (71-1312 VT-99), with 1.30 yr/mm from 1312 to the YD hiatus at 12,025 VT-99 and with 0.76 yr/mm from the YD hiatus to the lowermost age of 14,158 VT-99. Model B is calculated using the same parameters as for Model A and with the same treatment of outliers.

The resulting posterior model shows similarities to Model A, having a maximum mean age of 14,456 [min.: 14,236, max.: 14,749] cal. BP at a depth of 12.93 m and a mean 95% confidence interval of 456 years with a maximum of 1064 years at 8.78 m, i.e. within the period of lowest radiocarbon dating density (Fig. 34).

The **tie point-based integration (Model C)** is based on the approach used by Shanahan et al. (2012). We include 43 marker layers with related VT-99 ages and cumulative errors as normal distributed tie points into the model, which adds to the dates used in Models A and B and sums up to 89 dates. This approach increases the amount of chronological information and fills areas with larger gaps between radiocarbon dates. The model was run with default settings provided by Bacon (Fig. 5A3C). Bacon recognizes the outliers in the same way as by previously described models.

Model C results in a maximum age of 14,614 [min.: 14,332, max.: 14,919] cal. BP (at 12.93 m) with a mean 95% confidence interval of 329 years, which is better than for Models A and B. A maximum age range of 749 years is given at a depth of 9.18 m, which is also slightly better than for previously presented models. However, Model C produced MCMC iterations with highest noise and it was difficult to cut out a well-mixed section (Fig. 5A3C, upper left panel).

The **segmented and parameter-based integration (Model D)** is a more complex method of varve integration used by Bonk et al. (2021) and was adapted for the HZM19 profile by dividing the varve chronology of VT-99 into four sections. This separation is based on variations of

counting uncertainty, radiocarbon sampling density and an increasing offset of VT-99 to the latest calibration curve IntCal20 (Fig. A32).

Section 1 (0 – 5.98 m) and Section 3 (6.70 – 9.90 m) are transferred and interpolated based on VT-99 marker layers, as they are consistent with calibrated radiocarbon data (Section 1) and have well-preserved varves with small counting errors of  $\pm 0.7\%$  (Section 3). Section 2 (5.98 – 6.70 m) and Section 4 (9.9 – 14.6 m) are reported as showing higher counting uncertainties (Section 2) or increasing differences between VT-99 and the calibration curve (Section 4). Thus, we replace the varve chronology in Sections 2 and 4 with Bayesian age-depth modelling (Fig. 5A3D). Section 4 also contains very dense radiocarbon dates (Hajdas et al., 2000), which increase the predictability of Bacon (Fig. 34).

Section 1 is based on linear interpolation for ages of the sediment surface ( $-69 \pm 1$  cal. BP), three dates derived by Cs-137 analyses (Table A75) and 25 ages of marker layers with a basal age of  $3704 \pm 134$  cal. BP at the position of HZM-25 (Table A52).

The modelled Section 2, previously identified as a section with sedimentation rates  $> 2.86$  yr/mm and therefore a source of high counting uncertainties and underestimation of varve ages (Zolitschka et al., 2000), consists of five radiocarbon dates (Table A52) and the basal age of Section 1 ( $3704 \pm 134$  cal. BP) as anchor point for Section 2. To reduce the resulting gap between first and second sections, we reduce the error estimation for the anchor point to  $\pm 70$  years ( $\pm 0.5\sigma$ ). As there is a major change in sedimentation rates within this section, we calculated a boundary similar as in Model B using the marker layers of this section (Fig. 5A3D). This allows defining a boundary at the depth of 6.29 m with adjusted accumulation means of  $3.33$  yr/mm above (5.98 – 6.29 m) and  $1.59$  yr/mm below (6.29 – 6.70 m), using published sedimentation rate data (Zolitschka et al., 2000). Based on suggestions by the software, the “thick”-parameter was set to 4 mm. The resulting model covers an age range from 3709 [min.: 3591, max.: 3825] to 5419 [min.: 5329, max.: 5548] cal. BP (Fig. 5A3D section 2).

Section 3 interpolates 16 marker layers (Table A52), which are treated as a floating chronology. The placement of the anchor point relates to the basal age of the lowermost calibrated radiocarbon date (HZM-4.3) in Section 2 (Table A52) and the maximum sum of the four calibrated radiocarbon PDFs within this part with a summed probability of 0.076 at  $5450 \pm 165$  cal. BP (Fig. A4-A).






Figure 6: Age-depth model for HZM19 based on Model D with Sections 1 and 3 based on VT-99 (section numbers at the right) and Sections 2 and 4 based on Bayesian modelling (shaded).

In comparison to the original VT-99 this approach results in a shift of +65 years for all marker layers within Section 3 (Fig. A4-B). Thus, a basal age of  $10,619 \pm 213$  cal. BP is obtained for Section 3.

The basal age of Section 3 is implemented as the anchor tie-point for the Bacon calculation of Section 4 with a reduced error of 100 years to tighten both sections closer to each other. In

addition to the difficulties based on missing sediment within the YD, this section is the source of highest counting uncertainties for VT-99. Section 4 is based on 25 radiocarbon dates and the latest age estimation for the LST (Table A52). As in Section 2, we adjusted the sedimentation rate prior (= 0.94 yr/mm) based on VT-99 accumulation rate data (Zolitschka et al., 2000). The Bacon software suggests a segment length of 30 mm that we applied. The resulting model covers an age range from 10,663 [min.: 10,457, max.: 10,864] to 14,485 [min.: 14,287, max.: 14,721] cal. BP at 12.93 m (Fig. 5A3D, Section 4).

If all sections are merged, the continuous age-depth relationship forming Model D (Fig. 65) consists of 63% VT-99 ages and 37% Bacon modelled ages with in total 80 missing years between the sections, as it is not possible to determine the exact start and end age of the models. This segmented and parameter-based integration model results in a maximum age of 14,485 [min.: 14,287, max.: 14,721] cal. BP (at 12.93 m) with a mean age uncertainty of 229 years, which is the smallest of all four tested models. The maximum age range is 447 years at 11.09 m depth and thus considerably smaller compared to those of Models A to C (Table A6).

#### **2.1.1 4.4 Comparison of model output with VT-99**

The comparison of all presented models differs in means and accuracies of predicted ages along the core (Fig. 76A1; B1; C1; D1), which becomes more evident in comparison with VT-99 (Fig. 76A2,3; B2,3; C2,3; D2,3). These differences in mean modelled age and mean VT-99 age vary in direction and amplitude (Fig. 76A2; B2; C2; D2). The largest age differences during the Holocene occur in Model A and B with up to 300 years between 4 and 6 m depth (Fig. 76A2; B2). The defined boundary in Model B results in large differences within the boundary area, predicting much younger ages than VT-99. Due to the small cumulative counting uncertainty of VT-99 in the upper part of the profile, the mean of Model B outranges the VT-99 error in most sections above 6 m (Fig. 76B2).

**Formatiert:** Einzug: Links: 0.63 cm, Keine Aufzählungen oder Nummerierungen




Figure 7: Results of Models A, B, C, D plotted against composite depth (1), compared to VT-99 as the difference of mean ages (Model mean - VT-99 mean) (2) and plotted vs. VT-99 confidence intervals (CI) (3).

The approach used for Model C reduces the difference between VT-99 and the model, probably a result of increased dating density (Fig. 76C2). This approach also leads to less over- and underestimations of the model's mean age and the VT-99 age range (Fig. 76C2). Only the segmented insertion of VT-99 in model D results in comparable ages during the Holocene (Fig. 76D2)

In the Late Glacial part below 11 m, all models produce ages constantly older than VT-99 (Fig. 76A2, B2, C2, D2). The age differences are even higher (up to 477 years), when the Bacon prior for accumulation rates was not adjusted to VT-99 (Fig. 76A2, C2). In the other cases the maximum age differences are 369 and 354 years for Model B and D, respectively (Fig. 76B2, D2). Hajdas et al. (2000) already observed a shift between the varve ages of radiocarbon dated samples and

**Formatiert:** Beschriftung, Zeilenabstand: einfach, Vom nächsten Absatz trennen

calibrated ages using the INTCAL98 calibration curve (Stuiver et al., 1998) and discuss the difference using the LST age estimation from Meerfelder Maar (12,880 VT). However, no adjustment has been made to fit the VT-99 ages to the calibration curve. With the LST dated to 13,006  $\pm$  9 cal. BP (Reinig et al., 2021) and the use of the INTCAL20 calibration curve, an underestimation of VT-99 compared to the calibration curve is still existing (Fig. A32). Therefore, a correction of ages older than 12,800 cal. BP is needed to ensure comparability of HZM19 to other sites.

In order to find the best method to transfer VT-99 to HZM19 and to improve the chronology by using Bayesian modelling, a closer look to each model's accuracy is necessary (Fig. 76A3, B3, C3, D3). In comparison to the cumulative VT-99 counting error, Models A and B show maximum differences in age uncertainties up to +655 and +665 years, respectively (Table A6). Especially below 9.82 m, both models predict ages with larger uncertainties than the estimated counting error for VT-99, particularly with increasing distance to radiocarbon dated levels. Therefore, no improvement in accuracy of age estimations is observed when using the parameter-based approach (Model B).

The tie point-based Model C also predicts larger uncertainties than VT-99 below 9.82 m (Fig. 76C), whereas the overall difference of the age range is reduced to a mean of 47 years with a maximum of +401 years (Table A6). Only the segmented and parameter-based Model D shows no significantly enlarged age uncertainties and an overall improved mean age range as it adapts the cumulative error of the varve chronology in Sections 1 and 3 (Table A6). The overall improvement occurs in Sections 2 and 4, which is the result of more detailed prior settings for the model run. However, all age models result in more accurate age estimations in the Late Glacial part, where the cumulative counting error is higher and radiocarbon dating sampling is dense. But still we see that Models C and D perform best within this section, as they predict ages with constantly lower uncertainty ranges than VT-99. This is in contrast to the other models, which show increased and therefore larger uncertainties at a depth of ca. 11 m. As we calculate this section in Model D with the same data like for Model A and B, we assume that the better adjustment of the sedimentation rate mean prior of Model D influences the model's accuracy. In terms of accuracy, there are no general improvements in calculating a single model for the entire record, but improvements are realised by adjusting the priors in a more detailed way.

### 3.2.3 4.5 Comparison of model output with the common isochrones UMT and LST and the YD biozone

The tephra layers of UMT and LST have been identified for sediments from Holzmaar and Meerfelder Maar (Brauer et al., 1999). The varve age of 11,000 VT-99 for UMT was derived from the Holzmaar chronology (Zolitschka, 1998b), while the YD hiatus of this site did not allow any calendar-year estimation for LST. As no such hiatus exists between these two isochrones at Meerfelder Maar, the age for the LST was derived as 1880 varve years older than UMT, i.e. as 12,880 VT-99. A recent study presents a new and 126 years older age for the LST (Reinig et al., 2021). This age of 13,006 cal. BP was implemented for the calculation of Models A-D.

When we compare all models, the age estimations for UMT and LST are close to the published ages with the UMT dated ca. 20-50 years earlier and thus matches well within the 95% confidence interval (Fig. 8A5, Table A6). Due to the new age of LST, the distances between both isochrones vary from 2030 (Model D) to 2057 (Model C) years, which is 150-177 years more than counted for Meerfelder Maar (Fig. 8A5).

The main differences occur in prediction of the end of the YD that defines the transition to the Holocene. The rapid cooling and subsequent warming left behind easy to recognize traces in many European lake records increasing the comparability between sites. The entire YD is not covered by HZM19 due to a technical gap. Nevertheless, we are able to estimate depth and time range based on detailed pollen investigations (Leroy et al., 2000). Using VT-99, Leroy et al. (2000) date the onset of the YD, i.e. the Allerød/Younger Dryas transition (AL/YD) to 12,606 VT-99 and the Younger Dryas/Preboreal (YD/PB) transition to 11,632 VT-99 with a 320 years hiatus at 12,025 VT-99. For HZM19 these boundaries occur at 10.88 m (YD/PB), at 11.26 m (AL/YD) and the hiatus at 11.11 m (Fig. 76, 8A5).

All model runs predict a YD duration in the range of 1012 (Model C) to 1073 (Model D) years, which is longer than the 974 years given by VT-99 (Fig. 8A5, Table A6). However, the predicted times are closer to its duration counted for Meerfelder Maar (1080 years) (Brauer et al., 1999) or the even longer time spans detected for Lake Gosciadz (1150 years) (Bonk et al., 2021).

Moreover, both onset and the end of the YD transitions have been predicted within the 95% confidence interval comparable to VT-99 (Fig. 8A5, Table A6) and to the Meerfelder Maar record. Only the AL/YD transition varies between 12,694 (Model C) and 12,737 (Model B) cal. BP and, thus, is predicted earlier than for VT-99 (12,606 VT-99). However, this age range still covers the age estimations from Lake Gosciadz [12,620 [min.: 12,389, max.: 12,753] cal. BP] and Meerfelder Maar [12,680 [min.: 12,640, max.: 12,720] cal. BP] (Fig. 8A5). In difference, the YD/PB transition varies between 11,655 (Model D) and 11,723 (Model B) cal. BP, which is slightly earlier than

Formatiert: Keine Aufzählungen oder Nummerierungen

estimated by Meerfelder Maar (11,600 [min.: 11,570, max.: 11,630] cal. BP) and much earlier than the age estimation for Lake Gosciarz (11,470 [min.: 11,264, max.: 11,596] cal. BP) (Fig. 8A5). These discrepancies between the boundaries of the YD biozone obtained by VT-99 and those obtained by the model runs are probably related to the new and 126-year older age for the LST, which is included with all models. Thus, age discrepancies are attenuating towards the UMT with 110 years at the AL/YD transition and 57 years at the YD/PB transition (Fig. 8A5).

#### 4.5. Evaluation of the different varve integration techniques

All models predict convincing age estimations for the isochrones of LST and UMT, whereas the prediction of the YD between both isochrones remains somewhat ambiguous, due to a documented hiatus and too few radiocarbon ages being available for this biozone.

In terms of accuracy and precision, the varve-integration technique applied in Model D, introduced by Bonk et al. (2021), results in most convincing age estimations for HZM19. Especially in terms of accuracy, none of the completely Bayesian modelled age-depth relationships improved the small age uncertainties of VT-99 in the upper part. Only in sections with markedly higher radiocarbon sampling density or in sections with high varve counting uncertainty the Bacon models perform better and result in more accurate age estimations than VT-99.

In comparison, Model B shows nearly no improvement over the approach without varve integration (Model A). The reason is probably the low-resolution definition of sedimentation rate changes (boundaries) for HZM19, which does not reflect the complex accumulation history. Also Vandergoes et al. (2018) reject this integration model. We suggest that this form of varve integration is more useful for less complex and for shorter sediment profiles records.

Better results are observed applying Model C, which is actually the easiest to apply. The accuracy is improved compared to Models A and B as the dating density increases significantly. Based on the Bayesian approach, this leads to smaller age ranges as higher uncertainties occur with increasing distances to dated levels. The resulting mean age is more constrained by VT-99. The accuracy might be improved by additional adjustments of the sedimentation-rate prior (here: based on VT-99). However, varve ages inserted as tie points are included with normal distribution. Therefore, they should not be interpreted as independent measurements with not a normally distributed PDFs. Bayesian statistics could weight tie points too much when they are included too densely. Therefore, this approach should be interpreted with care.

The best result in precision and especially accuracy is achieved by the segmented and parameter-based Model D. This approach is the most challenging, but and makes advantage of both, the high

accuracy of varve counting and the Bayesian approach for densely radiocarbon dated sections. The main difference to the other models is that Model D replaces the sections of lower dating accuracy with modelled ages that incorporate varve information and radiocarbon measurements, which result in a much better performance.

For upcoming geochemical and geophysical studies of the HZM19 record, we will use Model D. As parts of VT-99 (63%) are included in the new chronology, we will refer to it as chronology “VT-22”, which delivers highly accurate age estimations for each depth of the [Holocene](#) sediment profile HZM19. Altogether, this will improve the comparability of the Holzmaar record with other sites.

## **5.6. Conclusion**

As limnogeological and varve studies proceed, new techniques for sediment analysis develop. Thus, previous studies can be improved by reinvestigation. However, many of the previously studied sediment cores are not available for analysis anymore. We expect such cases to happen more frequently in the future. Rarely, the rather time-consuming and expensive chronological studies, especially if the counting of varves is involved, will be funded a second time. This increases the need for finding best ways to adapt varve chronologies obtained during previous studies and to transfer them efficiently and precisely to new sediment cores.

For the well-dated Holzmaar record, we tested three different approaches for the integration of varve counting and radiocarbon dating using Bayesian modelling and applied them to the new composite profile from Holzmaar (HZM19). We conclude that all models result in accurate and precise age estimations. However, with higher dating density and more prior settings used to adjust the Bacon model runs, the model output is enhanced. This is confirmed by results of Model D, which improved and corrected the age estimations considerably. In contrast, Models B and C show nearly no improvement ~~over~~ [compared to](#) VT-99 just like the output of Model A without varve integration.


Multiple varve counting is still one of the best approaches of building a reliable chronology for lacustrine sediment archives. However, the occurrence of hiati or errors in varve counts lead to larger uncertainties with increasing depth that need to be corrected by using independent dating techniques. Therefore, if varve and radiocarbon data are available, like it is the case for Holzmaar, the transfer of both to form a new and integrating chronology is the best option.

For this study of varve integration, we use Bacon. The parameter adjustment of Bacon is complex and especially beginners have problems to understand each single parameter and the effect it has on model results. We compare different models and settings, which helps to decide selecting the best suited approach and to consider the parameters that have to be adjusted. Afterall, we suggest

851 to increase the independent dating density and to adjust prior settings as detailed as possible to  
852 gain a more precise chronology for the varve-integration attempt.

853 Optimizing the Holzmaar chronology is the first step in order to provide a precise and robust age-  
854 depth model for upcoming and high-resolution multi-proxy investigations to unveil all the  
855 environmental details recorded by the varved sediment archive of Holzmaar.

856 **Appendix**



857

858 *Figure A11-54. Determination of the technical gap for HZM19 during the YD. This gap exists between sections HZM19-10-*  
859 *07 and HZM19-08-10 and is bridged by section HZM90-H5u from an earlier [coring campaign](#).*  
860 *[coring campaign](#).*  
861

- Formatiert:** Schriftart: 9 Pt., Kursiv, Schriftfarbe: Text 2
- Feldfunktion geändert**
- Formatiert:** Schriftart: 9 Pt., Kursiv, Schriftfarbe: Text 2
- Feldfunktion geändert**
- Formatiert:** Schriftart: 9 Pt., Kursiv, Schriftfarbe: Text 2, Rechtschreibung und Grammatik prüfen
- Formatiert:** Schriftart: 9 Pt., Kursiv, Schriftfarbe: Text 2








Figure A2: Correlation of HZM84-B/C and HZM19. Positions of marker layers (ML indicated to the left) are marked as solid lines and connected by dotted lines between both profiles. Positions of radiocarbon dates (numbers indicated in rectangular boxes to the right) are marked as solid circles. Grey dotted horizontal lines refer to Cs-137 dated depths. Positions of Ulmener Maar Tephra (UMT), Laacher See Tephra (LST) and the technical gap are indicated.




Figure A32: Radiocarbon ages vs. Intcal98 and Intcal20 calibrated ages between 10,000 and 15,000 cal BP. Black circles show radiocarbon ages from Holzmaar vs. VT-99 age (*reworked samples excluded*). An underestimation of these ages occurs after 12,500 cal BP, where VT-99 seems to be too young.




Figure A4: Calculations for the floating VT-99 chronology of Model D, section 3. A: Calculation of the anchoring age for the varve chronology based on matched and summed calibrated probability density function values of all radiocarbon samples within this section. The maximum summed probability occurs at an anchor age of 5450 cal BP. B: Original VT-99 (black line) vs. floating VT-99 (+65 years, red dotted line) with calibrated radiocarbon samples vs. depth.

**Formatiert:** Schriftart: (Standard) +Textkörper (Cambria), 9 Pt., Nicht Fett, Kursiv, Schriftfarbe: Text 2




Figure A5: Close-up plots for the Lateglacial / Early Holocene transition for Model A, B, C and D with VT-99 mean age (black solid line) and error (shaded in gray) for comparison. Horizontal lines as labelled in (a). Vertical lines refer to the Younger Dryas transitions for each Model (solid lines), while dotted lines refer to mean ages derived by different sites (Lake Gosciaz in blue: Bonk et al., 2021; Meerfelder Maar in red: Brauer et al., 1999).

Formatiert: Nicht vom nächsten Absatz trennen

Formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Englisch (Vereinigtes Königreich)

Feldfunktion geändert

Formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Englisch (Vereinigtes Königreich)

Table A1: Core section and composite depths of lithozones H1 to H12 for HZM19

Lithozone	From		To			Biozone	Human phase
	Section	Section depth [mm]	Composite depth [mm]	Section	Section depth [mm]	Composite depth [mm]	
H12	HZM19_07_01	138	11	HZM19_08_01	700	1057	Last century Middle Ages / Little Ice Age Migration Period / Early Middle Ages Iron Age / Roman Period
H11	HZM19_08_01	700	1058	HZM19_08_03	520	3081	
H10	HZM19_08_03	520	3081	HZM19_08_04	710	4308	
H9	HZM19_08_04	710	4308	HZM19_08_05	750	5535	
H8	HZM19_08_05	750	5535	HZM19_10_05	480	9280	
H7	HZM19_10_05	480	9280	HZM19_11_06	588	9852	
H6	HZM19_11_06	588	9852	HZM19_08_10	140	10025	
H5	HZM19_08_10	140	10025	HZM19_08_10	860	10745	
H4	HZM19_08_10	860	10745	HZM19_08_10	970	10855	
H3	HZM19_08_10	970	10855	HZM19_10_07	300	11258	
H2	HZM19_10_07	300	11258	HZM19_10_08	859	12859	
H1	HZM19_10_08	860	12860	HZM19_07_17	920	14643	

Formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Beschriftung, Zeilenabstand: einfach

Formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Englisch (Vereinigtes Königreich)

890 Table A221: Error (1 sigma) estimations for different varve quality periods for the Holzmaar record (Zolitschka, 1998b),  
891 updated from VT-95 to VT-99.

Varve quality period	VT-99 (duration in years)	Error
A	0 – 2800	±4.0 %
B	2800 – 5300	±2.6 %
C	5300 – 11,600	±0.7 %
D	11,600 – 14,158	±5.9 %
Entire record	0 – 14,158	±2.5 %

892

893

Table A3: List of markers used for correlation of core sections in HZM19

Correlation Marker	Section 1	Section 1 depth (mm)	Section 2	Section 2 depth (mm)
CM 1	HZM19-07-01	970	HZM19-08-01	500
CM 2	HZM19-08-01	755	HZM19-07-02	235
CM 3	HZM19-08-01	795	HZM19-07-02	285
CM 4	HZM19-08-01	935	HZM19-08-02	390
CM 5	HZM19-07-03	230	HZM19-08-02	850
CM 6	HZM19-07-03	585	HZM19-08-03	80
CM 7	HZM19-07-03	665	HZM19-08-03	155
CM 8	HZM19-07-03	925	HZM19-08-03	390
CM 9	HZM19-07-04	30	HZM19-08-03	515
CM 10	HZM19-07-04	45	HZM19-08-03	525
CM 11	HZM19-07-04	85	HZM19-08-03	570
CM 12	HZM19-07-04	155	HZM19-08-03	640
CM 13	HZM19-07-04	255	HZM19-08-03	730
CM 14	HZM19-07-04	800	HZM19-08-04	230
CM 15	HZM19-10-01	90	HZM19-08-04	990
CM 16	HZM19-10-01	290	HZM19-08-05	40
CM 17	HZM19-10-01	320	HZM19-08-05	70
CM 18	HZM19-10-01	340	HZM19-08-05	90
CM 19	HZM19-10-01	410	HZM19-08-05	155
CM 20	HZM19-10-01	670	HZM19-08-05	390
CM 21	HZM19-10-01	925	HZM19-08-05	645
CM 22	HZM19-10-02	60	HZM19-08-05	805
CM 23	HZM19-10-02	180	HZM19-08-05	925
CM 24	HZM19-10-02	245	HZM19-08-05	990
CM 25	HZM19-10-02	570	HZM19-08-06	290
CM 26	HZM19-10-02	815	HZM19-08-06	535
CM 27	HZM19-10-02	990	HZM19-08-06	700
CM 28	HZM19-10-03	10	HZM19-08-06	855
CM 29	HZM19-10-03	790	HZM19-08-07	520
CM 30	HZM19-10-03	830	HZM19-08-07	560
CM 31	HZM19-10-04	180	HZM19-08-07	975
CM 32	HZM19-10-04	320	HZM19-08-08	80
CM 33	HZM19-10-04	425	HZM19-08-08	175
CM 34	HZM19-10-04	505	HZM19-08-08	260
CM 35	HZM19-10-04	870	HZM19-08-08	640
CM 36	HZM19-10-05	150	HZM19-08-08	975
CM 37	HZM19-10-05	850	HZM19-11-06	380
CM 38	HZM19-08-10	135	HZM19-11-06	780
CM 39	HZM19-08-10	180	HZM19-11-06	820
CM 40	HZM19-08-10	355	HZM19-11-06	995
GAP				
CM 41	HZM19-10-07	250	HZM19-11-07	170
CM 42	HZM19-10-07	750	HZM19-11-07	650
CM 43	HZM19-10-07	905	HZM19-11-07	800
CM 44	HZM19-10-07	970	HZM19-11-07	860
CM 45	HZM19-10-08	395	HZM19-11-07	300

Formatiert: Beschriftung, Nicht vom nächsten Absatz trennen



895

<a href="#">CM 46</a>	<a href="#">HZM19-10-08</a>	<a href="#">580</a>	<a href="#">HZM19-11-08</a>	<a href="#">45</a>
<a href="#">CM 47</a>	<a href="#">HZM19-10-08</a>	<a href="#">620</a>	<a href="#">HZM19-11-08</a>	<a href="#">85</a>
<a href="#">CM 48</a>	<a href="#">HZM19-07-17</a>	<a href="#">100</a>	<a href="#">HZM19-11-08</a>	<a href="#">1250</a>

Table A4.3: Core section depths of the composite profile HZM19 with resulting composite end depths for each core.

Core section	From [mm]	To [mm]	Length [mm]	Composite core section end depth [mm]
HZM19_07_01	138	800	662	662
HZM19_08_01	305	755	451	1113
HZM19_07_02	243	924	681	1794
HZM19_08_02	380	839	459	2254
HZM19_07_03	229	912	683	2936
HZM19_08_03	375	714	339	3275
HZM19_07_04	243	800	557	3833
HZM19_08_04	235	994	759	4592
HZM19_10_01	90	913	823	5415
HZM19_08_05	630	930	299	5715
HZM19_10_02	183	877	693	6409
HZM19_08_06	596	957	361	6770
HZM19_10_03	87	827	740	7510
HZM19_08_07	562	971	409	7919
HZM19_10_04	179	870	691	8611
HZM19_08_08	641	967	326	8937
HZM19_10_05	137	859	722	9659
HZM19_11_06	395	655	260	9919
HZM19_08_10	35	974	939	10859
Technical gap			129	10988
HZM19_10_07	30	810	780	11768
HZM19_11_07	710	1012	302	12071
HZM19_10_08	72	902	830	12902
HZM19_11_08	326	1245	919	13822
HZM19_07_17	100	920	820	14643

899 Table A52: Marker layers (in italics) and radiocarbon dates (Hajdas et al., 2000, 1995 plus one unpublished radiocarbon  
 900 date) vs. composite depth of HZM19. The calibrated median  $^{14}\text{C}$  age is calculated using OxCal with the IntCal20 calibration  
 901 curve. Inconsistent calibrated ages are shown in brackets.

Marker layer and $^{14}\text{C}$ sample ID	HZM19 depth (m)	VT-99 Age (cal BP)	VT-99 cumulative $\pm 1\sigma$ error (yrs)	$^{14}\text{C}$ age (BP)	$^{14}\text{C} \pm 1\sigma$ error (yrs)	Calibrated $^{14}\text{C}$ median age (cal BP)	$^{14}\text{C} \pm 1\sigma$ error (yrs)
<i>ML-36/1</i>	1.16	141	6				
<i>ML-36</i>	1.45	209	8				
<i>ML-35/1</i>	1.81	334	13				
<i>ML-35</i>	2.12	442	18				
<i>ML-34</i>	2.44	572	23				
<i>ML-33/2</i>	2.62	657	26				
<i>ML-33/1</i>	2.69	685	27				
HZM-1.1	2.90	796	32	685	40	644	41
HZM-1.2	2.91	802	32	795	40	708	29
HZM-1.3	2.93	810	32	975	90	869	94
<i>ML-33</i>	2.94	819	33				
<i>ML-32</i>	3.29	985	39				
<i>ML-31/1</i>	3.54	1248	50				
HZM-2.2+3	4.01	1569	63	1565	55	1451	57
<i>ML-31</i>	4.17	1710	68				
<i>ML-30</i>	4.29	1789	72				
<i>ML-29</i>	4.59	1984	79				
<i>ML-28</i>	4.91	2219	89				
HZM-3.1	5.16	2433	97	2405	60	2469	112
<i>ML-27</i>	5.17	2449	98				
HZM-3.3*	5.19	2450	98	2750	60	(2850)	66
<i>ML-26</i>	5.43	2593	104				
HZM-23*	5.45	2595	104	2720	60	(2826)	58
HZM-24	5.61	2754	110	2620	65	2743	101
<i>ML-25/1</i>	5.77	3147	121				
HZM-25	5.97	3704	136	3465	70	3730	96
<i>ML-25</i>	6.11	3992	143				
<i>ML-24</i>	6.21	4420	154				
HZM-26*	6.23	4616	159	4100	90	4624	127
<i>ML-23</i>	6.51	5083	171				
<i>ML-22</i>	6.68	5286	177				
HZM-4.1	6.69	5334	177	4575	65	5243	131
HZM-4.2	6.70	5359	177	4730	70	5462	85
HZM-4.3	6.71	5385	178	4675	70	5409	95
<i>ML-21</i>	6.78	5520	179				
<i>ML-20</i>	6.84	5619	179				
<i>ML-19</i>	7.05	5977	182				
<i>ML-18/2</i>	7.51	6328	184				
<i>ML-18/1</i>	7.70	6590	186				
<i>ML-18</i>	8.06	7274	191				
HZM-5.3	8.13	7428	192	6455	70	7363	68
<i>ML-17/3</i>	8.42	7870	195				
<i>ML-17/2</i>	8.72	8338	198				

ML-17/1	9.03	8943	203				
HZM-6.1	9.33	9649	207	8800	95	9851	170
ML-17	9.40	9746	208				
ML-16	9.66	10169	211				
HZM-7	9.92	10464**	213	9465	45	10705	130
ML-15	9.92	10554	214				
ML-14	10.03	10681	215				
HZM-8	10.07	10708	215	9495	55	10773	148
ML-13	10.24	10999	217				
HZM-9 (UMT)	10.25	11008	217	9560	49	10923	121
HZM-40	10.27	11048	217	9550	80	10901	148
HZM-41	10.33	11109	218	9665	100	10998	154
HZM-42	10.38	11145	218	9565	100	10912	160
HZM-43	10.46	11226	219	9830	100	11264	178
ML-12	10.48	11232	219				
HZM-44	10.52	11267	219	9805	190	11243	329
HZM-45	10.59	11322	219	9905	80	11357	138
HZM-46	10.64	11357**	219	10060	80	11584	159
HZM-10.1	10.67	11339**	219	10085	80	11630	165
HZM-47	10.70	11400	220	10110	110	11680	231
ML-11	10.73	11453	220				
HZM-48*	10.78	11534	221	11040	140	(12959)	120
HZM-50	10.99	11942	241	10080	110	11628	214
ML-9	11.02	11943	241				
HZM-12	11.10	12354	266	10520	90	12509	181
HZM-51	11.14	12530	276	10350	90	12203	194
ML-8	11.20	12578	279				
HZM-13*	11.38	12769	290	11295	85	(13197)	74
ML-7	11.41	12778	291				
HZM-14*	11.48	12861	296	11780	100	(13647)	112
ML-6	11.56	12880	297				
ML-5	11.70	12880	297				
HZM-30	11.74	12925	299	11250	110	13158	109
ML-4	11.86	13087	309				
HZM-16*	12.03	13130	311	13140	140	(15766)	212
HZM-32	12.19	13445	330	11770	135	13642	150
HZM-17	12.26	13472	332	12100	110	13984	183
ML-3	12.40	13646**	339				
HZM-35	12.78	13985	362	12570	130	14858	286
ML-2	12.86	14152**	369				
HZM-18	12.90	14156	372	12430	110	14586	249
ML-1	12.90	14156**	372				
HZM-100***	12.92	14157	372	12380	85	14492	228
HZM-19	12.93	14158	372	12555	80	14879	221

\* Dates described to contain reworked organic material or being fractionated during graphitization (see Hajdas et al., 1995).

\*\* VT-99 dates excluded from modelling due to inconsistencies in documentation.

\*\*\* unpublished radiocarbon age (KIA-1460)

908 Table A6: Age estimations for VT-99 and Models A-D with their 95% confidence intervals in brackets for Ulmener Maar  
909 Tephra (UMT), Younger Dryas/Preboreal-transition (YD/PB), YD duration, Allerød/Younger Dryas-transition (AL/YD),  
910 predicted YD hiatus with duration and position, Laacher See Tephra (LST), Maximum model age at 12.93 m with its mean  
911 and maximum age ranges and position of the maximum age range and maximum difference between VT-99 and each of  
912 the model ranges.

Chronology	VT-99	A	B	C	D	Formatiert: Abstand Nach: 0 Pt., Zeilenabstand: einfach
Age of UMT	10999 [10782, 11216]	10961 [10784, 11090]	10965 [10787, 11093]	10952 [10788, 11067]	10981 [10829, 11088]	Formatiert: Abstand Nach: 0 Pt., Zeilenabstand: einfach
YD/PB transition	11632	11674 [11461, 11965]	11723 [11486, 12070]	11682 [11494, 11913]	11655 [11499, 11845]	Formatiert: Abstand Nach: 0 Pt., Zeilenabstand: einfach
YD duration	974	1038	1014	1012	1073	Formatiert: Abstand Nach: 0 Pt., Zeilenabstand: einfach
AL/YD transition	12606	12712 [12517, 12880]	12737 [12562, 12880]	12694 [12475, 12869]	12728 [12595, 12838]	Formatiert: Abstand Nach: 0 Pt., Zeilenabstand: einfach
Duration of YD hiatus	320	623	603	583	686	Formatiert: Abstand Nach: 0 Pt., Zeilenabstand: einfach
End of YD hiatus	12025	11863 [11571, 12269]	11952 [11623, 12502]	11901 [11646, 12207]	11854 [11651, 12098]	Formatiert: Abstand Nach: 0 Pt., Zeilenabstand: einfach
Age of LST	12880 [12583, 13177]	13010 [12984, 13042]	13010 [12985, 13043]	13009 [12984, 13037]	13011 [12984, 13043]	Formatiert: Abstand Nach: 0 Pt., Zeilenabstand: einfach
Maximum model age (at 12.93 m)	14158 [13786, 14530]	14615 [14339, 14926]	14456 [14236, 14749]	14614 [14332, 14919]	14485 [14287, 14721]	Formatiert: Abstand Nach: 0 Pt., Zeilenabstand: einfach
Mean age range	282	468	456	329	229	Formatiert: Abstand Nach: 0 Pt., Zeilenabstand: einfach
Maximum age range	744	1056	1064	749	447	Formatiert: Abstand Nach: 0 Pt., Zeilenabstand: einfach
Max. age range position (m)	12.93	8.86	8.78	9.18	11.09	Formatiert: Abstand Nach: 0 Pt., Zeilenabstand: einfach
Maximum difference to VT-99 age range	0	655	665	401	0	Formatiert: Abstand Nach: 0 Pt., Zeilenabstand: einfach

913

914

915 *Table A75: Additional dates for the HZM19 chronology with composite depths, ages (cal. BP) and errors used for Bacon*  
 916 *calculations. LST age with error is from Reinig et al. (2020).*

Event	HZM19 comp. depth (cm)	Age (cal. BP)	error
Sediment surface	0.00	-69	1
Chernobyl accident	47.20*	-36	1
Maximum atomic bomb tests	67.20*	-13	1
First atomic bomb tests	79.20*	0	1
Laacher See Tephra	1160.00	13,006	9

917 *\* 22 cm subtracted due to slump event documented by Pb-210 data.*

918

## Data Availability

The results of the different age-depth models carried out for the lacustrine sediment record from Holzmaar are accessible via the PANGAEA data archiving and publication system at <https://doi.org/10.1594/PANGAEA.949393>~~https://doi.org/10.1594/PANGAEA.xxxxxx~~.

## Author contributions

SB and BZ conducted the fieldwork and conceptualized the study. SB described and sampled the sediment, modified and run the Bayesian age-depth models, visualized the data and drafted the first version of the manuscript. WT measured and interpreted lead and cesium data. All authors contributed to the writing and to revising of the manuscript.

## Competing interests

The contact author declares that neither she nor her co-authors have any competing interests.

## Disclaimer

## Acknowledgments

We like to thank Christian Ohlendorf, Rafael Stiens and An-Sheng Lee for participating in the coring campaign of 2019 and also for subsequent help with core opening, sediment preparations and scanning in the GEOPOLAR lab. Furthermore, we want to thank Maarten Blaauw, Arne Ramisch and Alicja Bonk for helpful discussions.

## References

- Anderson, R. Y. and Dean, W. E.: Lacustrine varve formation through time, *Palaeogeogr. Palaeoclimatol.* 62, 215–235, [https://doi.org/10.1016/0031-0182\(88\)90055-7](https://doi.org/10.1016/0031-0182(88)90055-7), 1988.
- Baier, J., Lücke, A., Negendank, J. F. W., Schleser, G.-H., and Zolitschka, B.: Diatom and geochemical evidence of mid- to late Holocene climatic changes at Lake Holzmaar, West-Eifel (Germany), *Quatern. Int.*, 113, 81–96, [https://doi.org/10.1016/S1040-6182\(03\)00081-8](https://doi.org/10.1016/S1040-6182(03)00081-8), 2004.
- Battarbee, R. W., Howells, G. D., Skeffington, R. A., Bradshaw, A. D., Battarbee, R. W., Mason, B. J., Renberg, I., and Talling, J. F.: The causes of lake acidification, with special reference to the role of acid deposition, *Philos. T. Roy. Soc. B.*, 327, 339–347, <https://doi.org/10.1098/rstb.1990.0071>, 1990.
- Berglund, B. E. (Ed.): *Handbook of Holocene Palaeoecology and Palaeohydrology*, John Wiley & Sons Ltd., Chichester, 869 pp., 1986.
- Blaauw, M.: Methods and code for ‘classical’ age-modelling of radiocarbon sequences, *Quat. Geochronol.*, 5, 512–518, <https://doi.org/10.1016/j.quageo.2010.01.002>, 2010.

Formatiert: Block

950 Blaauw, M. and Christen, J. A.: Flexible paleoclimate age-depth models using an autoregressive  
 951 gamma process, *Bayesian Anal.*, 6, 457–474, <https://doi.org/10.1214/11-BA618>, 2011.

952 Blaauw, M., Christen, J. A., Bennett, K. D., and Reimer, P. J.: Double the dates and go for Bayes —  
 953 Impacts of model choice, dating density and quality on chronologies, *Quaternary Sci. Rev.*, 188,  
 954 58–66, <https://doi.org/10.1016/j.quascirev.2018.03.032>, 2018.

955 Blaauw, M., Christen, J. A., Lopez, M. A. A., Vazquez, J. E., V. O. M. G., Belding, T., Theiler, J., Gough,  
 956 B., and Karney, C.: rbacon: Age-Depth Modelling using Bayesian Statistics, CRAN [code], 2021.

957 Bonk, A., Müller, D., Ramisch, A., Kramkowski, M. A., Noryśkiewicz, A. M., Sekudewicz, I.,  
 958 Gąsiorowski, M., Lubierda-Durnaś, K., Słowiński, M., Schwab, M., Tjallingii, R., Brauer, A., and  
 959 Błaskiewicz, M.: Varve microfacies and chronology from a new sediment record of Lake Gościąg  
 960 (Poland), *Quaternary Sci. Rev.*, 251, 106715, <https://doi.org/10.1016/j.quascirev.2020.106715>,  
 961 2021.

962 Brauer, A.: Weichselzeitliche Seesedimente des Holzmaars - Warvenchronologie des Hochglazials  
 963 und Nachweis von Klimaschwankungen, Ph.D. thesis, Universität Trier, Documenta naturae, 85,  
 964 1–210, 1994.

965 Brauer, A., Hajdas, I., Negendank, J. F. W., Rein, B., Vos, H., and Zolitschka, B.: Warvenchronologie -  
 966 eine Methode zur absoluten Datierung und Rekonstruktion kurzer und mittlerer solarer  
 967 Periodizitäten, *Geowissenschaften*, 12, 325–332, 1994.

968 Brauer, A., Endres, C., Günter, C., Litt, T., Stebich, M., and Negendank, J. F. W.: High resolution  
 969 sediment and vegetation responses to Younger Dryas climate change in varved lake sediments  
 970 from Meerfelder Maar, Germany, *Quaternary Sci. Rev.*, 18, 321–329,  
 971 [https://doi.org/10.1016/S0277-3791\(98\)00084-5](https://doi.org/10.1016/S0277-3791(98)00084-5), 1999.

972 Bronk Ramsey, C.: Deposition models for chronological records, *Quaternary Sci. Rev.*, 27, 42–60,  
 973 <https://doi.org/10.1016/j.quascirev.2007.01.019>, 2008.

974 Brooks, S. P. and Gelman, A.: General Methods for Monitoring Convergence of Iterative  
 975 Simulations, *J. Comput. Graph. Stat.*, 7, 434–455,  
 976 <https://doi.org/10.1080/10618600.1998.10474787>, 1998.

977 Büchel, G.: Maars of the Westeifel, Germany, in: *Paleolimnology of European Maar Lakes*, edited  
 978 by: Negendank, J. F. W. and Zolitschka, B., Springer, Berlin, Heidelberg, 1–13,  
 979 <https://doi.org/10.1007/BFb0117585>, 1993.

980 Cohen, A. S.: *Paleolimnology. The history and evolution of lake systems*, Oxford University Press,  
 981 Oxford, 500 pp., 2003.

982 Davies, S. M.: Cryptotephra: the revolution in correlation and precision dating, *Journal of*  
 983 *Quaternary Science*, 30, 114–130, <https://doi.org/10.1002/jqs.2766>, 2015.

984 De Geer, G.: A geochronology of the last 12,000 years., *Eleventh International Geological Congress*,  
 985 Stockholm, 1, 241–253, 1912.

986 Dean, W. E., Bradbury, J. P., Anderson, R. Y., and Barnosky, C. W.: The Variability of Holocene  
 987 Climate Change: Evidence from Varved Lake Sediments, *Science*, 226, 1191–1194,  
 988 <https://doi.org/10.1126/science.226.4679.1191>, 1984.

989 Fortin, D., Praet, N., McKay, N. P., Kaufman, D. S., Jensen, B. J. L., Haeussler, P. J., Buchanan, C., and  
 990 De Batist, M.: New approach to assessing age uncertainties – The 2300-year varve chronology

Formatiert: Deutsch (Deutschland)

Formatiert: Deutsch (Deutschland)



991 from Eklutna Lake, Alaska (USA), *Quaternary Sci. Rev.*, 203, 90–101,  
992 <https://doi.org/10.1016/j.quascirev.2018.10.018>, 2019.

1p93 García, M. L., Birlo, S., and Zolitschka, B.: Paleoenvironmental changes of the last 16,000 years  
994 based on diatom and geochemical stratigraphies from the varved sediment of Holzmaar (West-  
995 Eifel Volcanic Field, Germany), *Quaternary Science Reviews*, 293, 107691,  
996 <https://doi.org/10.1016/j.quascirev.2022.107691>, 2022.

1p97 Goslar, T., Kuc, T., Ralska-Jasiewiczowa, M., Różanski, K., Arnold, M., Bard, E., van Geel, B., Pazdur,  
998 M., Szeroczyńska, K., Wicik, B., Więckowski, K., and Walanus, A.: High-resolution lacustrine record  
999 of the late glacial/holocene transition in central Europe, *Quaternary Sci. Rev.*, 12, 287–294,  
1000 [https://doi.org/10.1016/0277-3791\(93\)90037-M](https://doi.org/10.1016/0277-3791(93)90037-M), 1993.

1p01 Hajdas, I., Zolitschka, B., Ivy-Ochs, S. D., Beer, J., Bonani, G., Leroy, S. A. G., Negendank, J. W.,  
1002 Ramrath, M., and Suter, M.: AMS radiocarbon dating of annually laminated sediments from lake  
1003 Holzmaar, Germany, *Quaternary Sci. Rev.*, 14, 137–143, [https://doi.org/10.1016/0277-3791\(94\)00123-S](https://doi.org/10.1016/0277-3791(94)00123-S), 1995.

1p05 Hajdas, I., Bonani, G., and Zolitschka, B.: Radiocarbon Dating of Varve Chronologies: Soppensee  
1006 and Holzmaar Lakes after Ten Years, *Radiocarbon*, 42, 349–353,  
1007 <https://doi.org/10.1017/S0033822200030290>, 2000.

1p08 Hajdas-Skowronek, I.: Extension of the radiocarbon calibration curve by AMS dating of laminated  
1009 sediments of Lake Soppensee and Lake Holzmaar, Ph. D thesis, ETH Zurich,  
1010 <https://doi.org/10.3929/ETHZ-A-000916163>, 1993.

1p11 Haslett, J. and Parnell, A.: A simple monotone process with application to radiocarbon-dated depth  
1012 chronologies, *J. R. Stat. Soc. C-Appl.*, 57, 399–418, <https://doi.org/10.1111/j.1467-9876.2008.00623.x>, 2008.

1p14 Jenny, J.-P., Koirala, S., Gregory-Eaves, I., Francus, P., Niemann, C., Ahrens, B., Brovkin, V., Baud, A.,  
1015 Ojala, A. E. K., Normandeau, A., Zolitschka, B., and Carvalhais, N.: Human and climate global-scale  
1016 imprint on sediment transfer during the Holocene, *P. Natl. Acad. Sci.*, 116, 22972–22976,  
1017 <https://doi.org/10.1073/pnas.1908179116>, 2019.

1p18 Kelts, K., Briegel, U., Ghilardi, K., and Hsu, K.: The limnogeology-ETH coring system, *Schweiz. Z.*  
1019 *Hydro.*, 48, 104–115, <https://doi.org/10.1007/BF02544119>, 1986.

1p20 Kienel, U., Schwab, M. J., and Schettler, G.: Distinguishing climatic from direct anthropogenic  
1021 influences during the past 400 years in varved sediments from Lake Holzmaar (Eifel, Germany), *J.*  
1022 *Paleolimnol.*, 33, 327–347, <https://doi.org/10.1007/s10933-004-6311-z>, 2005.

1p23 Lacourse, T. and Gajewski, K.: Current practices in building and reporting age-depth models,  
1024 *Quaternary Res.*, 96, 28–38, <https://doi.org/10.1017/qua.2020.47>, 2020.

1p25 Lamoureux, S.: Varve Chronology Techniques, in: *Tracking Environmental Change Using Lake*  
1026 *Sediments: Basin Analysis, Coring, and Chronological Techniques*, edited by: Last, W. M. and Smol,  
1027 J. P., Springer Netherlands, Dordrecht, 247–260, [https://doi.org/10.1007/0-306-47669-X\\_11](https://doi.org/10.1007/0-306-47669-X_11),  
1028 2001.

1p29 Last, W. M. and Smol, J. P. (Eds.): *Tracking Environmental Change Using Lake Sediments*, vol. 1,  
1030 *Basin Analysis, Coring, and Chronological Techniques*, Springer Netherlands, Dordrecht, 548 pp.,  
1031 2001a.

- 1p32 Last, W. M. and Smol, J. P. (Eds.): Tracking Environmental Change Using Lake Sediments, vol. 2,  
1033 Physical and Geochemical Methods, Springer Netherlands, Dordrecht, 504 pp., 2001b.
- 1p34 Leroy, S. A. G., Zolitschka, B., Negendank, J. F. W., and Seret, G.: Palynological analyses in the  
1035 laminated sediment of Lake Holzmaar (Eifel, Germany): duration of Lateglacial and Preboreal  
1036 biozones, *Boreas*, 29, 52–71, <https://doi.org/10.1111/j.1502-3885.2000.tb01200.x>, 2000.
- 1p37 Litt, T., Schölzel, C., Kühl, N., and Brauer, A.: Vegetation and climate history in the Westeifel  
1038 Volcanic Field (Germany) during the past 11 000 years based on annually laminated lacustrine  
1039 maar sediments, *Boreas*, 38, 679–690, <https://doi.org/10.1111/j.1502-3885.2009.00096.x>, 2009.
- 1p40 Lorenz, V.: Explosive Volcanism of the West Eifel Volcanic Field/Germany, in: Developments in  
1041 Petrology, vol. 11, edited by: Kornprobst, J., Elsevier, 299–307, <https://doi.org/10.1016/B978-0-444-42273-6.50026-2>, 1984.
- 1p43 Lorenz, V., Lange, T., and Büchel, G.: Die Vulkane der Westeifel, *Jahresberichte und Mitteilungen*  
1044 *des Oberrheinischen Geologischen Vereins*, 102, 379–411,  
1045 <https://doi.org/10.1127/jmogv/102/0022>, 2020.
- 1p46 Lotter, A. F.: Absolute Dating of the Late-Glacial Period in Switzerland Using Annually Laminated  
1047 Sediments, *Quaternary Res.*, 35, 321–330, [https://doi.org/10.1016/0033-5894\(91\)90048-A](https://doi.org/10.1016/0033-5894(91)90048-A),  
1048 1991.
- 1p49 Lücke, A., Schleser, G. H., Zolitschka, B., and Negendank, J. F. W.: A Lateglacial and Holocene organic  
1050 carbon isotope record of lacustrine palaeoproductivity and climatic change derived from varved  
1051 lake sediments of Lake Holzmaar, Germany, *Quaternary Sci. Rev.*, 22, 569–580,  
1052 [https://doi.org/10.1016/S0277-3791\(02\)00187-7](https://doi.org/10.1016/S0277-3791(02)00187-7), 2003.
- 1p53 Martin-Puertas, C., Walsh, A. A., Blockley, S. P. E., Harding, P., Biddulph, G. E., Palmer, A., Ramisch,  
1054 A., and Brauer, A.: The first Holocene varve chronology for the UK: Based on the integration of  
1055 varve counting, radiocarbon dating and tephrostratigraphy from Diss Mere (UK), *Quat.*  
1056 *Geochronol.*, 61, 101134, <https://doi.org/10.1016/j.quageo.2020.101134>, 2021.
- 1p57 Meyer, W.: Zur Entstehung der Maare in der Eifel, *Z. Dtsch. Ges. Geowiss.*, 141–155,  
1058 <https://doi.org/10.1127/zdgg/136/1985/141>, 1985.
- 1p59 Meyer, W.: *Geologie der Eifel*, 4th ed., Schweizerbart Science Publishers, Stuttgart, Germany, 2013.
- 1p60 Meyer, W. and Stets, J.: Pleistocene to Recent tectonics in the Rhenish Massif (Germany), *Neth. J.*  
1061 *Geosci.*, 81, 217–221, <https://doi.org/10.1017/S0016774600022460>, 2002.
- 1p62 Muggeo, V. M. R.: segmented: Regression Models with Break-Points / Change-Points Estimation,  
1063 CRAN [code], 2022.
- 1p64 Nesje, A., Søgne, K., Elgersma, A., and Dahl, S. O.: A Piston Corer for Lake Sediments, *Norsk. Geogr.*  
1065 *Tidsskr.*, 41, 123–125, <https://doi.org/10.1080/00291958708621986>, 1987.
- 1p66 Olsson, I. U.: Radiometric dating, John Wiley and Sons, United Kingdom, 1986.
- 1p67 O'Sullivan, P. E.: Annually-laminated lake sediments and the study of Quaternary environmental  
1068 changes — a review, *Quaternary Sci. Rev.*, 1, 245–313, [https://doi.org/10.1016/0277-3791\(83\)90008-2](https://doi.org/10.1016/0277-3791(83)90008-2), 1983.
- 1p70 Pearson, G. W., Pilcher, J. R., Baillie, M. G. L., and Hillam, J.: Absolute radiocarbon dating using a low  
1071 altitude European tree-ring calibration, *Nature*, 270, 25–28, <https://doi.org/10.1038/270025a0>,  
1072 1977.

Formatiert: Deutsch (Deutschland)

Formatiert: Deutsch (Deutschland)

- 1073 Prasad, S. and Baier, J.: Tracking the impact of mid- to late Holocene climate change and  
1074 anthropogenic activities on Lake Holzmaar using an updated Holocene chronology, *Global Planet.*  
1075 *Change*, 122, 251–264, <https://doi.org/10.1016/j.gloplacha.2014.08.020>, 2014.
- 1076 R Core Team: R: A Language and Environment for Statistical Computing, 2021.
- 1077 Reimer, P. J., Austin, W. E. N., Bard, E., Bayliss, A., Blackwell, P. G., Bronk Ramsey, C., Butzin, M.,  
1078 Cheng, H., Edwards, R. L., Friedrich, M., Grootes, P. M., Guilderson, T. P., Hajdas, I., Heaton, T. J.,  
1079 Hogg, A. G., Hughen, K. A., Kromer, B., Manning, S. W., Muscheler, R., Palmer, J. G., Pearson, C., van  
1080 der Plicht, J., Reimer, R. W., Richards, D. A., Scott, E. M., Southon, J. R., Turney, C. S. M., Wacker, L.,  
1081 Adolphi, F., Büntgen, U., Capano, M., Fahrni, S. M., Fogtmann-Schulz, A., Friedrich, R., Köhler, P.,  
1082 Kudsk, S., Miyake, F., Olsen, J., Reinig, F., Sakamoto, M., Sookdeo, A., and Talamo, S.: The INTCAL20  
1083 Northern Hemisphere Radiocarbon Age Calibration Curve (0–55 cal kBP), *Radiocarbon*, 62, 725–  
1084 757, <https://doi.org/10.1017/RDC.2020.41>, 2020.
- 1085 Reinig, F., Wacker, L., Jöris, O., Oppenheimer, C., Guidobaldi, G., Nievergelt, D., Adolphi, F.,  
1086 Cherubini, P., Engels, S., Esper, J., Land, A., Lane, C., Pfan, H., Remmele, S., Sigl, M., Sookdeo, A., and  
1087 Büntgen, U.: Precise date for the Laacher See eruption synchronizes the Younger Dryas, *Nature*,  
1088 595, 66–69, <https://doi.org/10.1038/s41586-021-03608-x>, 2021.
- 1089 Renberg, I. and Hansson, H.: A pump freeze corer for recent sediments, *Limnol. Oceanogr.*, 38,  
1090 1317–1321, <https://doi.org/10.4319/lo.1993.38.6.1317>, 1993.
- 1091 Renberg, I., Persson, M. W., and Emteryd, O.: Pre-industrial atmospheric lead contamination  
1092 detected in Swedish lake sediments, *Nature*, 368, 323–326, <https://doi.org/10.1038/368323a0>,  
1093 1994.
- 1094 Saarnisto, M.: Long varve series in Finland, *Boreas*, 14, 133–137, <https://doi.org/10.1111/j.1502-3885.1985.tb00905.x>, 1985.
- 1096 Saarnisto, M.: Annually laminated lake sediments, in: *Handbook of Holocene palaeoecology and*  
1097 *palaeohydrology*, edited by: Berglund, B. E., John Wiley and Sons Ltd, Chichester, 343–370, 1986.
- 1098 Scharf, B.: *Limnologische Beschreibung, Nutzung und Unterhaltung von Eifelmaaren*, Ministerium  
1099 für Umwelt und Gesundheit Rheinland-Pfalz, Mainz, 1987.
- 1100 Scharf, B. W. and Oehms, M.: Physical and chemical characteristics, in: *Limnology of Eifel maar*  
1101 *lakes. Ergebnisse der Limnologie*, vol. 38, edited by: Scharf, B. W. and Björk, S., E. Schweizerbart,  
1102 Stuttgart, 63–83, 1992.
- 1103 Schmincke, H.-U.: The Quaternary Volcanic Fields of the East and West Eifel (Germany), in: *Mantle*  
1104 *Plumes: A Multidisciplinary Approach*, edited by: Ritter, J. R. R. and Christensen, U. R., Springer,  
1105 Berlin, Heidelberg, 241–322, [https://doi.org/10.1007/978-3-540-68046-8\\_8](https://doi.org/10.1007/978-3-540-68046-8_8), 2007.
- 1106 Schmincke, H.-U.: *Vulkane der Eifel*, 2nd ed., Springer, Berlin, Heidelberg, 2014.
- 1107 Schnurrenberger, D., Russell, J., and Kelts, K.: Classification of lacustrine sediments based on  
1108 sedimentary components, *J. Paleolimnol.*, 29, 141–154,  
1109 <https://doi.org/10.1023/A:1023270324800>, 2003.
- 1110 Shanahan, T. M., Beck, J. W., Overpeck, J. T., McKay, N. P., Pigati, J. S., Peck, J. A., Scholz, C. A., Heil, C.  
1111 W., and King, J.: Late Quaternary sedimentological and climate changes at Lake Bosumtwi Ghana:  
1112 New constraints from laminae analysis and radiocarbon age modeling, *Palaeogeogr. Palaeoclimatol.*,  
1113 361–362, 49–60, <https://doi.org/10.1016/j.palaeo.2012.08.001>, 2012.

Formatiert: Deutsch (Deutschland)

Formatiert: Deutsch (Deutschland)

Formatiert: Deutsch (Deutschland)

- 1114 Sirocko, F., Dietrich, S., Veres, D., Grootes, P. M., Schaber-Mohr, K., Seelos, K., Nadeau, M.-J., Kromer,  
1115 B., Rothacker, L., Röhrner, M., Krbetschek, M., Appleby, P., Hambach, U., Rolf, C., Sudo, M., and Grim,  
1116 S.: Multi-proxy dating of Holocene maar lakes and Pleistocene dry maar sediments in the Eifel,  
1117 Germany, *Quaternary Sci. Rev.*, 62, 56–76, <https://doi.org/10.1016/j.quascirev.2012.09.011>,  
1118 2013.
- 1119 Stuiver, M., Reimer, P. J., Bard, E., Beck, J. W., Burr, G. S., Hughen, K. A., Kromer, B., McCormac, G.,  
1120 Van Der Plicht, J., and Spurk, M.: INTCAL98 Radiocarbon Age Calibration, 24,000–0 cal BP,  
1121 *Radiocarbon*, 40, 1041–1083, <https://doi.org/10.1017/S0033822200019123>, 1998.
- 1122 Telford, R., Heegaard, E., and Birks, H.: All age–depth models are wrong: but how badly?,  
1123 *Quaternary Sci. Rev.*, 23, 1–5, <https://doi.org/10.1016/j.quascirev.2003.11.003>, 2004.
- 1124 Trachsel, M. and Telford, R. J.: All age–depth models are wrong, but are getting better, Holocene,  
1125 27, 860–869, <https://doi.org/10.1177/0959683616675939>, 2017.
- 1126 Turkey, C. S. M. and Lowe, J. J.: Tephrochronology, in: *Tracking Environmental Change Using Lake*  
1127 *Sediments: Basin Analysis, Coring, and Chronological Techniques*, edited by: Last, W. M. and Smol,  
1128 J. P., Springer Netherlands, Dordrecht, 451–471, [https://doi.org/10.1007/0-306-47669-X\\_16](https://doi.org/10.1007/0-306-47669-X_16),  
1129 2001.
- 1130 Vandergoes, M. J., Howarth, J. D., Dunbar, G. B., Turnbull, J. C., Roop, H. A., Levy, R. H., Li, X., Prior,  
1131 C., Norris, M., Keller, L. D., Baisden, W. T., Ditchburn, R., Fitzsimons, S. J., and Bronk Ramsey, C.:  
1132 Integrating chronological uncertainties for annually laminated lake sediments using layer  
1133 counting, independent chronologies and Bayesian age modelling (Lake Ohau, South Island, New  
1134 Zealand), *Quaternary Sci. Rev.*, 188, 104–120, <https://doi.org/10.1016/j.quascirev.2018.03.015>,  
1135 2018.
- 1136 Wright, H. E., Mann, D. H., and Glaser, P. H.: Piston Corers for Peat and Lake Sediments, *Ecology*,  
1137 65, 657–659, <https://doi.org/10.2307/1941430>, 1984.
- 1138 Zolitschka, B.: Spätquartäre Sedimentationsgeschichte des Meerfelder Maars (Westeifel).  
1139 Mikrostratigraphie jahreszeitlich geschichteter Seesedimente, *E&G Quaternary Sci. J.*, 38, 87–93,  
1140 <https://doi.org/10.3285/eg.38.1.08>, 1988.
- 1141 Zolitschka, B.: Jahreszeitlich geschichtete Seesedimente aus dem Holzmaar und dem Meerfelder  
1142 Maar, *Z. Dtsch. Ges. Geowiss.*, 25–33, <https://doi.org/10.1127/zdgg/140/1989/25>, 1989.
- 1143 Zolitschka, B.: Jahreszeitlich geschichtete Seesedimente ausgewählter Eifelmaare -  
1144 paläolimnologische Untersuchung als Beitrag zur spät- und postglazialen Klima- und  
1145 Besiedlungsgeschichte, Ph.D. thesis, Universität Trier, *Documenta naturae*, 60, 226, 1990.
- 1146 Zolitschka, B.: Absolute dating of late Quaternary Lacustrine sediments by high resolution varve  
1147 chronology, 214, 59–61, <https://doi.org/10.1007/BF00050932>, 1991.
- 1148 Zolitschka, B.: Climatic change evidence and lacustrine varves from maar lakes, Germany, *Clim.*  
1149 *Dynam.*, 6, 229–232, <https://doi.org/10.1007/BF00193535>, 1992.
- 1150 Zolitschka, B.: A 14,000 year sediment yield record from western Germany based on annually  
1151 laminated lake sediments, *Geomorphology*, 22, 1–17, [https://doi.org/10.1016/S0169-555X\(97\)00051-2](https://doi.org/10.1016/S0169-555X(97)00051-2), 1998a.
- 1153 Zolitschka, B.: Paläoklimatische Bedeutung laminierter Sedimente: Holzmaar (Eifel, Deutschland),  
1154 Lake C2 (Northwest-Territorien, Kanada) und Lago Grande di Monticchio (Basilicata, Italien),  
1155 *Habil. thesis, Universität Potsdam, Relief Boden Paläoklima*, 13, 176, 1998b.

Formatiert: Deutsch (Deutschland)

Formatiert: Deutsch (Deutschland)

- 1156 Zolitschka, B., Haverkamp, B., and Negendank, J. F. W.: Younger Dryas Oscillation — Varve Dated  
 1157 Microstratigraphic, Palynological and Palaeomagnetic Records from Lake Holzmaar, Germany, in:  
 1158 The Last Deglaciation: Absolute and Radiocarbon Chronologies, edited by: Bard, E. and Broecker,  
 1159 W. S., Springer Berlin Heidelberg, Berlin, Heidelberg, 81–101, [https://doi.org/10.1007/978-3-](https://doi.org/10.1007/978-3-642-76059-4_6)  
 1160 [642-76059-4\\_6](https://doi.org/10.1007/978-3-642-76059-4_6), 1992.
- 1161 Zolitschka, B., Brauer, A., Negendank, J. F. W., Stockhausen, H., and Lang, A.: Annually dated late  
 1162 Weichselian continental paleoclimate record from the Eifel, Germany, *Geology*, 28, 783–786,  
 1163 [https://doi.org/10.1130/0091-7613\(2000\)28<783:ADLWCP>2.0.CO;2](https://doi.org/10.1130/0091-7613(2000)28<783:ADLWCP>2.0.CO;2), 2000.
- 1164 Zolitschka, B., Francus, P., Ojala, A. E. K., and Schimmelmann, A.: Varves in lake sediments – a  
 1165 review, *Quaternary Sci. Rev.*, 117, 1–41, <https://doi.org/10.1016/j.quascirev.2015.03.019>, 2015.
- 1166