

# Marine reservoir ages for coastal West Africa

Guillaume Soulet<sup>1</sup>, Philippe Maestrati<sup>2</sup>, Serge Gofas<sup>2,3</sup>, Germain Bayon<sup>1</sup>, Fabien Dewilde<sup>1</sup>, Maylis Labonne<sup>4</sup>, Bernard Dennielou<sup>1</sup>, Franck Ferraton<sup>4</sup>, Giuseppe Siani<sup>5</sup>

1 Ifremer, Univ Brest, CNRS, Geo-Ocean UMR6538, F-29280, Plouzané, France

2 Muséum National d'Histoire Naturelle, Paris, DGD-Collections, France

3 Departamento de Biología Animal, Facultad de Ciencias, Universidad de Málaga, Málaga, Spain

4 MARBEC, Univ Montpellier, CNRS, Ifremer, IRD, Montpellier, France

5 GEOPS, UMR 8148 Université Paris-Saclay, Orsay, France

Correspondence to: Guillaume Soulet (gsoulet@ifremer.fr)

10 **Abstract.** We measured the <sup>14</sup>C age of pre-bomb suspension-feeding bivalves of known-age from coastal West Africa (n=30) across a latitudinal transect extending from 33°N to 15°S. The specimens are from the collections of the Muséum National d'Histoire Naturelle (Paris, France). They were carefully chosen to ensure that the specimens were alive when collected or died not long before collection. From the <sup>14</sup>C-dating of these known-age bivalves, we calculated the marine reservoir age (as ΔR and R values) for each specimen. ΔR values were calculated relative to the Marine20 calibration curve and the R values relative to Intcal20 or SHcal20 calibration curves. Except for five outliers, the ΔR and R values were quite homogenous to a weighted mean value of  $-77-72 \pm 47-42$  <sup>14</sup>C yrs (1sd, n = 2524), and of  $400-406 \pm 59-56$  <sup>14</sup>C yrs (1sd, n = 2524), respectively. These values are typical of low latitude marine reservoir age values. Five suspension-feeding species living in five different ecological habitats were studied. For localities where different species were available, the results yielded similar results whatever the species considered suggesting that in these locations the habitat has only a limited impact on the marine reservoir age reconstruction. We show that our measured marine reservoir ages follow the declining trend of the global marine reservoir age starting ca. 1900 AD, suggesting that marine reservoir age of coastal West Africa is driven, at least at first order, by the global carbon cycle and climate the atmospheric CO<sub>2</sub> <sup>14</sup>C ageing due to fossil fuel burning rather than by local effects. Each outlier was discussed. Local upwelling conditions or Sub-fossil specimens likely may explain the older <sup>14</sup>C age and thus larger marine reservoir age measured for these samples. *Bucardium ringens* might not a best choice for marine reservoir age reconstructions.

## 1 Introduction

The marine reservoir age (R) at a given calendar time/year (t) is the difference between the radiocarbon age (<sup>14</sup>C) of the dissolved inorganic carbon (DIC) of the ocean (<sup>14</sup>C<sub>m</sub>) and that of atmospheric CO<sub>2</sub> (<sup>14</sup>C<sub>atm</sub>) (Stuiver et al., 1986; Stuiver and Braziunas, 1993)(Stuiver and Polach, 1977; Ascough et al., 2005; Soulet et al., 2016; Skinner and Bard, 2022; Soulet, 2015):

$$30 \quad R(t) = {}^{14}\text{C}_m(t) - {}^{14}\text{C}_{\text{atm}}(t) \quad (1)$$

At global scale, the marine reservoir age of the surface mixed layer of the ocean is set by the exchange of “young” CO<sub>2</sub> at the atmosphere-ocean interface, plus the exchange of DIC between oceanic surface waters and deep waters that contain large amounts of “old” DIC (Bard, 1988; Skinner and Bard, 2022). [Box models have been used to study the distribution of radiocarbon in Earth’s system since the 1950s \(Craig, 1957; Revelle and Suess, 1957; Arnold and Anderson, 1957; Siegenthaler, 1983\).](#) The <sup>14</sup>C age of the global ocean over time, i.e., the Marine20 calibration curve (Heaton et al., 2020), has been modelled using the global carbon cycle box model BICYCLE (Köhler et al., 2006; Köhler and Fischer, 2006, 2004; Köhler et al., 2005) and the Northern Hemisphere atmospheric <sup>14</sup>C calibration curve (IntCal20; Reimer et al., 2020). While the global marine calibration curve (Marine20) is widely used to derive calibrated ages from <sup>14</sup>C dating of marine samples, it does not account for local marine <sup>14</sup>C offsets due to, for instance, continental carbon inputs to the coastal ocean, regional winds, and changes in the oceanic circulation and climate (Bard, 1988; Alves et al., 2018; Skinner and Bard, 2022; Heaton et al., 2023). Hence, the usefulness of the ΔR metric (Stuiver et al., 1986; Stuiver and Braziunas, 1993; Reimer and Reimer, 2017) that is the difference between the <sup>14</sup>C age of any marine sample (<sup>14</sup>C<sub>m</sub>) and that of the marine calibration curve (<sup>14</sup>C<sub>Marine20</sub>) at the same time (t):

$$\Delta R(t) = {}^{14}\text{C}_m(t) - {}^{14}\text{C}_{\text{Marine20}}(t) \quad (2)$$

The local marine reservoir age offset (ΔR) is known to vary largely as demonstrated by pre-bomb values ranging between –500 to +2000 <sup>14</sup>C years (Reimer and Reimer, 2001) depending on the location. [Most larger-larger ΔR values are located at high-latitudes while values close to ΔR = 0 <sup>14</sup>C years are located at low latitudes \(Bard, 1988; Bard et al., 1994\).](#)

From a geochronological perspective (i.e., calibration of marine <sup>14</sup>C dates and building age-depth models from marine <sup>14</sup>C dates), knowing ΔR(t) is of crucial interest to correct marine <sup>14</sup>C dates for a local <sup>14</sup>C offset compared to the global marine calibration curve and hence a pre-requirement to derive accurate calendar ages. Reconstructing ΔR(t) values from unstudied areas is also valuable as it could contribute to deriving regional/local marine calibration curves from the global one using 3-D large-scale ocean circulation model (Butzin et al., 2017; Alves et al., 2019).

From a carbon cycle perspective, the R(t) and ΔR(t) are also very important as they reflect <sup>14</sup>C disequilibria between the ocean and the atmosphere and hence they are key proxies to understand local variations of the global carbon cycle, and its evolution over time with changing climate and environment (Skinner et al., 2015, 2010; Lindsay et al., 2016; Soulet et al., 2011; Siani et al., 2001; Schefuß et al., 2016; Heaton et al., 2021).

On a whole, efforts to estimate R(t) and ΔR(t) values wherever possible are valuable to the understanding of both modern and past carbon cycle, and the reconstruction of climate and environmental changes based on sedimentary archives.

Pre-bomb R(t) and ΔR(t) values for coastal West Africa are very sparse. According to the Marine Reservoir Correction Database (Reimer and Reimer, 2001; <http://calib.org/marine/>; last seen 15/11/2022), from Oran on the Mediterranean coast of Algeria (Siani et al., 2000) to Hondelkip Bay on the Atlantic coast of South Africa (Dewar et al., 2012), only a few marine reservoir ages from Mauritania and Senegal were reported (Ndeye, 2008) (Fig. 1). For Mauritania, collection sites were Nouadhibou (formerly Port-Étienne, 2 samples), the area of the Cape Timiris – El Mamghar (3 samples). Two samples were collected from an unknown location from coastal Mauritania. For Senegal, collection sites were restricted to the Dakar area

65 (Almadies, Dakar harbour, Gorée Island and Rufisque; 5 samples). Thirteen additional samples were from unknown locations from coastal Senegal.

In this study we report new marine reservoir age values (n=30) based on the  $^{14}\text{C}$  dating of bivalves with a known pre-bomb collection date and collected across a latitudinal transect extending from Mohammedia (Morocco, 33°N) to Moçâmedes (Angola, 15°S). Our suite of sample includes specimens from Mauritania, Senegal, Republic of Guinea, Sierra Leone, Ivory  
70 Coast, Benin, Gabon and Republic of Congo (Fig. 1, Table S1 in the Supplement). We used specimens of five different species: *Senilia senilis*, *Bucardium ringens*, *Donax rugosus*, *Ostrea stentina* and *Pseudochama gryphina*. We briefly discuss our results in the context of local environmental setting of the studied bivalves and regional oceanography of the Eastern Atlantic Ocean.

## 2 Material and methods

### 2.1 Material

75 Bivalve shells were selected from the collections of the Muséum National d'Histoire Naturelle (MNHN) (Paris, France). We carefully chose pre-bomb specimens of known-age and ensured that they were collected alive or very soon after death. For example, specimens with articulated valves and exhibiting flesh remains inside the shell were clearly collected alive. For *Senilia senilis*, the presence of the fragile periostracum provides evidence that the specimen was collected fresh. For *Bucardium ringens*, remains of the hinge ligament indicate that the bivalve death occurred not long before collection. The collection date  
80 was also carefully checked. Below, we provide background information for the five different bivalve species investigated in this study. Additional information for each sample is given in section 3.1.

*Senilia senilis* (Linnaeus, 1758) can be found from Mauritania to northern Angola. It lives in fine sand in estuaries, creeks or lagoon with regular tidal influence from the lower intertidal zone to about 2 meters water depth. The ~~speeiespecies~~ is tolerant to seasonal salinity changes (von Cosel and Gofas, 2019). *S. senilis* is a suspension feeder that lives in the top 5-10-cm layer  
85 of the sediment (Okera, 1976; Catry et al., 2017).

*Bucardium ringens* (Bruguère, 1789) occurs from Mauritania to southern Angola. It lives in clean fine sand and mixed sand on open coast from shallow (5-10 meters depth) to about 50 meters depth. Shells and valves are commonly cast ashore on beaches but live-taken specimens are rare (von Cosel and Gofas, 2019). *B. ringens* is likely a suspension feeder as typically are cardiiids (Herrera et al., 2015).

90 *Donax rugosus* (Linnaeus, 1758) occurs from Mauritania to Ghana and from northern Angola to southern Angola. It lives in mixed and coarse sand in the surf zone of open beaches (von Cosel and Gofas, 2019). *D. rugosus* is a suspension feeder (Smith, 1971).

*Ostrea stentina* (Payraudeau, 1826) can be found from southern Portugal to Ghana, then from Gabon to northern Angola. It is common and occurs on various types of hard substrate such as rocks, stones, pebbles and other oysters from 1 to 30 meters  
95 depths. It can also be found in lagoons, inlets and creeks under marine condition (von Cosel and Gofas, 2019). *O. stentina* is a suspension feeder (Türkmen et al., 2005).

*Pseudochama gryphina* (Lamarck, 1819) occurs from Southern Portugal to Mauritania and from Gabon to southern Angola (von Cosel and Gofas, 2019) and lives on hard substrate such as rocks and stones in clear water offshore about 10 to 60 meters water depth. *P. gryphina* is a suspension feeder (Sessa et al., 2013).

100 A small piece (30-100 mg) of the outermost layers of each shell was cut using a Dremel™ rotary tool fitted with a cut-off wheel. We focused on the external part of the shell to ensure that we sampled and dated the most recent part (likely the last few months) of the specimen. The shell carbonate samples were then sonicated and rinsed in deionized water at least 5 times. Samples were coarsely crushed and split into a subsample for stable isotopic analysis and a subsample for <sup>14</sup>C analysis.

## 2.2 Radiocarbon measurements

105 Samples were washed with dilute HNO<sub>3</sub> (0.01M) for 15 min then rinsed to neutral pH. Then the shell carbonate was converted into CO<sub>2</sub> following LMC14 laboratory (Laboratoire de Mesure du Radiocarbone, Saclay, France) standard phosphoric acid hydrolysis procedure (Tisnérat-Laborde et al., 2001; Dumoulin et al., 2017). The CO<sub>2</sub> was then converted to graphite (Cottureau et al., 2007; Dumoulin et al., 2017) and analyzed for its <sup>14</sup>C composition by Accelerator Mass Spectrometry (AMS) using the Artémis <sup>14</sup>C AMS facility (Moreau et al., 2013). Results are corrected for the <sup>13</sup>C/<sup>12</sup>C ratio as measured on the AMS (Santos et al., 2007) and are reported in the F<sup>14</sup>C notation (Reimer et al., 2004). F<sup>14</sup>C is identical to the A<sub>SN</sub>/A<sub>ON</sub> metric (Stuiver and Polach, 1977), and the <sup>14</sup>a<sub>N</sub> notation (Mook and van der Plicht, 1999). Corresponding conventional <sup>14</sup>C ages reported in <sup>14</sup>C years Before Present (AD 1950) were calculated according to:

$$^{14}\text{C} = -8033\ln(\text{F}^{14}\text{C}) \quad (3)$$

## 2.3 Stable carbon isotopes

115 Stable carbon and oxygen isotopic analyses of the dated samples were performed at the Pôle Spectrométrie Océan (PSO, Plouzané, France) using a MAT-253 (Thermo Scientific) stable isotope ratio mass spectrometer (IRMS) coupled with a Kiel IV Carbonate Device (Thermo Scientific). The measurements are reported versus Vienna Pee Dee Belemnite standard (VPDB) defined with respect to two international carbonate standards: NBS-19 ( $\delta^{18}\text{O} = -2.20 \text{ ‰}$  and  $\delta^{13}\text{C} = +1.95 \text{ ‰}$ ) and NBS-18 ( $\delta^{18}\text{O} = -23.20 \text{ ‰}$  and  $\delta^{13}\text{C} = -5.01 \text{ ‰}$ ). The mean external reproducibilities (1 $\sigma$ ), based on repeated measurements of an in-house standard, was  $\pm 0.04 \text{ ‰}$  and  $\pm 0.02 \text{ ‰}$  for  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  values, respectively. Note that our samples integrate a seasonal variability of up 0.5 to 1‰ as shown by several investigations of growth layers in shells (e.g., Carré et al., 2005; Jones et al., 2007, 2010).

## 2.4 Marine Reservoir Age calculation

125 The marine reservoir age R of the selected shells is calculated according to equation (1) where t is the collection year as known from the museum records (Table S1 in the Supplement and section results), <sup>14</sup>C<sub>m</sub> is the measured shell <sup>14</sup>C age, and <sup>14</sup>C<sub>atm</sub> is the <sup>14</sup>C age of the atmosphere. For shells picked from the northern hemisphere, <sup>14</sup>C<sub>atm</sub> is obtained from the IntCal20 calibration

curve (Reimer et al., 2020). For shells from the southern hemisphere, we used instead the southern hemisphere calibration curve SHCal20 (Hogg et al., 2020). The uncertainty is calculated (Soulet, 2015) according to:

$$\sigma_{R(t)} = \sqrt{\sigma_{14C_m(t)}^2 + \sigma_{14C_{atm}(t)}^2} \quad (4)$$

130 Note that mean SHCal20 offset compared to IntCal20 is estimated to be  $36 \pm 27$   $^{14}C$  yrs. Thus, the R values calculated with IntCal20 or SHCal20 are essentially the same if one takes uncertainties into account.

The local marine reservoir offset  $\Delta R$  of the selected shells is calculated according to equation (2) where t is the collection year as known from the museum records (Table S1 in the Supplement and section results),  $^{14}C_m$  is the measured shell  $^{14}C$  age, and  $^{14}C_{Marine20}$  is the  $^{14}C$  age of the global marine calibration curve. The uncertainty is calculated as follows:

$$135 \quad \sigma_{\Delta R(t)} = \sqrt{\sigma_{14C_m(t)}^2 + \sigma_{14C_{Marine20}(t)}^2} \quad (5)$$

Note that Reimer and Reimer (2017) do not propagate the uncertainty of Marine20 calibration curve.

### 3 Results and Discussion

#### 3.1 Radiocarbon measurements results

The results are shown in [Table 1](#), Table S1 in the Supplement. ~~Below we provide the detailed list of the samples used in this~~  
140 ~~study.~~ We classified the samples by location with corresponding geographic coordinates, then by species. Code numbers “MNHN-IM-2022-xxxx” allows one to find the samples in the collections of the MNHN of Paris (France). Code numbers “SacA-xxxxx” are the radiocarbon laboratory number for the sample.

##### 3.1.1 Samples from Morocco

MNHN-IM-2022-4615

145 *Ostrea stentina*

Mohammedia (33.71°N, 7.37°W)

~~Articulated specimen with remains of flesh.~~

SacA-68834:  $482 \pm 18$  BP

$F^{14}C = 0.9418 \pm 0.0021$

150  $\delta^{13}C = 1.19$  ‰ VPDB

Collection date: 1921

Collector: Jacques de Lépiney

~~Museum label: Ostrea stentina Payr, Fedhala 1921, donat J de Lépiney 1939~~

155 MNHN-IM-2022-4609

Mis en forme : Français (France)

*Ostrea stentina*

El Jadida, beach (33.25°N, 8.49°W)

An isolated valve looking quite fresh.

SacA-68828: 817 ± 18 BP

160  $F^{14}C = 0.9033 \pm 0.0020$

$\delta^{13}C = -0.71 \text{‰ VPDB}$

Collection date: October 26<sup>th</sup> 1909

Collector: Louis Gentil

Museum label: Plage de Mazagan 26-octobre 1909, Maroc, Louis Gentil

165

MNHN-IM-2022-4608

*Ostrea stentina*

Lagoon of Sidi-Moussa (32.98°N, 8.75°W)

Articulated specimen with remains of flesh.

170 SacA-68827: 1058 ± 19 BP

$F^{14}C = 0.8766 \pm 0.0021$

$\delta^{13}C = -0.48 \text{‰ VPDB}$

Collection date: 1924

Collector: Jacques de Lépiney

175 Museum label: Ostrea stentina Payr., lagune de Sidi-Moussa (région de Mazagan), 1924, donat. J de Lépiney 1939

**3.1.2 Samples from Mauritania**

MNHN-IM-2022-4612

*Ostrea stentina*

Nouadhibou, Pointe Chacal (20.91°N, 17.04°W)

180 Articulated specimen.

SacA-68831: 514 ± 17 BP

$F^{14}C = 0.9381 \pm 0.0020$

$\delta^{13}C = 2.28 \text{‰ VPDB}$

Collection date: 1948

185 Collector: Roger Sourie

Museum label: Port Etienne (Pointe des Chacals) M. Sourie, 1948

- MNHN-IM-2022-4610  
*Ostrea stentina*  
190 Cansado Bay (20.88°N, 17.04°W)  
Articulated specimen with remains of flesh.  
SacA-68829: 516 ± 17 BP  
 $F^{14}C = 0.9378 \pm 0.0020$   
 $\delta^{13}C = 2.09 \text{‰ VPDB}$   
195 Collection date: 1911-1912  
Collector: Mission Gruvel  
Museum label: *Ostrea stentina* Payr. = *lacerans* Hanl., Baie de Cansado, Mission Gruvel, 1911-1912
- MNHN-IM-2022-4599  
200 *Bucardium ringens*  
Nouadhibou (20.88°N, 17.04°W)  
An isolated valve of a juvenile with remains of the hinge ligament.  
SacA-68811: 863 ± 18 BP  
 $F^{14}C = 0.8981 \pm 0.0020$   
205  $\delta^{13}C = 0.38 \text{‰ VPDB}$   
Collection date: 1908  
Collector: Mission Gruvel  
Museum label: *Cardium ringens* Gmelin; Port Etienne; 1908; Mission Gruvel
- MNHN-IM-2022-4603  
210 *Donax rugosus*  
Ndiago, beach (16.17°N, 16.51°W)  
Articulated specimen with remains of flesh and hinge ligament.  
SacA-68815: 518 ± 19 BP  
215  $F^{14}C = 0.9376 \pm 0.0022$   
 $\delta^{13}C = 0.68 \text{‰ VPDB}$   
Collection date: January 21<sup>st</sup> 1908  
Collector: Mission Gruvel  
Museum label: *Donax rugosus* Linné, N'Diago plage, 21.1.08, Mission Gruvel

220 **3.1.3 Samples from Senegal**

MNHN-IM-2022-4607

*Donax rugosus*

Saint Louis (16.02°N, 16.51°W)

Articulated specimen with remains of flesh and hinge ligament.

225 SacA-68819: 574 ± 18 BP

F<sup>14</sup>C = 0.9310 ± 0.0021

δ<sup>13</sup>C = 1.29 ‰ VPDB

Collection date: December 1901

Collector: Buchet's mission

230 Museum label: Sénégal, Saint Louis, Coquilles; Donax, M<sup>em</sup> Buchet, X<sup>bis</sup> 1901

MNHN-IM-2022-4592

*Senilia senilis*

Dakar, backwaters of the "Marigot de Hann" (14.74°N, 17.39°W)

235 Articulated specimen with remains of flesh and well-preserved periostracum.

SacA-68824: 560 ± 17 BP

F<sup>14</sup>C = 0.9327 ± 0.0020

δ<sup>13</sup>C = -0.29 ‰ VPDB

Collection date: May 1908

240 Collector: Mission Gruvel

Museum label: Area (*Senilia*) *senilis* Linné, Marigot de Hann V. 1908, se vend sur le marché de Dakar env. 2 sous la douzaine, Mission Gruvel

Note: The Marigot of Hann seems to have been a creek more or less connected to the ocean. It was drawn on a map of Dakar in 1905 but does not exist any longer. The map can be accessed from the Gallica website managed by the Bibliothèque

245 Nationale de France: <https://gallica.bnf.fr/ark:/12148/btv1b53197802m>

MNHN-IM-2022-4593

*Senilia senilis*

Dakar, Bay of Hann, Pointe Bel Air, beach at low tide (14.71°N, 17.42°W)

250 Articulated specimen with remains of flesh and well-preserved periostracum.

SacA-68825: 544 ± 18 BP

F<sup>14</sup>C = 0.9346 ± 0.0020



- 255  $\delta^{13}\text{C} = -0.07\text{‰}$  VPDB  
Collection date: December 1<sup>st</sup> 1909  
Collector: Mission Gruvel  
Museum label: *Area (Senilia) senilis* Linné, Baie de Hann, Pointe de Bel-Air, plage à basse mer, M-Gruvel, I.XII.1909
- MNHN-IM-2022-4606  
*Donax rugosus*  
260 Dakar, Bay of Hann (14.71°N, 17.42°W)  
Articulated specimen with remains of flesh and hinge ligament.  
SacA-68818: 526 ± 18 BP  
 $F^{14}\text{C} = 0.9366 \pm 0.0022$   
 $\delta^{13}\text{C} = -0.33\text{‰}$  VPDB
- 265 Collection date: April 1908  
Collector: Mission Gruvel  
Museum label: *Donax rugosus* Linné, Baie de Hann à basse mer, IV-08, Mission-Gruvel
- MNHN-IM-2022-4598  
270 *Bucardium ringens*  
Dakar, Bay of Hann at low tide (14.71°N, 17.42°W)  
An isolated valve with remains of the hinge ligament.  
SacA-68810: 628 ± 17 BP  
 $F^{14}\text{C} = 0.9247 \pm 0.0019$
- 275  $\delta^{13}\text{C} = -0.68\text{‰}$  VPDB  
Collection date: April 1908  
Collector: Mission-Gruvel  
Museum label: *Cardium ringens* Gmelin, Baie de Hann à basse mer, IV.08, Mission-Gruvel
- 280 MNHN-IM-2022-4616  
*Ostrea stentina*  
Dakar, beach of Hann, posts of the pontoon (14.71°N, 17.42°W)  
An isolated valve with remains of flesh  
SacA-68835: 539 ± 19 BP
- 285  $F^{14}\text{C} = 0.9351 \pm 0.0022$   
 $\delta^{13}\text{C} = -1.56\text{‰}$  VPDB

Collection date: 1947

Collector: Roger Sourie

Museum label: *Ostrea stentina* Payr-Dakar (plage de Hann, piles du ponton) M Sourie 1947

290 **3.1.4 Samples from Republic of Guinea**

MNHN-IM-2022-4601

*Bucardium ringens*

Los Islands, Roume Island at low tide (9.46°N, 13.79°W)

A fresh-looking isolated valve.

295 SacA-68813: 857 ± 39 BP

$F^{14}C = 0.8988 \pm 0.0043$

$\delta^{13}C = -0.40\text{‰ VPDB}$

Collection date: December 20<sup>th</sup> 1909

Collector: Mission Gruvel

300 Museum label: *Cardium ringens* Gmelin; Ile Roumé, archipel de Los, à basse mer, 20.XII.09, Mission Gruvel

MNHN-IM-2022-4618

*Pseudochama gryphina*

Los Islands, Tamara Island (9.46°N, 13.83°W)

305 An articulated specimen.

SacA-68820: 502 ± 19 BP

$F^{14}C = 0.9395 \pm 0.0022$

$\delta^{13}C = 1.52\text{‰ VPDB}$

Collection date: 1909-1910

310 Collector: Mission Gruvel

Museum label: *Chama gryphina* Lm, Tamara, Guinée, mission Gruvel, 1909-1910

Note: It is possible that this sample was also collected in December 1909 as sample MNHN-IM-2022-4601

**3.1.5 Sample from Sierra Leone**

MNHN-IM-2022-4611

315 *Ostrea stentina*

Near Cape Saint Ann (7.56°N, 12.94°W)

Articulated specimen with remains of flesh.

SacA-68830: 464 ± 18 BP

- 320  $F^{14}C = 0.9439 \pm 0.0021$   
 $\delta^{13}C = 1.04 \text{‰ VPDB}$   
Collection date: 1912  
Collector: Mission Gruvel  
Museum label: Sierra Léone près Cap Ste Anne, m. Gruvel, 1912
- 3.1.6 Samples from Benin**
- 325 MNHN-IM-2022-4591  
*Senilia senilis*  
Ahémé Lake (6.42°N, 1.96°E)  
Articulated specimen with well-preserved periostracum  
SacA-68823: 414 ± 18 BP
- 330  $F^{14}C = 0.9498 \pm 0.0021$   
 $\delta^{13}C = 4.76 \text{‰ VPDB}$   
Collection date: February 1910  
Collector: Mission Gruvel  
Museum label: Arca (*Senilia*) *senilis* Linné, Lac Ahémé-Dahomey, Mission Gruvel, II.1910
- 335 MNHN-IM-2022-4600  
*Bucardium ringens*  
Cotonou, dredging at a water depth of 20-25 meters (6.33°N, 2.39°E)  
A fresh isolated valve of a juvenile.
- 340 SacA-68812: 606 ± 18 BP  
 $F^{14}C = 0.9273 \pm 0.0020$   
 $\delta^{13}C = 0.26 \text{‰ VPDB}$   
Collection date: February 1910  
Collector: Mission Gruvel
- 345 Museum label: *Cardium ringens*, Cotonou, mer, II.1910, sac 372, Mission Gruvel  
Note: The information that the sample came from a dredging at 20-25 meters water depth in front of Cotonou can be found in Dautzenberg (1912).
- 350 MNHN-IM-2022-4602  
*Bucardium ringens*  
La Bouche du Roi, Grand Popo, beach (6.29°N, 1.92°E)

An isolated valve with remains of the hinge ligament.

SacA-68814: 527 ± 17 BP

$F^{14}C = 0.9365 \pm 0.0020$

355  $\delta^{13}C = -0.25\text{‰ VPDB}$

Collection date: February 1910

Collector: Mission Gruvel

Museum label: *Cardium ringens* Gmelin, Bouche du Roi, Gd Popo, plage, II.1910, Mission Gruvel.

360 Note: Three labels mention the same location, but one label mentions the Catumbella estuary (Angola, June 17th 1910). We believe the sample is from Grand Popo.

### 3.1.7 Samples from Ivory Coast

MNHN-IM-2022-4595

*Bucardium ringens*

Grand Bassam, beach (5.19°N, 3.73°W)

365 An isolated valve with remains of the hinge ligament.

SacA-68807: 507 ± 18 BP

$F^{14}C = 0.9388 \pm 0.0021$

$\delta^{13}C = -0.29\text{‰ VPDB}$

Collection date: 1909–1910

370 Collector: Mission Gruvel

Museum label: *Cardium ringens* Gmelin, plage de Gd Bassam 1909–10, mission Gruvel.

MNHN-IM-2022-4597

*Bucardium ringens*

375 Jaqueville, beach (5.19°N, 4.42°W)

An isolated valve of a juvenile with remains of the hinge ligament.

SacA-68809: 995 ± 18 BP

$F^{14}C = 0.8835 \pm 0.0020$

$\delta^{13}C = -0.03\text{‰ VPDB}$

380 Collection date: January 10<sup>th</sup> 1910

Collector: Mission Gruvel

Museum label: *Cardium ringens* Gmel., Jaqueville Côte d'Ivoire, plage, 19.I.10, Mission Gruvel.

### 3.1.8 Sample from Gabon

MNHN-IM-2022-4613

385 *Ostrea stentina*

Port-Gentil (0.71°S, 8.79°E)

An isolated valve with remains of flesh.

SacA-68832: 497 ± 19 BP

$F^{14}C = 0.9401 \pm 0.0022$

390  $\delta^{13}C = -2.24 \text{‰ VPDB}$

Collection date: 1948

Collector: Charles Roux' mission

Museum label: Port Gentil, M.Roux, 1949.

395 Note: Charles Roux writes in 1949 (Roux, 1949) that he was in the Port-Gentil area during the year 1948. We can understand that he was already back to France in 1949. Hence, the collection date must be 1948.

### 3.1.9 Samples from Republic of Congo

MNHN-IM-2022-4614

*Ostrea stentina*

Loango (4.66°S, 11.80°E)

400 An isolated valve with remains of flesh.

SacA-68833: 571 ± 19 BP

$F^{14}C = 0.9314 \pm 0.0022$

$\delta^{13}C = -1.38 \text{‰ VPDB}$

Collection date: 1890

405 Collector: Augusto Nobre

Museum label: Ostrea stentina Payr Loango M. Nobre 1890

MNHN-IM-2022-4604

*Donax rugosus*

410 Pointe-Noire (4.76°S, 11.84°E)

A fresh isolated valve from a juvenile specimen.

SacA-68816: 447 ± 18 BP

$F^{14}C = 0.9459 \pm 0.0021$

$\delta^{13}C = -0.84 \text{‰ VPDB}$

415 Collection date: December 1936—April 1937  
Collector: Edgard Aubert de la Rüe  
Museum label: Pte Noire, Aubert de la Rüe, 1937  
Note: Edgard Aubert de la Rüe was in Congo from 18/12/1936 to 16/04/1937 as evidences by his field books kept in the archives of the Musée du Quai Branly (files 2AP/62 to 2AP/64)

420 **3.1.10 Samples from Angola**

MNHN IM 2022-4590

*Senilia senilis*

Cabinda (5.55°S, 12.20°E)

Articulated specimen with well preserved periostracum.

425 SacA 68822: 584 ± 17 BP

$F^{14}C = 0.9298 \pm 0.0020$

$\delta^{13}C = -1.15\text{‰ VPDB}$

Collection date: 1885-1887

Collector: Paul Hesse

430 Museum label: Cabinda, Cabinda, Angola; C.R. Boettger coll. 1909

Note: The shell was donated by Caesar R. Boettger to the MNHN in 1909 (Oliver and von Cosel, 1992) but collected earlier by Paul Hesse when Hesse was leaving in Banana (Democratic Republic of Congo) south of Cabinda (Boettger, 1912). Boettger (1912, p. 110) writes that Hesse's collection includes a number of *Senilia senilis* specimens from Cabinda. The collection date is unfortunately not provided. However, Hesse was employed by a trading company in Banana by the end/beginning 1884/1885 since at least after March 1886 (Westhoff, 1886). Mollusc specimens reported in Boettger (1912) were collected by Hesse between 1885 and 1886. Also, Hesse collected reptile specimens in Cabinda in 1885 and 1887 (Boettger, 1898). Thus, we believe that the MNHN specimen must have been collected between 1885 and 1887.

MNHN IM 2022-4594

440 *Senilia senilis*

Cabinda (5.55°S, 12.20°E)

An isolated valve with well preserved periostracum.

SacA 68826: 536 ± 19 BP

$F^{14}C = 0.9355 \pm 0.0022$

445  $\delta^{13}C = -3.11\text{‰ VPDB}$

Collection date: June 6<sup>th</sup> 1921

Collector: Unknown

Museum label [Staat collection]: *Area senilis* Lin., Cabenda, Africa, Guinea, bought just over 1d on June 6th 1921 in Grays Jun rd. (some of the specimens at B. M. are more than double the size of mine).

450

MNHN IM 2022 4589

*Senilia senilis*

Luanda, beach (8.78°S, 13.27°E)

Articulated specimen with well-preserved periostracum.

455

SacA-68821: 538 ± 19 BP

$F^{14}C = 0.9353 \pm 0.0022$

$\delta^{13}C = -0.16 \text{‰ VPDB}$

Collection date: May 18<sup>th</sup> 1910

Collector: Mission Gruvel

460

Museum label: *Area (Senilia) senilis* Linné, St Paul de Loanda, plage, Mission Gruvel, 18.V.1910.

MNHN IM 2022 4605

*Donax rugosus*

Luanda, beach (8.82°S, 13.21°E)

465

Articulated specimen with remains of flesh and hinge ligament.

SacA-68817: 528 ± 18 BP

$F^{14}C = 0.9364 \pm 0.0021$

$\delta^{13}C = -0.82 \text{‰ VPDB}$

Collection date: May 18<sup>th</sup> 1910

470

Collector: Mission Gruvel

Museum label: *Donax rugosus* Linné, St Paul de Loanda plage, 18.V.10, Mission Gruvel.

MNHN IM 2022 4596

*Bucardium ringens*

475

Bay of Lobito, near the peninsula (12.33°S, 13.56°E)

An isolated valve.

SacA-68808: 595 ± 17 BP

$F^{14}C = 0.9286 \pm 0.0020$

$\delta^{13}C = -1.51 \text{‰ VPDB}$

480

Collection date: June 1910

Collector: Mission Gruvel

Mis en forme : Français (France)

Museum label: *Cardium ringens* Gmelin, Baie de Lobito côté presqu'île, VI.1910, mission Gruvel.

MNHN-IM 2022-4617

485 *Ostrea stentina*

Moçâmedes (15.18°S, 12.14°E)

A fresh isolated valve.

SacA 68836: 568 ± 18 BP

F<sup>14</sup>C = 0.9317 ± 0.0021

490 δ<sup>13</sup>C = -1.75 ‰ VPDB

Collection date: 1910

Collector: Mission Gruvel

Museum label: Mossamédès, m Gruvel, 1910

### 3.2 West African marine reservoir ages

495 The vast majority of the calculated ΔR values, with a weighted mean average value of  $-77-72 \pm 47-42$  <sup>14</sup>C yrs (1sd, n = 2524), which corresponds to an averaged weighted mean R value of  $400-406 \pm 59-56$  <sup>14</sup>C yrs (1sd, n = 2524), are typical of low latitudes marine reservoir age values (Bard, 1988; Bard et al., 1994) (Table S1 in the Supplement, Fig. 1). Note that all averaged R and ΔR values were calculated according to the methodology recommended in the Marine Reservoir Correction Database (Reimer and Reimer, 2001; <http://calib.org/marine/AverageDeltaR.html>; last seen 31/05/2023). Our results agree perfectly  
500 with those already obtained (Ndeye, 2008) from the Nouadhibou-Cansado Bay area (Mauritania; Nh in Fig. 1) and the Dakar area (Senegal; Dk in Fig. 1); the only two areas that we can compare our results to.

No significant interspecific differences were observed. This is best illustrated for the localities where reservoir age values were obtained from at least two different species for the same calendar time. In the Dakar area (Senegal; Dk in Fig. 1) for years 1908-1909 AD, we present data for 5 species (*Bucardium ringens*, *Donax rugosus*, *Mactra glabrata*, *Ostrea stentina*, *Senilia senilis*) (Ndeye, 2008; this study) all clustering within a range of [413;546] ([min; max]) with an average ΔR value of  $-16-18 \pm 56$  <sup>14</sup>C yrs (1sd, n = 6) (corresponding to an average R value of  $478-465 \pm 55$  <sup>14</sup>C yrs). This was also the case for Luanda (Angola; Lu in Fig. 1) in the year 1910 AD, with two species (*Donax rugosus* and *Senilia senilis*) yielding the same reservoir age values. This was further supported for the area of Nouadhibou-Cansado Bay (Mauritania) showing the same pattern (Ndeye, 2008; this study), although one sample over-out of four was likely an outlier (*Bucardium ringens* with # MNHN-IM-  
510 2022-4599). The fact that species living in very different ecological habitats (e.g., *Senilia senilis* in lagoons/semi-enclosed bays and *Donax rugosus* on beaches exposed to heavy surf; see also section material) show similar reservoir age values (R or ΔR) suggests that the habitat only exerts a minor influence on measured reservoir age in this region. The fact that all investigated species in this study correspond to suspension feeders further implies that suspension feeders are suitable material for reservoir age reconstruction.



515 Unlike semi-isolated basins such as the [Baltic Sea \(Lougheed et al., 2013\)](#) and Black Sea (Soulet et al., 2019), where the radiocarbon system is closely linked to the local [oxygen/carbon stable isotopic system respectively](#), the open-ocean coastal region of West Africa is characterized by the lack of any relationship between reservoir age values (R or  $\Delta R$ ) and stable oxygen and carbon isotope compositions ( $r^2$  of 0.02 and 0.001, respectively), as inferred from our results.

### 3.3 Marine reservoir evolution over time

520 The local marine reservoir age were averaged over 5-yr windows ([1886-1890 AD]-[1891-1895 AD] and so on), excluding the five outliers discussed in section 3.5. Sample with radiocarbon lab # AA-70015 (see Table S1 in the Supplement) is a single value from 1916 AD and was averaged with samples from years 1912 AD. We also calculated global marine reservoir age as the difference between the Marine20 and IntCal20 calibration curves. The evolution of the marine reservoir age of coastal West Africa (pink symbols in Fig. 2) shows a similar trend as that of the global marine reservoir age (black line in Fig. 2) with values declining steadily with time since ca. 1900 AD.

525 The  $^{14}\text{C}$  age evolution of the global ocean (Marine20 calibration curve; Heaton et al., 2020) is constructed using the global carbon cycle model BICYCLE (Köhler et al., 2006; Köhler and Fischer, 2006, 2004; Köhler et al., 2005). This box model incorporates a globally averaged atmospheric box and modules of the terrestrial (7 boxes) and oceanic (10 boxes) components of the carbon cycle. It is driven by temporal changes in the boundary conditions mimicking changing climate and simulates changes in the carbon cycle including  $^{14}\text{C}$ . To construct the Marine20 calibration curve, the BICYCLE model was revised to allow the atmospheric  $\text{CO}_2$  and  $\text{F}^{14}\text{C}$  to be specified externally (Heaton et al., 2020). [While the modelled Marine20 \(global surface ocean\) radiocarbon age suggest constant values between 1900 and 1950 AD, our measured marine reservoir age R indicates instead a decreasing trend during that period, as a consequence of increasing atmospheric Intcal20 radiocarbon age. This observation of decreasing trend for R in West Africa between 1900 and 1950 AD could possibly reflect atmospheric  \$^{14}\text{CO}\_2\$  ageing following enhanced fossil fuel emissions to the atmosphere by burning \(e.g., Suess, 1955; Tans et al., 1979\)](#) The global marine reservoir age evolution calculated as the difference between Marine20 and Intcal20 calibration curves (Fig.1) is thus also an output of the revised BICYCLE model. The fact that the trend in our measured marine reservoir age is similar to that of the globally modelled ones strongly suggest that global carbon cycle and climate are the first order drivers of the coastal West African marine reservoir age rather than local effects.

### 540 3.4 Marine reservoir age off equatorial Ogooué and Congo rivers

Large rivers draining equatorial Africa as the Ogooué and Congo rivers inject massive amounts of freshwater into the Atlantic Ocean (Lambert et al., 2015; Milliman and Farnsworth, 2011) leading to extensive sea surface salinity negative anomalies (Martins and Stammer, 2022). The sea surface salinity negative anomalies are associated with net primary productivity positive anomalies that are likely caused by the nutrient-rich river plumes from the Ogooué and Congo Rivers (Martins and Stammer, 2022). From a radiocarbon perspective, such net primary productivity positive anomalies should imply an increased uptake of atmospheric  $\text{CO}_2$  through intensified biological pump. As a result, the reservoir age should be lower than average. The Congo

River represents the second largest supplier of dissolved organic carbon (DOC) to [the](#) global ocean with ~5% of the land to ocean DOC flux (Spencer et al., 2016; Coynel et al., 2005; Richey et al., 2022). The DOC exported by the Congo river is <sup>14</sup>C-modern (Marwick et al., 2015; Spencer et al., 2012) and experiments showed that 45% of the Congo River DOC can be photo-mineralized by sunlight (Spencer et al., 2009; Richey et al., 2022). Dissolved inorganic carbon (DIC) released from the photo-mineralisation of the Congo River DOC should also be <sup>14</sup>C-modern. Thus, this modern DOC-derived DIC should impact the marine reservoir age towards values lower compared to average. There is a lack of available data to estimate the age and flux of dissolved CO<sub>2</sub> discharged by the Congo river into the ocean (Richey et al., 2022). Nevertheless, the marine reservoir age value measured at Port-Gentil (Gabon) close to the Ogooué river outlet is lower than [the regional weighted mean valueaverage value](#) ( $\Delta R = -106 \pm 63$  <sup>14</sup>C years, corresponding to  $R = 329 \pm 21$  <sup>14</sup>C yrs) (PG in Fig. 1). The marine reservoir age measured in Pointe-Noire (Republic of Congo) ~150 km north of the Congo river outlet is also lower than [the regional weighted mean value-average](#) ( $\Delta R = -156 \pm 64$  <sup>14</sup>C yrs;  $R = 289 \pm 20$  <sup>14</sup>C yrs) (PN in Fig. 1). These values could be interpreted as having been influenced by the Ogooué and Congo Rivers discharges. However, all other localities close to the Congo River outlet had marine reservoir age close to the [average-regional weighted mean value](#) (Lo, Ca and Lu, in Fig. 1). Instead the lower values observed in Port-Gentil (Gabon) and Pointe-Noire (Republic of Congo) are from years 1948 and 1937 suggesting that these lower values are in line with the declining global marine reservoir evolution linked to [the atmospheric CO<sub>2</sub> <sup>14</sup>C ageing linked to <sup>14</sup>C-dead input from fossil fuel burning \(Suess effect\) global-elimite-and-carbon-eyele-changes](#)-(see section 3.3). The impact of the African equatorial rivers on the local/regional coastal marine reservoir age, if any, cannot be inferred from our results.

### 3.5 Outlier specimens

Mean marine reservoir age values (R and  $\Delta R$ ) are provided for West Africa based on our data, excluding 5 samples. These particular samples display much larger values with  $\Delta R$  values ranging from 209 to 454 <sup>14</sup>C yrs or R values ranging from 701 to 912 <sup>14</sup>C yrs. Three specimens out of the five outlier samples correspond to *Bucardium ringens* specimens. We analysed 8 *Bucardium ringens* specimens. These 3 outlier specimens display reservoir age (R and  $\Delta R$ ) values that clearly disagree with neighbouring data (Nouadhibou-Cansado Bay, Loos Islands and Ivory Coast areas; Nh, LI and IC in Fig. 1). The Museum number of these specimens are MNHN-IM-2022-4597, MNHN-IM-2022-4599 and MNHN-IM-2022-4601. We do not expect that these larger values compared to those for neighbouring individuals come from the species feeding practice as they are all suspension feeders like all other investigated specimens. Similarly, we showed that the difference in the habitat [in this region](#) does not impact the [speeiespecies](#) reservoir ages. Instead *Bucardium ringens* lives in the open coast from 5-10 meters to about 50 meters depth. Shells are commonly cast ashore on beaches but live-taken specimens are rare (von Cosel and Gofas, 2019). [Two-One of these the samples that are outliers were-was](#) collected at low tide (Roume Island in the Loos Islands; Republic of Guinea)[-or on a beach \(Jacqueville; Ivory Coast\) and was devoid of any remain of flesh or hinge ligament](#). It is thus possible that [these-this](#) outlier samples [were-was a](#) transported subfossil samples that died a century or more before collection date. [The two other outlier samples had small remain of the hinge ligament \(Nouadhibou; Mauritania and Jacqueville; Ivory Coast\). It may be possible that these samples are also subfossil specimens. In this case, the hinge ligament must have been partially](#)

580 preserved owing to very favourable environmental conditions (Forman et al., 2004; Huntley et al., 2021). Alternatively, these  
outliers are not subfossil specimens and unlike the other studied species here, the habitat may exert an influence on R and  $\Delta R$   
values measured in *B. ringens*. Nevertheless, Finally, we cannot fully rule out that these higher values represent some sub-  
585 annual variability of up to 200  $^{14}\text{C}$  in the local marine reservoir age as evidenced elsewhere (Jones et al., 2007, 2010). Although,  
five *Bucardium ringens* samples out of eight displayed reservoir age values in agreement with the neighbouring reservoir age  
values, this specie might not be the best suited for reservoir age reconstruction or for sediment/archaeological dating.  
The two remaining outliers are *Ostrea stentina* specimens from the El Jadida area (Morocco; eJ in Fig. 1). The sample from  
El Jadida beach was a single valve looking fresh and collected from the beach (museum # MNHN-IM-2022-4609). Based on  
the older  $^{14}\text{C}$  age of this specimen, we cannot rule out that this sample could actually be a subfossil specimen. The specimen  
with museum # MNHN-IM-2022-4608 collected in the Sidi Moussa lagoon (south of El Jadida) was a specimen with the  
590 articulated valves and remains of flesh was still inside the shell, meaning the specimen was still alive when collected. Variations  
in the reservoir age could be explained by coastal upwelling that impacts some regions of the Atlantic coast of Morocco and  
Western Sahara (Freudenthal et al., 2001; Barton et al., 1998). Upwelled waters can be depleted in  $^{14}\text{C}$  relative to the sea  
surface potentially causing larger reservoir age values (R or  $\Delta R$ ) like off Portugal (Monge Soares, 1993; Monge Soares and  
Alveirinho Dias, 2006), California (Kennett et al., 1997), Peru (Kennett et al., 2002; Fontugne et al., 2004; Jones et al., 2007,  
595 2010) or Southern Arabian coast (Southon et al., 2002). Conversely, upwelled waters can also be nutrient-rich causing  
intensified ocean  $\text{CO}_2$  uptake through enhanced primary production and biological pump (Williams and Follows, 2011), in  
that case, one could expect low-latitude average or decreased reservoir age values (R or  $\Delta R$ ). Off Morocco and Western Sahara,  
the second hypothesis appears most likely as coastal upwelling in this area is known to bring nutrient-rich waters to the surface  
ocean (Barton et al., 1998; Freudenthal et al., 2001), although to our knowledge no direct measurement of the  $^{14}\text{C}$  content of  
600 coastal waters in this region has been published yet. However, according to recent studies the El Jadida area is only weakly  
impacted by upwelling (Lourenço et al., 2020; Cropper et al., 2014), suggesting average reservoir age values instead of larger  
ones. Another explanation could be linked to the local hydrology of the Sidi Moussa lagoon. Despite the lagoon is being  
permanently connected to the ocean, it receives waters from rainfall and resurgences that can have an impact on the salinity in  
the upstream section of the lagoon (Cheggour et al., 2001). As the surrounding rocks are calcareous sandstones (Manaan,  
605 2003), one could hypothesise that freshwaters feeding the lagoon might be depleted in  $^{14}\text{C}$  due to carbonate dissolution in the  
lagoon watershed causing a hardwater effect and thus a larger reservoir age. A last explanation could be due to an imperfect  
cleaning of the shell. For *Ostrea stentina*, sediment can be trapped between the growing layers of the shell. If this sediment  
contains old detrital carbonates and was not perfectly removed before  $^{14}\text{C}$  measurement, the  $^{14}\text{C}$  age of the shell will appear  
older, and the reservoir age larger. Additional reservoir age reconstructions from this region on different species would be  
610 require to validate the larger reservoir age values reconstructed from the El Jadida area.

Code de champ modifié

Mis en forme : Anglais (Royaume-Uni)

Mis en forme : Anglais (Royaume-Uni)

Code de champ modifié

#### 4 Conclusion

The analysis of pre-bomb suspension-feeding bivalves collected along coastal West Africa from 33°N to 15°S provides marine reservoir ages that are quite homogenous, with a mean  $\Delta R$  value of  $-77-72 \pm 47-42$   $^{14}\text{C}$  yrs (1sd, n = 2524) and a mean R value of  $400-406 \pm 59-56$   $^{14}\text{C}$  yrs (1sd, n = 2524). When including the robust dataset from Ndeye (2008), the resulting mean  $\Delta R$  and R values for coastal West Africa are  $-62-54 \pm 51$   $^{14}\text{C}$  years (1sd, n = 3332) and  $416-411 \pm 61$   $^{14}\text{C}$  years (1sd, n = 3332), respectively. We show that the marine reservoir age of coastal West Africa is mainly driven by the global carbon cycle and climate-atmospheric  $^{14}\text{C}$  changes rather than by local effects.

Our results for different species yield similar marine reservoir age values, indicating that the ecological habitat only has a second-order impact on the reservoir age reconstruction, if any. Nevertheless, we suspect that *Bucardium ringens* might not be best suited for marine reservoir age reconstruction as corresponding shells are typically not found alive on sample collecting sites. Additionally, ages obtained on *Ostrea stentina* could be possibly influenced by the presence of sediment within the growing shell layers if not fully removed after the cleaning process.

Despite these new data, large portions of the West African coast still remain to be investigated for reservoir age reconstructions, in particular off Western Sahara and Canarias Islands, Sierra Leone-Liberia, Nigeria and Namibia.

#### Author contribution statement

GSoulet designed the study and raised the funding. SG, PM, GSoulet and GSiani selected the specimens in the MNHN collections. GSoulet carried out the sample preparation with assistance of ML and FF. FD performed stable isotopes measurements. GSoulet performed reservoir calculations and analysed and discussed the data with GB, GSiani and SG. GSoulet wrote the manuscript with inputs from all co-authors.

#### Acknowledgements

All authors thank Michel Fontugne, an anonymous reviewer and Paula Reimer for reviewing the manuscript. We are grateful to the Muséum National d'Histoire Naturelle (Paris, France) for providing the samples. We thank the LMC14 staff (Laboratoire de Mesure du Carbone-14) and the CNRS-INSU ARTEMIS national radiocarbon AMS facility for providing radiocarbon measurements published in this study. The study was funded by the French national program INSU-LEFE (ResWA project – Pre-bomb Reservoir ages for the Western coast of Africa). I (GS) thank Ifremer in Brest and Sète for supporting my research. Finally, this study was carried out and the paper written during my part-time parental leave. I dedicate this article to Marian, with whom I have had a wonderful time during this year, growing up together, him as a toddler and me as a father.

### Competing interests

The authors declare that they have no conflict of interest.

### 640 Data availability

All data present in the paper are available in the text section 3.1 and in Table S1 in the Supplement.

### References

- Alves, E. Q., Macario, K., Ascough, P., and Bronk Ramsey, C.: The worldwide marine radiocarbon reservoir effect: Definitions, mechanisms, and prospects, *Rev. Geophys.*, 56, 278–305, <https://doi.org/10.1002/2017RG000588>, 2018.
- 645 Alves, E. Q., Macario, K. D., Urrutia, F. P., Cardoso, R. P., and Bronk Ramsey, C.: Accounting for the marine reservoir effect in radiocarbon calibration, *Quat. Sci. Rev.*, 209, 129–138, <https://doi.org/10.1016/j.quascirev.2019.02.013>, 2019.
- [Arnold, J. R. and Anderson, E. C.: The distribution of Carbon-14 in nature, \*Tellus\*, 9, 28–32, <https://doi.org/10.1111/j.2153-3490.1957.tb01850.x>, 1957.](https://doi.org/10.1111/j.2153-3490.1957.tb01850.x)
- Ascough, P., Cook, G., and Dugmore, A.: Methodological approaches to determining the marine radiocarbon reservoir effect, *Prog. Phys. Geogr. Earth Environ.*, 29, 532–547, <https://doi.org/10.1191/0309133305pp461ra>, 2005.
- 650 Bard, E.: Correction of accelerator mass spectrometry <sup>14</sup>C ages measured in planktonic foraminifera: Paleooceanographic implications, *Paleoceanography*, 3, 635–645, <https://doi.org/10.1029/PA003i006p00635>, 1988.
- Bard, E., Arnold, M., Mangerud, J., Paterne, M., Labeyrie, L., Duprat, J., Mélières, M.-A., Sønstegeard, E., and Duplessy, J.-C.: The North Atlantic atmosphere-sea surface <sup>14</sup>C gradient during the Younger Dryas climatic event, *Earth Planet. Sci. Lett.*, 126, 275–287, [https://doi.org/10.1016/0012-821X\(94\)90112-0](https://doi.org/10.1016/0012-821X(94)90112-0), 1994.
- 655 Barton, E. D., Aristegui, J., Tett, P., Cantón, M., García-Braun, J., Hernández-León, S., Nykjaer, L., Almeida, C., Almunia, J., Ballesteros, S., Basterretxea, G., Escánez, J., García-Weill, L., Hernández-Guerra, A., López-Laatzén, F., Molina, R., Montero, M. F., Navarro-Pérez, E., Rodríguez, J. M., van Lenning, K., Vélez, H., and Wild, K.: The transition zone of the Canary Current upwelling region, *Prog. Oceanogr.*, 41, 455–504, [https://doi.org/10.1016/S0079-6611\(98\)00023-8](https://doi.org/10.1016/S0079-6611(98)00023-8), 1998.
- 660 Boettger, C. R.: Zur Molluskenfauna des Kongogebiets, *Ann. la Société R. Zool. Malacol. Belgique*, 47, 89–117, 1912.
- Boettger, O.: Katalog der Reptilien-Sammlung im Museum der Senckenbergischen Naturforschenden Gesellschaft in Frankfurt am Main, II Teil (Schlangen), Gebrüder Knauer, Frankfurt am Main, 1898.
- Butzin, M., Köhler, P., and Lohmann, G.: Marine radiocarbon reservoir age simulations for the past 50,000 years, *Geophys. Res. Lett.*, 44, 8473–8480, <https://doi.org/10.1002/2017GL074688>, 2017.
- 665 [Carré, M., Bentaleb, I., Blamart, D., Ogle, N., Cardenas, F., Zevallos, S., Kalin, R. M., Ortlieb, L., and Fontugne, M.: Stable isotopes and sclerochronology of the bivalve \*Mesodesma donacium\*: Potential application to Peruvian paleoceanographic reconstructions, \*Palaeogeogr. Palaeoclimatol. Palaeoecol.\*, 228, 4–25, <https://doi.org/10.1016/j.palaeo.2005.03.045>, 2005.](https://doi.org/10.1016/j.palaeo.2005.03.045)

- Catry, T., Figueira, P., Carvalho, L., Monteiro, R., Coelho, P., Lourenço, P. M., Catry, P., Tchanchalam, Q., Catry, I., Botelho, M. J., Pereira, E., Granadeiro, J. P., and Vale, C.: Evidence for contrasting accumulation pattern of cadmium in relation to other elements in *Senilia senilis* and *Tagelus adansonii* from the Bijagós archipelago, Guinea-Bissau, *Environ. Sci. Pollut. Res.*, 24, 24896–24906, <https://doi.org/10.1007/s11356-017-9902-8>, 2017.
- Cheggour, M., Chafik, A., Langston, W. ., Burt, G. ., Benbrahim, S., and Texier, H.: Metals in sediments and the edible cockle *Cerastoderma edule* from two Moroccan Atlantic lagoons: Moulay Bou Selham and Sidi Moussa, *Environ. Pollut.*, 115, 149–160, [https://doi.org/10.1016/S0269-7491\(01\)00117-8](https://doi.org/10.1016/S0269-7491(01)00117-8), 2001.
- von Cosel, R. and Gofas, S.: *Marine Bivalves of Tropical West Africa: From Rio de Oro to Southern Angola*, edited by: MNHN and IRD, *Faune et Flore tropicales*, 48, Paris, Marseille, 1104 pp., 2019.
- Cottéreau, E., Arnold, M., Moreau, C., Baqué, D., Bavay, D., Caffy, I., Comby, C., Dumoulin, J.-P., Hain, S., Perron, M., Salomon, J., and Setti, V.: Artemis, the new <sup>14</sup>C AMS at LMC14 in Saclay, France, *Radiocarbon*, 49, 291–299, <https://doi.org/10.1017/S0033822200042211>, 2007.
- Coynel, A., Seyler, P., Etcheber, H., Meybeck, M., and Orange, D.: Spatial and seasonal dynamics of total suspended sediment and organic carbon species in the Congo River, *Global Biogeochem. Cycles*, 19, n/a-n/a, <https://doi.org/10.1029/2004GB002335>, 2005.
- Craig, H.: The natural distribution of radiocarbon and the exchange time of carbon dioxide between atmosphere and sea, *Tellus*, 9, 1–17, <https://doi.org/10.1111/j.2153-3490.1957.tb01848.x>, 1957.
- Cropper, T. E., Hanna, E., and Bigg, G. R.: Spatial and temporal seasonal trends in coastal upwelling off Northwest Africa, 1981–2012, *Deep Sea Res. Part I Oceanogr. Res. Pap.*, 86, 94–111, <https://doi.org/10.1016/j.dsr.2014.01.007>, 2014.
- Dautzenberg, P.: Mission Gruvel sur la côte occidentale d'Afrique (1909-1910): Mollusques marins. *Annales de l'Institut Océanographique*, Paris (Nouvelle Série), 5(3): 1-111, pl. 1-3, 1912.
- Dewar, G., Reimer, P. J., Sealy, J., and Woodborne, S.: Late-Holocene marine radiocarbon reservoir correction (ΔR) for the west coast of South Africa, *The Holocene*, 22, 1481–1489, <https://doi.org/10.1177/0959683612449755>, 2012.
- Dumoulin, J.-P., Comby-Zerbino, C., Delqué-Količ, E., Moreau, C., Caffy, I., Hain, S., Perron, M., Thellier, B., Setti, V., Berthier, B., and Beck, L.: Status report on sample preparation protocols developed at the LMC14 laboratory, Saclay, France: From Sample Collection to <sup>14</sup>C AMS Measurement, *Radiocarbon*, 59, 713–726, <https://doi.org/10.1017/RDC.2016.116>, 2017.
- Fontugne, M., Carré, M., Bentaieb, I., Julien, M., and Lavallée, D.: Radiocarbon reservoir age variations in the south Peruvian upwelling during the Holocene, *Radiocarbon*, 46, 531–537, <https://doi.org/10.1017/S003382220003558X>, 2004.
- Forman, S.: A review of postglacial emergence on Svalbard, Franz Josef Land and Novaya Zemlya, northern Eurasia, *Quat. Sci. Rev.*, 23, 1391–1434, <https://doi.org/10.1016/j.quascirev.2003.12.007>, 2004.
- Freudenthal, T., Neuer, S., Meggers, H., Davenport, R., and Wefer, G.: Influence of lateral particle advection and organic matter degradation on sediment accumulation and stable nitrogen isotope ratios along a productivity gradient in the Canary Islands region, *Mar. Geol.*, 177, 93–109, [https://doi.org/10.1016/S0025-3227\(01\)00126-8](https://doi.org/10.1016/S0025-3227(01)00126-8), 2001.

- Heaton, T. J., Köhler, P., Butzin, M., Bard, E., Reimer, R. W., Austin, W. E. N., Bronk Ramsey, C., Grootes, P. M., Hughen, K. A., Kromer, B., Reimer, P. J., Adkins, J., Burke, A., Cook, M. S., Olsen, J., and Skinner, L. C.: Marine20—The Marine Radiocarbon Age Calibration Curve (0–55,000 cal BP), *Radiocarbon*, 62, 779–820, <https://doi.org/10.1017/RDC.2020.68>,  
705 2020.
- Heaton, T. J., Bard, E., Bronk Ramsey, C., Butzin, M., Köhler, P., Muscheler, R., Reimer, P. J., and Wacker, L.: Radiocarbon: A key tracer for studying Earth’s dynamo, climate system, carbon cycle, and Sun, *Science* (80-. ), 374, <https://doi.org/10.1126/science.abd7096>, 2021.
- Heaton, T. J., Bard, E., Bronk Ramsey, C., Butzin, M., Hatté, C., Hughen, K. A., Köhler, P., and Reimer, P. J.: A response to community questions on the Marine20 radiocarbon age calibration curve: Marine reservoir ages and the calibration of <sup>14</sup>C samples from the oceans, *Radiocarbon*, 65, 247–273, <https://doi.org/10.1017/RDC.2022.66>, 2023.
- Herrera, N. D., ter Poorten, J. J., Bieler, R., Mikkelsen, P. M., Strong, E. E., Jablonski, D., and Stepan, S. J.: Molecular phylogenetics and historical biogeography amid shifting continents in the cockles and giant clams (*Bivalvia: Cardiidae*), *Mol. Phylogenet. Evol.*, 93, 94–106, <https://doi.org/10.1016/j.ympev.2015.07.013>, 2015.
- 715 Hogg, A. G., Heaton, T. J., Hua, Q., Palmer, J. G., Turney, C. S., Southon, J., Bayliss, A., Blackwell, P. G., Boswijk, G., Bronk Ramsey, C., Pearson, C., Petchey, F., Reimer, P., Reimer, R., and Wacker, L.: SHCal20 Southern Hemisphere Calibration, 0–55,000 Years cal BP, *Radiocarbon*, 62, 759–778, <https://doi.org/10.1017/RDC.2020.59>, 2020.
- [Huntley, J. W., De Baets, K., Scarponi, D., Linehan, L. C., Epa, Y. R., Jacobs, G. S., and Todd, J. A.: Bivalve mollusks as hosts in the fossil record, in: \*The Evolution and Fossil Record of Parasitism\*, edited by: De Baets, K. and Huntley, J. W., Springer International Publishing, 251–287, \[https://doi.org/10.1007/978-3-030-52233-9\\\_8\]\(https://doi.org/10.1007/978-3-030-52233-9\_8\), 2021.](#)
- 720 Kennett, D. J., Ingram, B. L., Erlandson, J. M., and Walker, P.: Evidence for Temporal Fluctuations in Marine Radiocarbon Reservoir Ages in the Santa Barbara Channel, Southern California, *J. Archaeol. Sci.*, 24, 1051–1059, <https://doi.org/10.1006/jasc.1996.0184>, 1997.
- [Kennett, D. J., Lynn Ingram, B., Southon, J. R., and Wise, K.: Differences in <sup>14</sup>C age between stratigraphically associated charcoal and marine shell from the Archaic Period Site of Kilometer 4, Southern Peru: Old wood or old water?, \*Radiocarbon\*, 44, 53–58, <https://doi.org/10.1017/S0033822200064663>, 2002.](#)
- 725 Köhler, P. and Fischer, H.: Simulating changes in the terrestrial biosphere during the last glacial/interglacial transition, *Glob. Planet. Change*, 43, 33–55, <https://doi.org/10.1016/j.gloplacha.2004.02.005>, 2004.
- Köhler, P. and Fischer, H.: Simulating low frequency changes in atmospheric CO<sub>2</sub> during the last 740 000 years, *Clim. Past*, 2, 57–78, <https://doi.org/10.5194/cp-2-57-2006>, 2006.
- 730 Köhler, P., Fischer, H., Munhoven, G., and Zeebe, R. E.: Quantitative interpretation of atmospheric carbon records over the last glacial termination, *Global Biogeochem. Cycles*, 19, n/a-n/a, <https://doi.org/10.1029/2004GB002345>, 2005.
- Köhler, P., Muscheler, R., and Fischer, H.: A model-based interpretation of low-frequency changes in the carbon cycle during the last 120,000 years and its implications for the reconstruction of atmospheric Δ<sup>14</sup>C, *Geochemistry, Geophys. Geosystems*, 7, n/a-n/a, <https://doi.org/10.1029/2005GC001228>, 2006.
- 735

Lambert, T., Darchambeau, F., Bouillon, S., Alhou, B., Mbega, J.-D., Teodoru, C. R., Nyoni, F. C., Massicotte, P., and Borges, A. V.: Landscape control on the spatial and temporal variability of chromophoric dissolved organic matter and dissolved organic carbon in large African Rivers, *Ecosystems*, 18, 1224–1239, <https://doi.org/10.1007/s10021-015-9894-5>, 2015.

Lindsay, C. M., Lehman, S. J., Marchitto, T. M., Carriquiry, J. D., and Ortiz, J. D.: New constraints on deglacial marine radiocarbon anomalies from a depth transect near Baja California, *Paleoceanography*, 31, 1103–1116, <https://doi.org/10.1002/2015PA002878>, 2016.

[Lougheed, B. C., Filipsson, H. L., and Snowball, I.: Large spatial variations in coastal <sup>14</sup>C reservoir age – a case study from the Baltic Sea, \*Clim. Past\*, 9, 1015–1028, <https://doi.org/10.5194/cp-9-1015-2013>, 2013.](#)

745 Lourenço, C. R., Nicastro, K. R., McQuaid, C. D., Krug, L. A., and Zardi, G. I.: Strong upwelling conditions drive differences in species abundance and community composition along the Atlantic coasts of Morocco and Western Sahara, *Mar. Biodivers.*, 50, 15, <https://doi.org/10.1007/s12526-019-01032-z>, 2020.

Manaán, M.: Étude sédimentologique du remplissage de la lagune de Sidi Moussa, Maroc, PhD thesis, Université Chouaib Doukkali, El Jadida, <https://theses.hal.science/tel-00124571>, 2003.

Martins, M. S. and Stammer, D.: Interannual variability of the Congo River plume-induced sea surface salinity, *Remote Sens.*, 750 14, 1013, <https://doi.org/10.3390/rs14041013>, 2022.

Marwick, T. R., Tamooh, F., Teodoru, C. R., Borges, A. V., Darchambeau, F., and Bouillon, S.: The age of river-transported carbon: A global perspective, *Global Biogeochem. Cycles*, 29, 122–137, <https://doi.org/10.1002/2014GB004911>, 2015.

Milliman, J. D. and Farnsworth, K. L.: River discharge to the coastal ocean, Cambridge University Press, <https://doi.org/10.1017/CBO9780511781247>, 2011.

755 Monge Soares, A.: The <sup>14</sup>C content of marine shells: Evidence for variability in coastal upwelling off Portugal during the Holocene, in: *Isotope techniques in the study of past and current environmental changes in the hydrosphere and the atmosphere proceedings*, edited by: IAEA, Vienna, 471–485, 1993.

Monge Soares, A. M. and Alveirinho Dias, J. M.: Coastal upwelling and radiocarbon-evidence for temporal fluctuations in ocean reservoir effect off Portugal during the holocene, *Radiocarbon*, 48, 45–60, <https://doi.org/10.1017/S0033822200035384>, 760 2006.

Mook, W. G. and van der Plicht, J.: Reporting <sup>14</sup>C activities and concentrations, *Radiocarbon*, 41, 227–239, <https://doi.org/10.1017/S0033822200057106>, 1999.

Moreau, C., Caffy, I., Comby, C., Delqué-Količ, E., Dumoulin, J.-P., Hain, S., Quiles, A., Setti, V., Souprayen, C., Thellier, B., and Vincent, J.: Research and development of the Artemis <sup>14</sup>C AMS facility: Status report, *Radiocarbon*, 55, 331–337, 765 <https://doi.org/10.1017/S0033822200057441>, 2013.

Ndeye, M.: Marine reservoir ages in Northern Senegal and Mauritania coastal waters, *Radiocarbon*, 50, 281–288, <https://doi.org/10.1017/S0033822200033580>, 2008.

Okera, W.: Observations on some population parameters of exploited stocks of *Senilia senilis* (= *Arca senilis*) in Sierra Leone, *Mar. Biol.*, 38, 217–229, <https://doi.org/10.1007/BF00388935>, 1976.



- 770 Oliver, P. G. and von Cosel, R.: Taxonomy of tropical west African Bivalves. IV. Arcidae., *Bull. Muséum Natl. d'Histoire Nat.*, 14, 293–381, 1992.
- Reimer, P. J. and McCormac, F. G.: Marine radiocarbon reservoir corrections for the Mediterranean and Aegean Seas, *Radiocarbon*, 44, 159–166, <https://doi.org/10.1017/S0033822200064766>, 2002.
- Reimer, P. J. and Reimer, R. W.: A marine reservoir correction database and on-line interface, *Radiocarbon*, 43, 461–463, <https://doi.org/10.1017/S0033822200038339>, 2001.
- 775 Reimer, P. J., Brown, T. A., and Reimer, R. W.: Discussion: Reporting and calibration of post-bomb <sup>14</sup>C data, *Radiocarbon*, 46, 1299–1304, <https://doi.org/10.1017/S0033822200033154>, 2004.
- Reimer, P. J., Austin, W. E. N., Bard, E., Bayliss, A., Blackwell, P. G., Bronk Ramsey, C., Butzin, M., Cheng, H., Edwards, R. L., Friedrich, M., Grootes, P. M., Guilderson, T. P., Hajdas, I., Heaton, T. J., Hogg, A. G., Hughen, K. A., Kromer, B., 780 Manning, S. W., Muscheler, R., Palmer, J. G., Pearson, C., van der Plicht, J., Reimer, R. W., Richards, D. A., Scott, E. M., Southon, J. R., Turney, C. S. M., Wacker, L., Adolphi, F., Büntgen, U., Capano, M., Fahrni, S. M., Fogtmann-Schulz, A., Friedrich, R., Köhler, P., Kudsk, S., Miyake, F., Olsen, J., Reinig, F., Sakamoto, M., Sookdeo, A., and Talamo, S.: The IntCal20 northern hemisphere radiocarbon age calibration curve (0–55 cal kBP), *Radiocarbon*, 62, 725–757, <https://doi.org/10.1017/RDC.2020.41>, 2020.
- 785 Reimer, R. W. and Reimer, P. J.: An online application for  $\Delta R$  calculation, *Radiocarbon*, 59, 1623–1627, <https://doi.org/10.1017/RDC.2016.117>, 2017.
- [Revelle, R. and Suess, H. E.: Carbon dioxide exchange between atmosphere and ocean and the question of an increase of atmospheric CO<sub>2</sub> during the past decades. \*Tellus\*, 9, 18–27, <https://doi.org/10.1111/j.2153-3490.1957.tb01849.x>, 1957.](https://doi.org/10.1111/j.2153-3490.1957.tb01849.x)
- Richey, J. E., Spencer, R. G. M., Drake, T. W., and Ward, N. D.: Fluvial carbon dynamics across the land to ocean continuum of great tropical rivers, 391–412, <https://doi.org/10.1002/9781119657002.ch20>, 2022.
- 790 Roux, C.: Compte rendu sommaire d'une mission en Afrique Équatoriale Française, *Bull. du Muséum Natl. d'Histoire Nat.*, 21, 500–503, 1949.
- Santos, G. M., Southon, J. R., Griffin, S., Beaupre, S. R., and Druffel, E. R. M.: Ultra small-mass AMS <sup>14</sup>C sample preparation and analyses at KCCAMS/UCI Facility, *Nucl. Instruments Methods Phys. Res. Sect. B Beam Interact. with Mater. Atoms*, 259, 293–302, <https://doi.org/10.1016/j.nimb.2007.01.172>, 2007.
- 795 Schefuß, E., Eglinton, T. I., Spencer-Jones, C. L., Rullkötter, J., De Pol-Holz, R., Talbot, H. M., Grootes, P. M., and Schneider, R. R.: Hydrologic control of carbon cycling and aged carbon discharge in the Congo River basin, *Nat. Geosci.*, 9, 687–690, <https://doi.org/10.1038/ngeo2778>, 2016.
- Sessa, J. A., Callapez, P. M., Dinis, P. A., and Hendy, A. J. W.: Paleoenvironmental and paleobiogeographical implications of a middle Pleistocene mollusc assemblage from the marine terraces of Baía Das Pipas, southwest Angola, *J. Paleontol.*, 87, 1016–1040, <https://doi.org/10.1666/12-119>, 2013.
- 800 Siani, G., Paterne, M., Arnold, M., Bard, E., Métyvier, B., Tisnerat, N., and Bassinot, F.: Radiocarbon reservoir ages in the Mediterranean Sea and Black Sea, *Radiocarbon*, 42, 271–280, <https://doi.org/10.1017/S0033822200059075>, 2000.

- 805 Siani, G., Paterne, M., Michel, E., Sulpizio, R., Sbrana, A., Arnold, M., and Haddad, G.: Mediterranean Sea surface  
radiocarbon reservoir age changes since the Last Glacial Maximum, *Science* (80-. ), 294, 1917–1920,  
<https://doi.org/10.1126/science.1063649>, 2001.
- [Siegenthaler, U.: Uptake of excess CO<sub>2</sub> by an outcrop-diffusion model of the ocean, \*J. Geophys. Res.\*, 88, 3599,  
https://doi.org/10.1029/JC088iC06p03599, 1983.](https://doi.org/10.1029/JC088iC06p03599)
- 810 Skinner, L., McCave, I. N., Carter, L., Fallon, S., Scrivner, A. E., and Primeau, F.: Reduced ventilation and enhanced  
magnitude of the deep Pacific carbon pool during the last glacial period, *Earth Planet. Sci. Lett.*, 411, 45–52,  
<https://doi.org/10.1016/j.epsl.2014.11.024>, 2015.
- Skinner, L. C. and Bard, E.: Radiocarbon as a dating tool and tracer in paleoceanography, *Rev. Geophys.*, 60, 1–64,  
<https://doi.org/10.1029/2020RG000720>, 2022.
- 815 Skinner, L. C., Fallon, S., Waelbroeck, C., Michel, E., and Barker, S.: Ventilation of the deep Southern Ocean and deglacial  
CO<sub>2</sub> rise, *Science* (80-. ), 328, 1147–1151, <https://doi.org/10.1126/science.1183627>, 2010.
- Smith, D. A. S.: Polymorphism and population density in *Donax rugosus* (Lamellibranchiata: Donacidae), *J. Zool.*, 164, 429–  
441, <https://doi.org/10.1111/j.1469-7998.1971.tb01327.x>, 1971.
- Soulet, G.: Methods and codes for reservoir–atmosphere <sup>14</sup>C age offset calculations, *Quat. Geochronol.*, 29, 97–103,  
<https://doi.org/10.1016/j.quageo.2015.05.023>, 2015.
- 820 Soulet, G., Ménot, G., Garreta, V., Rostek, F., Zaragosi, S., Lericolais, G., and Bard, E.: Black Sea “Lake” reservoir age  
evolution since the Last Glacial — Hydrologic and climatic implications, *Earth Planet. Sci. Lett.*, 308, 245–258,  
<https://doi.org/10.1016/j.epsl.2011.06.002>, 2011.
- Soulet, G., Skinner, L. C., Beaupré, S. R., and Galy, V.: A note on reporting of reservoir <sup>14</sup>C disequilibria and age offsets,  
*Radiocarbon*, 58, 205–211, <https://doi.org/10.1017/RDC.2015.22>, 2016.
- 825 Soulet, G., Giosan, L., Flaux, C., and Galy, V.: Using stable carbon isotopes to quantify radiocarbon reservoir age offsets in  
the coastal black sea, *Radiocarbon*, 61, <https://doi.org/10.1017/RDC.2018.61>, 2019.
- Southon, J., Kashgarian, M., Fontugne, M., Metivier, B., and Yim, W. W. S.: Marine reservoir corrections for the Indian Ocean  
and Southeast Asia, *Radiocarbon*, 44, 167–180, <https://doi.org/10.1017/S0033822200064778>, 2002.
- Spencer, R. G. M., Stubbins, A., Hernes, P. J., Baker, A., Mopper, K., Aufdenkampe, A. K., Dyda, R. Y., Mwamba, V. L.,  
830 Mangangu, A. M., Wabakanghanzi, J. N., and Six, J.: Photochemical degradation of dissolved organic matter and dissolved  
lignin phenols from the Congo River, *J. Geophys. Res.*, 114, G03010, <https://doi.org/10.1029/2009JG000968>, 2009.
- Spencer, R. G. M., Hernes, P. J., Aufdenkampe, A. K., Baker, A., Gulliver, P., Stubbins, A., Aiken, G. R., Dyda, R. Y., Butler,  
K. D., Mwamba, V. L., Mangangu, A. M., Wabakanghanzi, J. N., and Six, J.: An initial investigation into the organic matter  
biogeochemistry of the Congo River, *Geochim. Cosmochim. Acta*, 84, 614–627, <https://doi.org/10.1016/j.gca.2012.01.013>,  
835 2012.
- Spencer, R. G. M., Hernes, P. J., Dinga, B., Wabakanghanzi, J. N., Drake, T. W., and Six, J.: Origins, seasonality, and fluxes  
of organic matter in the Congo River, *Global Biogeochem. Cycles*, 30, 1105–1121, <https://doi.org/10.1002/2016GB005427>,

2016.

Stuiver, M. and Braziunas, T. F.: Modeling atmospheric  $^{14}\text{C}$  influences and  $^{14}\text{C}$  ages of marine samples to 10,000 BC,

840 Radiocarbon, 35, 137–189, <https://doi.org/10.1017/S0033822200013874>, 1993.

Stuiver, M. and Polach, H. A.: Discussion: Reporting of  $^{14}\text{C}$  Data, Radiocarbon, 19, 355–363, <https://doi.org/10.1017/S0033822200003672>, 1977.

Stuiver, M., Pearson, G. W., and Braziunas, T.: Radiocarbon age calibration of marine samples back to 9000 cal yr BP, Radiocarbon, 28, 980–1021, <https://doi.org/10.1017/S0033822200060264>, 1986.

845 [Suess, H. E.: Radiocarbon concentration in modern wood, Science, 122, 415–417, https://doi.org/10.1126/science.122.3166.415.b, 1955.](https://doi.org/10.1126/science.122.3166.415.b)

[Tans, P. P., De Jong, A. F. M., and Mook, W. G.: Natural atmospheric  \$^{14}\text{C}\$  variation and the Suess effect, Nature, 280, 826–828, https://doi.org/10.1038/280826a0, 1979.](https://doi.org/10.1038/280826a0)

Tisnérat-Laborde, N., Poupeau, J. J., Tannau, J. F., and Paterne, M.: Development of a semi-automated system for routine preparation of carbonate samples, Radiocarbon, 43, 299–304, <https://doi.org/10.1017/S0033822200038145>, 2001.

850

Türkmen, A., Türkmen, M., and Tepe, Y.: Biomonitoring of heavy metals from Iskenderun Bay using two bivalve species *Chama pacifica* Broderip, 1834 and *Ostrea stentina* Payraudeau, 1826, Turkish J. Fish. Aquat. Sci., 5, 107–111, 2005.

Westhoff, F.: Die Fauna der Kongo-Mündung, Jahresbericht der Zool. Sekt. Westfälischen Prov. Vereins für Wiss. und Kunst für das Etatsjahr 1885-86, 38–40, 1886.

855 Williams, R. G. and Follows, M. J.: Ocean dynamics and the carbon cycle: Principles and mechanisms, Cambridge., Cambridge, 434 pp., 2011.






Figure 1: The geographic distribution of marine reservoir age values along the West African coast. A.  $\Delta R$  values. B. R values. Data shown in black are from this study. Other are selected results from previous studies discussed in the text, converted from their original format (conventional 14C ages and collection dates) to  $\Delta R$  and R values using the latest calibration curves Marine20 (Heaton et al., 2020) and Intcal20 or SHcal20 (Reimer et al., 2020; Hogg et al., 2020), respectively. Data in blue are from Ndeye (2008), data in green are from (Reimer and McCormac, 2002), data in purple are from (Siani et al., 2000) and data in pink are from (Dewar et al., 2012). eJ, Nh, Dk, LI, IC, PG, Lo, PN, Ca and Lu stand for el Jadida (Morocco), Nouadhibou (Mauritania), Dakar (Senegal), Loos Islands (Republic of Guinea), Ivory Coast, Port-Gentil (Gabon), Loango (Republic of Congo), Pointe Noire (Republic of Congo), Cabinda (Angola) and Luanda (Angola). The map was drawn using Ocean Data View (Schlitzer, Reiner, Ocean Data View, <https://odv.awi.de>, 2022).







870 Figure 2: **Left axis A:** The radiocarbon age evolution of the atmosphere (IntCal20; green curve with its light green 1- $\sigma$  envelope) and of the global ocean (Marine20; blue curve with its light blue 1- $\sigma$  envelope) between 1850 and 1950 AD. **Right axis B:** The global marine reservoir age (black curve with its grey 1- $\sigma$  envelope) calculated as the difference between Marine20 and Intcal20 curves. Pink symbols are the coastal West African marine reservoir age calculated averaging data over 5-yr windows. The reported error bars are the maximum of the standard deviation of the averaged data and the individual uncertainty of the averaged data.

875

Table 1 see version without tracks.

Mis en forme : Normal